

#### Home

2. Pyridium

# Pyridium 🖘

**Generic name:** phenazopyridine [ fen-AY-zoe-PIR-i-deen ]

Brand names: AZO Urinary Pain Relief, Azo-Gesic, Azo-Standard, Baridium, Prodium, ... show all 8 brands

Drug class: Miscellaneous genitourinary tract agents

Medically reviewed by Sanjai Sinha, MD. Last updated on Jul 15, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

## What is Pyridium?

Pyridium is a pain reliever that affects the lower part of your urinary tract (bladder and urethra).

Pyridium is used to treat urinary symptoms such as pain or burning, increased urination, and increased urge to urinate. These symptoms can be caused by infection, injury, surgery, catheter, or other conditions that irritate the bladder.

Pyridium will treat urinary symptoms, but **this medication will not treat a urinary tract infection.** Take any antibiotic that your doctor prescribes to treat an infection.

# **Warnings**

Do not take Pyridium if you are allergic to phenazopyridine, or if you have kidney disease. Pyridium will treat the symptoms of a urinary tract infection, but this medication does not treat the actual infection. Take any antibiotic that your doctor prescribes to treat your infection. To avoid stomach upset, take this medicine with food.

Pyridium will most likely darken the color of your urine to an orange or red color. This is a normal effect and is not cause for alarm unless you have other symptoms such as pale or yellowed skin, fever, stomach pain, nausea, and vomiting. Darkened urine may also cause stains to your underwear, which may or may not be removed by laundering.

Pyridium can also permanently stain soft contact lenses, and you should not wear them while taking this medicine.

Do not use Pyridium for longer than 2 days unless your doctor has told you to.

Stop taking this medicine and call your doctor at once if you have pale skin, fever, confusion, yellowing of your skin or eyes, increased thirst, swelling, or if you urinate less than usual or not at all.

# Before taking this medicine

To make sure Pyridium is safe for you, tell your doctor if you have ever had:

• liver disease:

- · diabetes; or
- a genetic enzyme deficiency called glucose-6-phosphate dehydrogenase (G6PD) deficiency.

Pyridium is not expected to harm an unborn baby. **Do not use this medicine without a doctor's advice if you are pregnant.** 

It is not known whether phenazopyridine passes into breast milk or if it could harm a nursing baby. **Do not use this** medicine without a doctor's advice if you are breast-feeding a baby.

Pyridium pregnancy and breastfeeding warnings (more detail)

# How should I take Pyridium?

Use Pyridium exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended.

Take Pyridium after meals.

Drink plenty of liquids while you are taking Pyridium.

Pyridium will most likely darken the color of your urine to an orange or red color. This is a normal effect and is not harmful. Darkened urine may also cause stains to your underwear that may be permanent.

Phenazopyridine can also permanently stain soft contact lenses, and you should not wear them while taking this medicine.

Do not use this medicine for longer than 2 days unless your doctor has told you to.

This medication can cause unusual results with urine tests. Tell any doctor who treats you that you are using this medicine.

Store at room temperature away from moisture and heat.

Detailed Pyridium dosage information

# What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

# What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

### What to avoid

Do not use this medication while wearing soft contact lenses. Phenazopyridine can permanently discolor soft contact lenses.

# Pyridium side effects

Get emergency medical help if you have any of these **signs of an allergic reaction to Pyridium:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using Pyridium and call your doctor at once if you have:

- little or no urinating;
- swelling, rapid weight gain;
- confusion, loss of appetite, pain in your side or lower back;
- fever, pale or yellowed skin, stomach pain, nausea and vomiting; or
- blue or purple appearance of your skin.

Common Pyridium side effects may include:

- headache;
- dizziness; or
- · upset stomach.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Pyridium side effects (more detail)

# Related/similar drugs

### **Ditropan**

Ditropan XL (oxybutynin extended release) is used to treat the symptoms of overactive bladder ...

#### **Reviews & ratings**

4.7 / 10

34 Reviews

View more

#### FEATURED

#### **Omvoh**

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

#### **Reviews & ratings**

1 Reviews

View more

### **Oxytrol**

Oxytrol patches reduces muscle spasms in the urinary tract and are used to treat symptoms of ...

#### **Reviews & ratings**

7.4 / 10

79 Reviews

View more

# What other drugs will affect Pyridium?

Other drugs may interact with phenazopyridine, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

1 Pyridium drug interactions (more detail)

# Does Pyridium interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

**Pyridium** 

Enter a drug name Add

# Frequently asked questions

• Why do you not take phenazopyridine for more than two days?

# More about Pyridium (phenazopyridine)

- Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (35)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: miscellaneous genitourinary tract agents
- Breastfeeding
- En español

### **Patient resources**

### Other brands

Azo Urinary Pain Relief, Azo-Standard, Uristat, Prodium, ... +5 more

### **Professional resources**

- Phenazopyridine monograph
- Phenazopyridine Tablets (FDA)

### Related treatment guides

- Interstitial Cystitis
- Dysuria

### **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Pyridium only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

#### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

### **DRUG STATUS**

### **Availability**

Rx and/or OTC

**Pregnancy & Lactation** 

ം Risk data available

**CSA Schedule\*** 

N/A Not a controlled drug

### **User Reviews & Ratings**

7.3 / 10

35 Reviews

### **Images**

### Pyridium 100 mg (P-D 180)





View larger images

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com
Advertising policy
Attribution & citations

### Terms & privacy

Terms of use Editorial policy Privacy policy

#### **Support**

Help center

Sitemap

Contact us














<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







 $\times$ 

Copyright © 2000-2025 Drugs.com. All rights reserved.