

Home

Cinacalcet

Cinacalcet 49

Generic name: cinacalcet [sin-ah-CAL-set]

Brand name: Sensipar

Dosage form: oral tablet (30 mg; 60 mg; 90 mg)

Drug class: Calcimimetics



Medically reviewed by Drugs.com on Jul 9, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is cinacalcet?

Cinacalcet decreases levels of parathyroid hormone (PTH), calcium, and phosphorous in the body.

Cinacalcet is used to treat hyperparathyroidism (overactive functioning of the parathyroid glands) in people who are on long-term dialysis for kidney disease.

Cinacalcet is also used to treat hypercalcemia (high levels of calcium in the blood) in people with cancer of the parathyroid gland, and in people with overactive parathyroid glands that cannot be treated surgically.

Cinacalcet may also be used for purposes not listed in this medication guide.

Cinacalcet side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Cinacalcet may cause serious side effects. Call your doctor at once if you have:

- numbness or tingly feeling around your mouth;
- muscle pain, tightness, or contraction;
- a seizure (convulsions);
- shortness of breath (even with mild exertion), swelling, rapid weight gain;
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds; or
- low levels of calcium in your blood--numbness or tingly feeling around your mouth, fast or slow heart rate, muscle tightness or contraction, overactive reflexes.

Common side effects of cinacalcet may include:

- nausea, vomiting; or
- diarrhea.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

• Cinacalcet side effects (more detail)

Related/similar drugs

Prolia

Prolia (denosumab) is an injection that is administered subcutaneously (under the skin) once every ...

Reviews & ratings

2.9 / 10

374 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Reclast

Reclast (zoledronic acid) is used to treat osteoporosis in women after menopause and also to treat ...

Reviews & ratings

5.1 / 10

275 Reviews

View more

Zometa

Zometa is used to treat high blood levels of calcium caused by cancer

and multiple myeloma. Learn ...

Reviews & ratings

4.8 / 10

43 Reviews

View more

Xgeva

Xgeva (denosumab) prevents bone fractures and other skeletal conditions in people with multiple ...

Reviews & ratings

6.1 / 10

9 Reviews

View more

Sensipar

Sensipar reduces elevated levels of parathyroid hormone in people with chronic renal disease who ...

Reviews & ratings

5.7 / 10

9 Reviews

View more

Zemplar

Zemplar (paricalcitol) is used to treat secondary hyperparathyroidism in people with chronic kidney ...

Reviews & ratings

Add a review

View more

Warnings

You should not take cinacalcet if you have low levels of calcium in your blood (hypocalcemia).

Before taking this medicine

You should not use cinacalcet if you are allergic to it, or if you have:

low levels of calcium in your blood (hypocalcemia).

To make sure cinacalcet is safe for you, tell your doctor if you have ever had:

- high or low blood pressure;
- heart disease, heart failure, heart rhythm disorder;
- personal or family history of long QT syndrome;
- a stomach ulcer or severe vomiting;
- liver disease;
- · seizures; or
- if you have kidney disease and you are **not** on dialysis.

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether cinacalcet passes into breast milk or if it could harm a nursing baby. You should not breast-feed while you are taking this medicine.

Cinacalcet is not approved for use by anyone younger than 18 years old.

How should I take cinacalcet?

Follow all directions on your prescription label. Your doctor may occasionally change your dose to make sure you get the best results. Do not use cinacalcet in larger or smaller amounts or for longer than recommended.

Cinacalcet works best if you take it with food or shortly after eating a meal.

Do not crush or break a cinacalcet tablet. Swallow the pill whole.

To be sure your calcium and phosphorous levels do not get too low, your blood will need to be tested 1 week after you start taking cinacalcet or whenever your dose is changed. You may also need blood tests on a regular basis during treatment.

Store at room temperature away from moisture, light, and heat.

Detailed Cinacalcet dosage information

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. **Do not** take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking cinacalcet?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

What other drugs will affect cinacalcet?

Many drugs can interact with cinacalcet. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide. Tell your doctor about all your current medicines and any medicine you start or stop using.

① Cinacalcet drug interactions (more detail)

Does cinacalcet interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

| cinacalcet | |
|-------------------|-----|
| + | |
| Enter a drug name | Add |

More about cinacalcet

- · Check interactions
- · Compare alternatives
- · Pricing & coupons
- Reviews (10)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- · Drug class: calcimimetics
- Breastfeeding
- En español

Patient resources

Other brands

Sensipar

Professional resources

Cinacalcet monograph

Cinacalcet (FDA)

Other brands

Sensipar

Related treatment guides

- Primary Hyperparathyroidism
- Secondary Hyperparathyroidism
- Hypercalcemia of Malignancy

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 7.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ঝ্য Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

5.7 / 10

10 Reviews

Images

Cinacalcet Hydrochloride 30 mg (H C6)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.