



[Home](#)

2. [Condylox](#)

Condylox

Generic name: [podofilox topical](#) [*poe-DOF-il-lox*]

Drug class: [Topical keratolytics](#)

[Medically reviewed](#) by Drugs.com. Last updated on Jul 10, 2024.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#)

What is Condylox?

Condylox prevents the growth of certain types of cells.

Condylox is used to treat [genital warts](#) that appear on the outside of the body, but this medicine is not a cure for genital warts.

Condylox may also be used for purposes not listed in this medication guide.

Warnings

Do not use Condylox on any warts or skin growths that have not been checked by your doctor.

Using more Condylox will not make it work better, and may increase side effects. Avoid getting this medication in your eyes. If this does happen, rinse with water and call your doctor. Do not place the gel or solution in your rectum, vagina, or urethra. If it does get into any of these areas, rinse with water.

Avoid having unprotected sex. Condylox is not a cure for genital warts, and it will not protect against sexually transmitted diseases such as chlamydia, gonorrhea, herpes, HIV, syphilis, and trichomoniasis. Talk with your doctor about safe ways to prevent transmission during sex.

Wash your hands before and after applying Condylox.

Before taking this medicine

You should not use Condylox if you are allergic to podofilox.

Do not use this medicine on any warts or skin growths that have not been checked by your doctor.

It is not known whether Condylox will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medicine.

It is not known whether podofilox topical passes into breast milk or if it could harm a nursing baby. You should not breast-feed while using this medicine.

Condylox is not approved for use by anyone younger than 18 years old.

 [Condylox pregnancy and breastfeeding warnings](#) (more detail)

How should I use Condylox?

Use Condylox exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not use this medicine in larger or smaller amounts or for longer than recommended.

Your healthcare provider may show you how to apply this medicine at home.

Using more of this medicine will not make it work better, and may increase side effects.

This medicine is usually applied in the morning and evening for 3 days in a row, followed by 4 days of not using the medicine. This 7-day treatment cycle should be repeated up to 4 times, or until you can no longer see any signs of wart tissue.

Wash your hands before applying Condylox. Also wash the skin area to be treated, and allow it to dry completely before applying the medicine.

Apply Condylox using the cotton applicator provided or with your finger. Apply the medicine only to the wart tissue and avoid getting it on healthy skin. If the genital wart is in a skin fold, spread the skin apart and apply the medicine. Allow the treated skin to dry before releasing the skin fold.

Wash your hands again after applying Condylox.

Call your doctor if your symptoms do not improve after 4 treatment cycles.

Condylox is not a cure for genital warts and it may not keep you from spreading this condition to others through vaginal, anal, or oral sex. You may develop new lesions during treatment with Condylox. For best results, keep using the medicine for the entire length of time prescribed by your doctor.

Store at room temperature away from moisture and heat. Do not freeze.

Condylox gel is flammable. Do not use it near high heat or open flame, or while you are smoking.

Dosing information

Usual Adult Dose for Condylomata Acuminata:

Solution or gel: Apply topically twice daily for 3 consecutive days, then discontinue for 4 consecutive days.

Duration: The cycle may be repeated until there is no visible wart tissue or for a maximum of four cycles.

 [Detailed Condylox dosage information](#)

What happens if I miss a dose?

Use the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not use extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may result if you absorb this medicine through your skin by applying too much of it. Symptoms may include nausea, vomiting, diarrhea, fever, flu symptoms, sores in your mouth and throat, numbness or tingling in your hands or feet, confusion, trouble breathing, or seizure (convulsions).

What should I avoid while using Condyllox?

Do not have sex on the days you apply Condyllox.

Avoid getting this medicine in your eyes. If this does happen, rinse with water and call your doctor. Do not place the gel or solution in your rectum, vagina, or urethra. If it does get into any of these areas, rinse with water.

Avoid having unprotected sex. Condyllox is not a cure for genital warts, and it will not protect against sexually transmitted diseases such as chlamydia, gonorrhea, herpes, HIV, syphilis, and trichomoniasis. Talk with your doctor about safe ways to prevent transmission of disease during sex.

Condyllox side effects

Get emergency medical help if you have **signs of an allergic reaction to Condyllox**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Wash off the medicine and call your doctor at once if you have a serious skin reaction such as severe itching, burning, oozing, bleeding, or skin changes where the medicine is applied.

Common Condyllox side effects may include:

- mild skin irritation (pain, swelling, burning, redness, itching, tingling, blisters, crusting, dryness, peeling) where the medicine was applied;
- headache, dizziness;
- sleep problems (insomnia);
- unpleasant skin odor;
- vomiting; or
- pain during sex.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Condyllox side effects](#) (more detail)

Related/similar drugs

Aldara

Aldara (imiquimod) is used to treat
actinic keratosis, basal cell carcinoma

actinic keratoses, basal cell carcinoma
and genital warts ...

Reviews & ratings

6.9 / 10

[135 Reviews](#)

[View more](#)

Veregen

Veregen is used to treat external
genital and anal warts in adult patients.
Learn about side ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide)
for type 2 diabetes treatment, weight
management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Zyclara

Zyclara (imiquimod) is used to treat
actinic keratoses on the face or
balding scalp and external ...

Reviews & ratings

6.8 / 10

[49 Reviews](#)

[View more](#)

Sinecatechins topical

Polyphenon E information from
Drugs.com, includes Polyphenon E
side effects interactions, indications

Reviews & ratings

[1 Reviews](#)

[View more](#)

What other drugs will affect Condyllox?

It is not likely that other drugs you take orally or inject will have an effect on topically applied podofilox. But many drugs can interact with each other. Tell each of your health care providers about all medicines you use, including prescription and over-the-counter medicines, vitamins, and herbal products.

Does Condyllox interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Condyllox

+

Enter a drug name

Add

References

1. [Condyllox Product Label](#)

More about Condyllox (podofilox topical)

- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(9\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [Drug class: topical keratolytics](#)
- [En español](#)

Professional resources

- [Condyllox prescribing information](#)
- [Podofilox \(Topical\) \(AHFS Monograph\)](#)

Related treatment guides

- [Condylomata Acuminata](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Condyllox only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 2.01.

DRUG STATUS

Availability

Rx Prescription only

[Pregnancy & Lactation](#)


 Risk data available

CSA Schedule*

N/A Not a controlled drug



Approval History

 Drug history at FDA



User Reviews & Ratings

5.9 / 10

[9 Reviews](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.