

Home

2. Hydrochlorothiazide/valsartan

Hydrochlorothiazide and valsartan 🖏

Generic name: hydrochlorothiazide and valsartan [*HYE-droe-klor-oh-THYE-a-zide-and-val-SAR-tan*]

Brand name: Diovan HCT

Dosage form: oral tablet (12.5 mg-160 mg; 12.5 mg-320 mg; 12.5 mg-80 mg; 25 mg-160 mg; 25 mg-320 mg)

Drug class: Angiotensin II inhibitors with thiazides



Medically reviewed by Drugs.com on Feb 5, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is hydrochlorothiazide and valsartan?

Hydrochlorothiazide is a diuretic (water pill). Valsartan is an angiotensin II receptor blocker (sometimes called an ARB blocker).

Hydrochlorothiazide and valsartan is a combination medicine used to treat high blood pressure (hypertension). Lowering blood pressure may lower your risk of a stroke or heart attack.

Hydrochlorothiazide and valsartan is given after other blood pressure medicines did not work or have stopped working.

Hydrochlorothiazide and valsartan may also be used for purposes not listed in this medication guide.

Hydrochlorothiazide and valsartan side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

In rare cases, hydrochlorothiazide and valsartan can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine.

Hydrochlorothiazide and valsartan may cause serious side effects. Call your doctor at once if you have:

- a light-headed feeling, like you might pass out;
- · eye pain, vision problems;
- · an unusual skin rash;
- little or no urinating;
- jaundice (yellowing of the skin or eyes);
- · easy bruising, unusual bleeding;

- · a seizure;
- **kidney problems**--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath; or
- signs of an electrolyte imbalance--dry mouth, increased thirst, confusion, feeling restless, vomiting, muscle pain or weakness, drowsiness, lack of energy, fast heartbeats, decreased urination.

Common side effects include:

- · cold symptoms such as stuffy nose, sneezing, sore throat;
- · dizziness; or
- · headache.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Hydrochlorothiazide / valsartan side effects (more detail)

Related/similar drugs

Vasotec

Vasotec (enalapril) is used to treat high blood pressure, heart failure and kidney problems caused ...

Reviews & ratings

4 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Lisinopril

Lisinopril is an ACE inhibitor. It is used to treat high blood pressure, congestive heart failure ...

Reviews & ratings

5.0 / 10

626 Raviawe

View more

Atenolol

Atenolol is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

Reviews & ratings

7.2 / 10

276 Reviews

View more

Valsartan

Valsartan is an angiotensin II receptor blocker (ARB) that may be used to treat high blood pressure ...

Reviews & ratings

5.7 / 10

152 Reviews

View more

Hydrochlorothiazide

HCTZ (hydrochlorothiazide) used to treat high blood pressure (hypertension) and edema. Includes ...

Reviews & ratings

5.2 / 10

185 Reviews

View more

Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

Reviews & ratings

6.4 / 10

56 Reviews

View more

Furosemide

Furosemide is a loop diuretic used to treat fluid retention and high blood pressure by increasing ...

Reviews & ratings

6.8 / 10

121 Reviews

View more

Losartan

Losartan is used to treat high blood pressure (hypertension). It is also used to lower the risk of ...

Reviews & ratings

4.5 / 10

578 Reviews

View more

Warnings

You should not use hydrochlorothiazide and valsartan if you are unable to urinate, or if you are allergic to sulfa drugs.

Do not use if you are pregnant. Stop using the medicine and tell your doctor right away if you become pregnant.

If you have diabetes, do not use hydrochlorothiazide and valsartan together with any medication that contains aliskiren (a blood pressure medicine).

Before taking this medicine

You should not use this medicine if you are allergic to valsartan or hydrochlorothiazide, or if you have:

- · decreased urination or are unable to urinate; or
- an allergy to sulfa drugs.

If you have diabetes, do not use hydrochlorothiazide and valsartan together with any medication that contains aliskiren (a blood pressure medicine).

You may also need to avoid taking hydrochlorothiazide and valsartan with aliskiren if you have kidney disease.

Tell your doctor if you have ever had:

• an electrolyte imbalance (such as low levels of potassium or magnesium, or high levels of calcium in your blood);

- liver disease;
- · kidney disease;
- · gallstones;
- · lupus;
- congestive heart failure;
- gout, or high levels of uric acid in your blood;
- asthma or allergies;
- high cholesterol or triglyceride levels;
- an allergy to penicillin;
- · glaucoma; or
- if you are on a low-salt diet.

Do not use if you are pregnant. Stop using the medicine and tell your doctor right away if you become pregnant. Valsartan can cause injury or death to the unborn baby if you take the medicine during your second or third trimester.

You should not breastfeed while using this medicine.

Hydrochlorothiazide and valsartan is not approved for use by anyone younger than 18 years old.

How should I take hydrochlorothiazide and valsartan?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

You may take hydrochlorothiazide and valsartan with or without food.

Call your doctor if you have ongoing vomiting or diarrhea, or if you are sweating more than usual. You can easily become dehydrated while taking this medication, which can lead to severely low blood pressure or a serious electrolyte imbalance.

Your blood pressure will need to be checked often. Your blood and urine may also be tested if you have been vomiting or are dehydrated.

If you need surgery, tell the surgeon ahead of time that you are using hydrochlorothiazide and valsartan.

It may take up to 4 weeks for this medication to control your blood pressure. **Keep using this medicine as directed, even if you feel well.** High blood pressure often has no symptoms. You may need to use blood pressure medication for the rest of your life.

Store at room temperature away from moisture and heat.

Hydrochlorothiazide and valsartan dosing information

Usual Adult Dose for Hypertension:

Initial dose: 1 tablet (12.5 mg-160 mg) orally once a day

Maintenance dose: Titrate as necessary (every 1 to 2 weeks) up to a maximum of 25 mg-320 mg orally once a day

Comment:

- -Hydrochlorothiazide-valsartan may be substituted for the titrated components.
- 1 Detailed Hydrochlorothiazide / valsartan dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking hydrochlorothiazide and valsartan?

Hydrochlorothiazide may increase your risk of skin cancer. Avoid sunlight or tanning beds. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors. Your doctor may want you to have skin examinations on a regular basis.

Do not use potassium supplements or salt substitutes, unless your doctor has told you to.

Drinking alcohol can further lower your blood pressure and may cause side effects.

Avoid driving or hazardous activity until you know how hydrochlorothiazide and valsartan will affect you. Your reactions could be impaired. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

What other drugs will affect hydrochlorothiazide and valsartan?

If you also take cholestyramine or colestipol, take your hydrochlorothiazide and valsartan dose 4 hours before or 4 hours after you take the other medicine.

Tell your doctor about all your current medicines. Many drugs can affect hydrochlorothiazide and valsartan, especially:

- cyclosporine, digoxin, digitalis, lithium, or ritonavir;
- all your heart or blood pressure medicines;
- insulin or oral diabetes medicine:
- a diuretic or "water pill";
- · opioid pain medicine;
- · a sleeping pill;
- a muscle relaxer;

- an antibiotic--rifabutin, rifampin, rifamycin, rifapentine;
- cancer medicine--cyclophosphamide, methotrexate; or
- NSAIDs (nonsteroidal anti-inflammatory drugs)--aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others.

This list is not complete and many other drugs may affect hydrochlorothiazide and valsartan. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Hydrochlorothiazide / valsartan drug interactions (more detail)

Does hydrochlorothiazide/valsartan interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

hydrochlorothiazide/valsartan
+
Enter a drug name
Add

More about hydrochlorothiazide / valsartan

- · Check interactions
- · Compare alternatives
- · Pricing & coupons
- Reviews (26)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: angiotensin II inhibitors with thiazides
- En español

Patient resources

- · Valsartan and hydrochlorothiazide advanced reading
- Valsartan and Hydrochlorothiazide

Other brands

Diovan HCT

Professional resources

• Valsartan and Hydrochlorothiazide prescribing information

Other brands

Diovan HCT

Related treatment guides

• High Blood Pressure

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.


Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 14.01.

DRUG STATUS

Availability Rx Prescription only **Pregnancy & Lactation** ঝ্ Risk data available CSA Schedule* N/A Not a controlled drug **Approval History**





Anti-Doping Classification

User Reviews & Ratings

6.8 / 10

26 Reviews

Images

Hydrochlorothiazide and Valsartan 12.5 mg / 160 mg (M V22)





Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





