

Home

2. Sanctura XR

Sanctura XR

Generic name: trospium [*tros-PEE-um*] **Drug class:** Urinary antispasmodics



Medically reviewed by Drugs.com on Aug 8, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions



The Sanctura XR brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be generic equivalents available.

What is Sanctura XR?

Sanctura XR relieves spasms of the bladder.

Sanctura XR is used to treat overactive bladder and symptoms of urinary incontinence, frequency, and urgency.

Sanctura XR may also be used for purposes not listed in this medication guide.

Sanctura XR side effects

Get emergency medical help if you have any of these **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using Sanctura XR and call your doctor at once if you have a serious side effect such as:

- · severe stomach pain or bloating;
- severe constipation;
- urinating less than usual or not at all; or
- · confusion, hallucinations.

Less serious side effects may include:

- dry mouth or throat;
- headache;
- mild constipation;
- upset stomach, gas;

- · dizziness, drowsiness, or
- dry eyes.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Sanctura XR side effects (more detail)

Related/similar drugs

Botox

Botox is used to treat chronic migraines, excessive sweating, bladder conditions, eye muscle ...

Reviews & ratings

5.7 / 10

473 Reviews

View more

Myrbetriq

Myrbetriq (mirabegron) is used to treat overactive bladder with symptoms of frequent or urgent ...

Reviews & ratings

4.8 / 10

276 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Detrol

Detrol is used to treat overactive bladder symptoms of urinary

frequency, urgency, and ...

Reviews & ratings

7.5 / 10

View more

VESIcare

17 Reviews

Vesicare is used to treat symptoms of overactive bladder such as incontinence and frequent ...

Reviews & ratings

5.6 / 10

142 Reviews

View more

Ditropan

Ditropan XL (oxybutynin extended release) is used to treat the symptoms of overactive bladder ...

Reviews & ratings

4.7 / 10

34 Reviews

View more

Toviaz

Toviaz (fesoterodine) is used to treat overactive bladder with symptoms of urinary frequency ...

Reviews & ratings

5.2 / 10

74 Reviews

View more

Warnings

You should not take Sanctura XR if you are allergic to Sanctura XR, or if you have untreated or uncontrolled narrow-angle glaucoma, a blockage in your digestive system, or if you are unable to urinate.

Take Sanctura XR on an empty stomach, at least 1 hour before a meal.

Avoid drinking alcohol within 2 hours before or after you take Sanctura XR. Drinking alcohol can increase certain side effects of trospium.

This medication may cause blurred vision and may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert and able to see clearly.

Avoid becoming overheated or dehydrated during exercise and in hot weather. Sanctura XR can decrease sweating, which makes it easier for you to have heat stroke. Drink plenty of fluids while you are taking this medication.

Before using Sanctura XR, tell your doctor if you regularly use other medicines that make you sleepy (such as cold or allergy medicine, sedatives, narcotic pain medicine, sleeping pills, muscle relaxers, and medicine for seizures, depression, or anxiety). They can add to sleepiness caused by this medicine.

Before taking this medicine

You should not use Sanctura XR if you are allergic to it, or if you have:

- untreated or uncontrolled narrow-angle glaucoma;
- · a blockage in your digestive system; or
- if you are unable to urinate.

To make sure you can safely take Sanctura XR, tell your doctor if you have any of these other conditions:

- glaucoma;
- · liver disease:
- · kidney disease;
- a stomach or intestinal disorder such as ulcerative colitis;
- a muscle disorder such as myasthenia gravis; or
- an enlarged prostate.

FDA pregnancy category C. It is not known whether Sanctura XR will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medication.

It is not known whether trospium passes into breast milk or if it could harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

Older adults may be more likely to have side effects from this medicine.

How should I take Sanctura XR?

Take exactly as prescribed by your doctor. Do not take in larger or smaller amounts or for longer than recommended. Follow the directions on your prescription label.

Take Sanctura XR on an empty stomach, at least 1 hour before a meal. Extended-release this medicine (**Sanctura XR**) should be taken once each morning, at least 1 hour before a meal.

Do not crush, chew, break, or open an extended-release capsule. Swallow it whole. Breaking or opening the pill may cause too much of the drug to be released at one time.

Store at room temperature away from moisture and heat.

Detailed Sanctura XR dosage information

What happens if I miss a dose?

Take the missed dose 1 hour before your next meal. Skip the missed dose if it is almost time for your next scheduled dose. **Do not** take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe forms of some of the side effects listed in this medication guide.

What should I avoid while taking Sanctura XR?

This medication may cause blurred vision and may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert and able to see clearly.

Avoid drinking alcohol within 2 hours before or after you take Sanctura XR. Drinking alcohol can increase certain side effects of trospium.

Avoid becoming overheated or dehydrated during exercise and in hot weather. Sanctura XR can decrease sweating, which makes it easier for you to have heat stroke. Drink plenty of fluids while you are taking this medication.

What other drugs will affect Sanctura XR?

Before using Sanctura XR, tell your doctor if you regularly use other medicines that make you sleepy (such as cold or allergy medicine, sedatives, narcotic pain medicine, sleeping pills, muscle relaxers, and medicine for seizures, depression, or anxiety). They can add to sleepiness caused by this medicine.

Tell your doctor about all other medicines you use, especially:

- atropine (Atreza, Sal-Tropine, and others);
- belladonna (Donnatal, and others);
- benztropine (Cogentin);
- dimenhydrinate (Dramamine);
- metformin (Actoplus Met, Avandamet, Glucophage, Glucovance, Janumet, Kombiglyze, Metaglip, PrandiMet);
- morphine (Kadian, MS Contin, Oramorph);
- procainamide (Procanbid, Pronestyl);
- tenofovir (Viread);

- vancomycin (Vancocin);
- bronchodilators such as ipratropium (Atrovent) or tiotropium (Spiriva);
- bladder or urinary medications such as darifenacin (Enablex), flavoxate (Urispas), oxybutynin (Ditropan, Oxytrol), tolterodine (Detrol), or solifenacin (Vesicare);
- irritable bowel medications such as dicyclomine (Bentyl), hyoscyamine (Hyomax), or propantheline (Pro Banthine); or
- ulcer medications such as glycopyrrolate (Robinul) or mepenzolate (Cantil).

This list is not complete and other drugs may interact with Sanctura XR. Tell your doctor about all medications you use. This includes prescription, over-the-counter, vitamin, and herbal products. Do not start a new medication without telling your doctor.

1 Sanctura XR drug interactions (more detail)

Does Sanctura XR interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Sanctura XR
+
Enter a drug name
Add

More about Sanctura XR (trospium)

- · Check interactions
- · Compare alternatives
- Reviews (8)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: urinary antispasmodics

Patient resources

Other brands

Sanctura

Professional resources

Trospium monograph

• Trospium Chloride Capsules (FDA)

Related treatment guides

- Interstitial Cystitis
- Urinary Frequency
- Urinary Incontinence

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

DRUG STATUS

Availability

Discontinued

Pregnancy & Lactation

্যু Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

8.4 / 10

8 Reviews

Images

Sanctura XR 60 mg (SAN 60)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

. .