



[Home](#)

2. [Hydrochlorothiazide](#)

Hydrochlorothiazide

Generic name: hydrochlorothiazide [*HYE-dro-KLOR-o-THY-a-zide*]

Brand names: HydroDIURIL, [Esidrix](#), [Microzide](#)

Dosage forms: oral capsule (12.5 mg), oral tablet (12.5 mg; 25 mg; 50 mg)

Drug class: [Thiazide diuretics](#)

Medically reviewed by [Sanjai Sinha, MD](#). Last updated on Mar 21, 2024.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

What is hydrochlorothiazide?

[Hydrochlorothiazide](#) is a thiazide diuretic (water pill) that helps prevent your body from absorbing too much salt, which can cause fluid retention.

Hydrochlorothiazide is used to treat [high blood pressure](#) (hypertension).

Hydrochlorothiazide is also used to treat [fluid retention \(edema\)](#) in people with congestive heart failure, cirrhosis of the liver, or kidney disorders, or edema caused by taking steroids or estrogen.

Warnings

You should not use hydrochlorothiazide if you are unable to urinate.

Before using hydrochlorothiazide, tell your doctor if you have liver disease, kidney disease, glaucoma, asthma or allergies, gout, diabetes, or if you are allergic to sulfa drugs or penicillin.

Avoid drinking alcohol, which can increase some of the side effects of hydrochlorothiazide .

Avoid becoming overheated or dehydrated during exercise and in hot weather. Follow your doctor's instructions about the type and amount of liquids you should drink. In some cases, drinking too much liquid can be as unsafe as not drinking enough.

There are many other drugs that can interact with hydrochlorothiazide. Tell your doctor about all medications you use. This includes prescription, over-the-counter, vitamin, and herbal products. Do not start a new medication without telling your doctor. Keep a list of all your medicines and show it to any healthcare provider who treats you.

If you are being treated for high blood pressure, keep using hydrochlorothiazide even if you feel fine. High blood pressure often has no symptoms.

Before taking this medicine

You should not use hydrochlorothiazide if you are allergic to it, or if you are unable to urinate.

To make sure hydrochlorothiazide is safe for you, tell your doctor if you have ever had:

- kidney disease;
- liver disease;
- gout;
- [glaucoma](#);
- low levels of potassium or sodium in your blood;
- high levels of calcium in your blood;
- a parathyroid gland disorder;
- diabetes; or
- an allergy to sulfa drugs or penicillin.

Tell your doctor if you are pregnant or plan to become pregnant. If you take hydrochlorothiazide during pregnancy, your newborn baby may develop [jaundice](#) or other problems.

You should not breastfeed while using this medicine.

Hydrochlorothiazide is not approved for use by anyone younger than 18 years old.

 [Hydrochlorothiazide pregnancy and breastfeeding warnings](#) (more detail)

How should I take hydrochlorothiazide?

Take hydrochlorothiazide exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Call your doctor if you have ongoing [vomiting](#) or [diarrhea](#), or if you are sweating more than usual. You can easily become dehydrated while taking hydrochlorothiazide, which can lead to severely [low blood pressure](#) or a serious electrolyte imbalance.

Your blood pressure will need to be checked often. Your blood and urine may be tested if you have been vomiting or are dehydrated.

If you need surgery, tell your surgeon you currently use this medicine. You may need to stop for a short time.

Keep using hydrochlorothiazide as directed, even if you feel well. High blood pressure often has no symptoms. You may need to use blood pressure medicine for the rest of your life.

Store at room temperature away from moisture, heat, and freezing. Keep the bottle tightly closed when not in use.

 [Hydrochlorothiazide patient tips](#) (more detail)

Hydrochlorothiazide dosing information

Usual Adult Dose for Edema:

Usual dose: 25 mg to 100 mg orally once or twice daily

Comments:

-Some patients respond to intermittent therapy, (i.e., administration on alternate days or on 3 to 5 days each week). Excessive response and undesirable electrolyte imbalance are less likely to occur with intermittent dosing.

Usual Adult Dose for Hypertension:

Initial dose: 25 mg orally once daily

Maintenance dose: May increase to 50 mg orally daily, as a single or 2 divided doses

Comments:

-Patients usually do not require doses in excess of 50 mg daily when used concomitantly with other antihypertensive agents.

Usual Adult Dose for Nephrocalcinosis:

Initial: 25 mg orally once daily

Maintenance dose: May increase to 50 mg twice daily

Usual Adult Dose for Diabetes Insipidus:

Initial: 50 mg orally once daily

Maintenance dose: May increase to 100 mg orally daily

Usual Pediatric Dose for Edema:

Less than 6 months: Up to 3 mg/kg/day (up to 1.5 mg/pound) orally in 2 divided doses

Less than 2 years: 1 to 2 mg/kg/day (0.5 to 1 mg/pound) orally daily as a single dose or in 2 divided doses

Maximum dose 37.5 mg per day

2 to 12 years: 1 to 2 mg/kg/day (0.5 to 1 mg/pound) orally daily as a single dose or in 2 divided doses

Maximum dose 100 mg per day

Usual Pediatric Dose for Hypertension:

Less than 6 months: Up to 3 mg/kg/day (up to 1.5 mg/pound) orally in 2 divided doses

Less than 2 years: 1 to 2 mg/kg/day (0.5 to 1 mg/pound) orally daily as a single dose or in 2 divided doses

Maximum dose 37.5 mg per day

2 to 12 years: 1 to 2 mg/kg/day (0.5 to 1 mg/pound) orally daily as a single dose or in 2 divided doses

Maximum dose 100 mg per day

 [Detailed Hydrochlorothiazide dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include [nausea](#), [dizziness](#), dry mouth, thirst, and muscle pain or weakness.

What should I avoid while taking hydrochlorothiazide?

Hydrochlorothiazide may increase your risk of skin cancer. Avoid sunlight or tanning beds. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors. Your doctor may want you to have skin examinations on a regular basis.

Drinking alcohol with hydrochlorothiazide can cause side effects.

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

Avoid becoming overheated or dehydrated during exercise, in hot weather, or by not drinking enough fluids. Follow your doctor's instructions about the type and amount of liquids you should drink. In some cases, drinking too much liquid can be as unsafe as not drinking enough.

Hydrochlorothiazide side effects

Get emergency medical help if you have **signs of an allergic reaction to hydrochlorothiazide** ([hives](#), difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, [sore throat](#), burning eyes, skin pain, red or purple [skin rash](#) with blistering and peeling).

Call your doctor at once if you have:

- a light-headed feeling;
- eye pain, vision problems;
- jaundice (yellowing of the skin or eyes);
- pale skin, easy bruising, unusual bleeding (nose, mouth, vagina, or rectum);
- shortness of breath, wheezing, cough with foamy mucus, chest pain;
- **dehydration symptoms** - feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and [dry skin](#); or
- **signs of an electrolyte imbalance** - increased thirst or urination, confusion, vomiting, [constipation](#), muscle pain, leg cramps, bone pain, lack of energy, irregular heartbeats, tingly feeling.

Common hydrochlorothiazide side effects may include:

- weakness;
- feeling like you might pass out;
- severe pain in your upper stomach spreading to your back, [nausea and vomiting](#);

- fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath; or
- electrolyte imbalance.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Hydrochlorothiazide side effects](#) (more detail)

Related/similar drugs

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Atenolol

Atenolol is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

Reviews & ratings

7.2 / 10

[276 Reviews](#)

[View more](#)

Valsartan

Valsartan is an angiotensin II receptor blocker (ARB) that may be used to treat high blood pressure ...

Reviews & ratings

5.7 / 10

[152 Reviews](#)

[View more](#)

Spironolactone

Spironolactone is a potassium-sparing

diuretic that is primarily used to treat heart failure, high ...

Reviews & ratings

6.9 / 10

[702 Reviews](#)

[View more](#)

Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

Reviews & ratings

6.4 / 10

[56 Reviews](#)

[View more](#)

Furosemide

Furosemide is a loop diuretic used to treat fluid retention and high blood pressure by increasing ...

Reviews & ratings

6.8 / 10

[121 Reviews](#)

[View more](#)

Losartan

Losartan is used to treat high blood pressure (hypertension). It is also used to lower the risk of ...

Reviews & ratings

4.5 / 10

[578 Reviews](#)

[View more](#)

What other drugs will affect hydrochlorothiazide?

Taking hydrochlorothiazide with other drugs that make you light-headed can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for [anxiety](#) or [seizures](#).

Tell your doctor about all your other medicines, especially:

- [cholestyramine](#), [colestipol](#);
- [insulin](#) or oral diabetes medicine;
- [lithium](#);
- other blood pressure medications;
- steroid medicine; or
- **NSAIDs (nonsteroidal anti-inflammatory drugs)** - [aspirin](#), [ibuprofen](#) ([Advil](#), [Motrin](#)), [naproxen](#) ([Aleve](#)), [celecoxib](#), [diclofenac](#), [indomethacin](#), [meloxicam](#), and others.

This list is not complete. Other drugs may affect hydrochlorothiazide, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible drug interactions are listed here.

 [Hydrochlorothiazide drug interactions](#) (more detail)

Does hydrochlorothiazide interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

+

Add

Popular FAQ

What is the best time of day to take blood pressure medication?



More FAQ

- [How long does hydrochlorothiazide stay in your system?](#)
- [Should you use a diuretic with diazoxide?](#)

[View more FAQ...](#)

References

1. [Hydrochlorothiazide Product Label](#)

More about hydrochlorothiazide

- [Check interactions](#)
- [Compare alternatives](#)

- [Pricing & coupons](#)
- [Reviews \(185\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)
- [Support group](#)
- [Drug class: thiazide diuretics](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

- [HCTZ drug information](#)

Other brands

[Microzide](#), [Aquazide H](#)

Professional resources

- [Hydrochlorothiazide monograph](#)
- [Hydrochlorothiazide \(FDA\)](#)
- [Hydrochlorothiazide Capsules \(FDA\)](#)
- [Hydrochlorothiazide Tablets \(FDA\)](#)

Other brands

[Microzide](#), [Esidrix](#), [Inzirgo](#)

Related treatment guides

- [Edema](#)
- [High Blood Pressure](#)
- [Diabetes Insipidus](#)
- [Nephrocalcinosis](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use hydrochlorothiazide only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal

circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 13.01.

DRUG STATUS

	Availability	
Rx	Prescription only	
	Pregnancy & Lactation	
	Risk data available	
	CSA Schedule*	
N/A	Not a controlled drug	▼
	Approval History	
	Drug history at FDA	▼
	WADA Class	
	Anti-Doping Classification	▼

User Reviews & Ratings

5.2 / 10

[185 Reviews](#)

Images

[Hydrochlorothiazide 25 mg \(U 128\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

- [About Drugs.com](#)
- [Advertising policy](#)
- [Attribution & citations](#)
- [Terms & privacy](#)**

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)


Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.