



[Home](#)

2. Mono-Linyah

Mono-Linyah

Generic name: [ethinyl estradiol and norgestimate](#) [*ETH-i-nil-ES-tra-DYE-ol-and-nor-JES-ti-mate*]

Brand names: [Estaylla](#), [Mili](#), Mono-Linyah, [Nymyo](#), [Sprintec](#), ... [show all 19 brands](#)

Drug class: [Contraceptives](#)



[Medically reviewed](#) by Drugs.com on Apr 4, 2025. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is Mono-Linyah?

Mono-Linyah is a combination [birth control pill](#) containing female hormones that prevent ovulation (the release of an egg from an ovary). Mono-Linyah also causes changes in your cervical mucus and uterine lining, making it harder for sperm to reach the uterus and harder for a fertilized egg to attach to the uterus.

Mono-Linyah is used as contraception to prevent pregnancy. **There are many available brands of this medicine. Not all brands are listed on this leaflet.**

Mono-Linyah may also be used for purposes not listed in this medication guide.

Birth control pills side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have:

- **signs of a stroke**--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance;
- **signs of a blood clot**--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood, pain or warmth in one or both legs;
- **heart attack symptoms**--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;
- **liver problems**--loss of appetite, upper stomach pain, tiredness, fever, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- severe headache, pounding in your neck or ears;
- swelling in your hands, ankles, or feet;
- changes in the pattern or severity of migraine headaches;

- a breast lump; or
- **symptoms of depression**--sleep problems, weakness, tired feeling, mood changes.

Common side effects of Mono-Linyah may include:

- stomach pain, gas, nausea, vomiting;
- breast tenderness;
- acne, darkening of facial skin;
- headache, nervousness, mood changes;
- problems with contact lenses;
- changes in weight;
- [breakthrough bleeding](#);
- vaginal itching or discharge; or
- rash.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Mono-Linyah side effects](#) (more detail)

Related/similar drugs

Premarin

Premarin is used to treat symptoms of menopause such as hot flashes, and vaginal dryness. Learn ...

Reviews & ratings

7.7 / 10

[84 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

Reviews & ratings

6.1 / 10

[2,624 Reviews](#)

[View more](#)

Depo Provera

Depo-Provera is used as contraception to prevent pregnancy. It is also used to reduce pain cause by ...

Reviews & ratings

4.2 / 10

[1,967 Reviews](#)

[View more](#)

Leuprolide

Leuprolide is used to treat endometriosis, prostate cancer, and early-onset puberty. Includes ...

Reviews & ratings

5.6 / 10

[404 Reviews](#)

[View more](#)

Minocycline

Minocycline is used to treat infections, such as urinary tract infections, acne and chlamydia ...

Reviews & ratings

5.9 / 10

[564 Reviews](#)

[View more](#)

Warnings

Do not use birth control pills if you are pregnant or if you have recently had a baby.

You should not use birth control pills if you have: uncontrolled [high blood pressure](#), [heart disease](#), [coronary artery disease](#), circulation problems (especially with diabetes), undiagnosed vaginal bleeding, [liver disease](#) or liver cancer, severe migraine headaches, if you also take certain hepatitis C medication, if you will have major surgery, if you smoke and are over 35, or if you have ever had a [heart attack](#), a stroke, a blood clot, jaundice caused by pregnancy or [birth control pills](#), or cancer of the breast, uterus/cervix, or vagina.

Taking birth control pills can increase your risk of blood clots, stroke, or heart attack.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. You should not take [birth control pills](#) if you smoke and are over 35 years old.

Before taking this medicine

Taking birth control pills can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have high blood pressure, diabetes, [high cholesterol](#), or if you are overweight. Your risk of stroke or blood clot is highest during your first year of taking birth control pills. Your risk is also high when you restart birth control pills after not taking them for 4 weeks or longer.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. Your risk increases the older you are and the more you smoke. You should not take combination birth control pills if you smoke and are over 35 years old.

Do not use if you are pregnant. Stop using Mono-Linyah and tell your doctor right away if you become pregnant, or if you miss two menstrual periods in a row. If you have recently had a baby, wait at least 4 weeks before taking birth control pills.

You should not take birth control pills if you have:

- untreated or uncontrolled high blood pressure;
- heart disease (chest pain, coronary artery disease, history of heart attack, stroke, or blood clot);
- an increased risk of having blood clots due to a heart problem or a hereditary blood disorder;
- circulation problems (especially if caused by diabetes);
- a history of hormone-related cancer, or cancer of the breast, uterus/cervix, or vagina;
- unusual vaginal bleeding that has not been checked by a doctor;
- liver disease or liver cancer;
- severe migraine headaches (aura, numbness, vision changes), especially if you are older than 35; or
- if you take any hepatitis C medication containing [ombitasvir/paritaprevir/ritonavir](#) (Technivie).

Tell your doctor if you have ever had:

- heart problems, high blood pressure, or if you are prone to having blood clots;
- high cholesterol or triglycerides, or if you are overweight;
- depression;

- a seizure or [migraine headache](#);
- diabetes, [gallbladder disease](#), [underactive thyroid](#);
- liver or kidney disease;
- jaundice caused by pregnancy or birth control pills;
- irregular menstrual cycles; or
- [fibrocystic breast disease](#), lumps, nodules, or an abnormal mammogram.

This medicine can slow breast milk production. Tell your doctor if you are breast-feeding.

How should I take birth control pills?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

You may need to use back-up birth control, such as condoms with spermicide, when you first start using this medicine. Follow your doctor's instructions.

Take one pill every day, no more than 24 hours apart. When the pills run out, start a new pack the following day. **You may get pregnant if you do not take one pill daily.**

Some birth control packs contain "reminder" pills to keep you on your regular cycle. Your period will usually begin while you are using these reminder pills.

You may have breakthrough bleeding, especially during the first 3 months. Tell your doctor if this bleeding continues or is very heavy.

Use a back-up birth control if you are sick with severe vomiting or diarrhea.

If you need major surgery or will be on long-term bed rest, you may need to stop using this medicine for a short time. Any doctor or surgeon who treats you should know that you are using estradiol and [levonorgestrel](#).

While taking birth control pills, you will need to visit your doctor regularly.

Store this medicine at room temperature away from moisture and heat.

 [Detailed Mono-Linyah dosage information](#)

What happens if I miss a dose?

Follow the instructions provided with your medicine. Missing a pill increases your risk of becoming pregnant.

If you miss 1 active pill, take 2 pills on the day you remember. Then take 1 pill per day for the rest of the pack.

If you miss 2 active pills in a row in Week 1 or 2, take 2 pills per day for 2 days in a row. Then take 1 pill per day for the rest of the pack. Use back-up birth control for at least 7 days following the missed pills.

If you miss 2 active pills in a row in Week 3, throw out the pack and start a new pack the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack that day.

If you miss 3 active pills in a row in Week 1, 2, or 3, throw out the pack and start a new pack on the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack that day.

If you miss 2 or more pills, you may not have a period during the month. **If you miss a period for 2 months in a row, call your doctor because you might be pregnant.**

If you miss a reminder pill, throw it away and keep taking one reminder pill per day until the pack is empty.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking birth control pills?

Do not smoke while taking birth control pills, especially if you are older than 35 years of age.

Birth control pills will not protect you from [sexually transmitted diseases](#), including HIV and AIDS. Using a condom is the only way to protect yourself from these diseases.

What other drugs will affect birth control pills?

Other drugs may affect birth control pills, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). **Some drugs can make birth control pills less effective, which may result in pregnancy.** Tell your doctor about all your current medicines and any medicine you start or stop using.

 [Mono-Linyah drug interactions](#) (more detail)

Does Mono-Linyah interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Mono-Linyah

+

Enter a drug name

Add

More about Mono-Linyah (ethinyl estradiol / norgestimate)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(122\)](#)
- [Drug images](#)

- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: contraceptives](#)
- [En español](#)

Patient resources

Other brands

[Sprintec](#), [Estarylla](#), [Tri-Sprintec](#), [Ortho Tri-Cyclen](#), ... +20 more

Professional resources

- [Mono-Linyah prescribing information](#)

Other brands

[Sprintec](#), [Estarylla](#), [Tri-Sprintec](#), [Ortho Tri-Cyclen](#), ... +21 more

Related treatment guides

- [Abnormal Uterine Bleeding](#)
- [Acne](#)
- [Birth Control](#)
- [Endometriosis](#)
- [Gonadotropin Inhibition](#)
- [Ovarian Cysts](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 17.04.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation



Risk data available

CSA Schedule*

N/A Not a controlled drug



Approval History



Drug history at FDA



User Reviews & Ratings

4.7 / 10

[122 Reviews](#)

Images

[Mono-Linyah ethinyl estradiol 0.035 mg / norgestimate 0.25 mg \(C3\)](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and

others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.