



[Home](#)

2. [Polyethylene glycol 335...](#)
3. [Consumer Information](#)

GaviLyte-N With Flavor Packs

Generic name: polyethylene glycol electrolyte solution [*pall-ee-ETH-il-eeen-GLYE-kol-ee-LEK-troe-lyte*]

Brand names: GaviLyte-C With Lemon Flavor Pack, GaviLyte-G With Lemon Flavor Pack, GaviLyte-N With Flavor Packs, [GoLYTELY](#), [MoviPrep](#), ... [show all 20 brands](#)

Drug class: [Laxatives](#)



[Medically reviewed](#) by Drugs.com on Jun 12, 2024. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is GaviLyte-N With Flavor Packs?

[GaviLyte-N](#) With Flavor Packs is a laxative solution that stimulates bowel movements. This medication also contains minerals to replace electrolytes that are passed from the body in the stool.

GaviLyte-N With Flavor Packs is used to clean the bowel before [colonoscopy](#), a barium x-ray, or other intestinal procedures.

GaviLyte-N With Flavor Packs may also be used for purposes not listed in this medication guide.

GaviLyte-N With Flavor Packs side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

GaviLyte-N With Flavor Packs may cause serious side effects. Call your doctor at once if you have:

- no bowel movement within 2 hours after use;
- vomiting;
- dizziness, feeling like you might pass out;
- little or no urination;
- a seizure; or
- **signs of an electrolyte imbalance**--increased thirst or urination, dry mouth, confusion, [constipation](#), [muscle pain](#) or weakness, leg cramps, irregular heartbeats, tingly feeling.

You may need to drink the liquid more slowly, or stop using it for a short time if you have certain side effects. **Call your doctor for instructions if you have:**

- gagging, choking, severe stomach pain or bloating;
- nausea, vomiting, headache, trouble drinking liquids, little or no urinating; or
- fever, sudden or severe stomach pain, severe diarrhea, rectal bleeding or bright red bowel movements.

Common side effects of GaviLyte-N With Flavor Packs may include:

- vomiting, stomach pain, indigestion, bloating;
- rectal pain or irritation;
- hunger, thirst, mild nausea;
- trouble sleeping; or
- dizziness, chills.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Polyethylene glycol 3350 with electrolytes side effects](#) (more detail)

Related/similar drugs

MiraLAX

MiraLAX (polyethylene glycol) is used as a laxative to treat occasional constipation or irregular ...

Reviews & ratings

7.2 / 10

[262 Reviews](#)

[View more](#)

Dulcolax

Dulcolax (bisacodyl) is a laxative used to treat constipation. Includes Dulcolax side effects ...

Reviews & ratings

5.3 / 10

[1,158 Reviews](#)

[View more](#)

FEATURED

OmvoH

OmvoH is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Movantik

Movantik reduces constipation caused by opioid pain medicine used to treat severe chronic pain. It ...

Reviews & ratings

4.9 / 10

[265 Reviews](#)

[View more](#)

Sutab

Sutab (sodium sulfate, magnesium sulfate, and potassium chloride) is used for colonoscopy ...

Reviews & ratings

6.3 / 10

[126 Reviews](#)

[View more](#)

Senna

Senna is a laxative is used treat constipation. Includes senna side effects, interactions and ...

Reviews & ratings

5.3 / 10

[308 Reviews](#)

[View more](#)

Warnings

Do not use GaviLyte-N With Flavor Packs if you have a perforated bowel, a [bowel obstruction](#) or severe constipation, or colitis or toxic megacolon. GaviLyte-N With Flavor Packs can cause dangerous or life-threatening side effects in people with these conditions.

Before taking this medicine

You should not use this medicine if you are allergic to polyethylene glycol or any other electrolyte solutions (such as [Pedialyte](#) or Gatorade), or if you have:

- a perforated bowel;
- a bowel obstruction or severe constipation; or
- colitis or toxic megacolon.

GaviLyte-N With Flavor Packs can cause dangerous or life-threatening side effects in people with these conditions.

People with [eating disorders](#) (such as anorexia or bulimia) should not use this medicine without the advice of a doctor.

Tell your doctor if you have ever had:

- heart problems, or a [heart attack](#);
- an electrolyte imbalance (such as low levels of potassium or sodium in your blood);
- kidney disease;
- a seizure;
- gastroesophageal reflux disease (GERD), [ulcerative colitis](#), or other stomach or bowel disorder;
- trouble swallowing, aspiration (accidentally inhaling food or drink);
- a genetic enzyme deficiency called glucose-6-phosphate dehydrogenase (G6PD) deficiency; or
- a drug or alcohol addiction.

Tell your doctor if you are pregnant or breast-feeding.

This medicine may contain phenylalanine. Talk to your doctor before using if you have [phenylketonuria](#) (PKU).

How should I take GaviLyte-N With Flavor Packs?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Polyethylene glycol electrolyte powder must be mixed with water into a solution before using it. Do not add any flavorings such as sugar, honey, artificial sweetener, fruit juices, or other beverages.

Shake the mixture well just before you measure a dose. Drink GaviLyte-N With Flavor Packs in the exact portions at the exact time intervals prescribed by your doctor.

This medicine comes with instructions about when and what to eat or drink on your first day of treatment. **Each brand may have different instructions.**

Do not drink GaviLyte-N With Flavor Packs if it has been less than 1 hour since you last ate solid food. For best results, take the medicine 2 to 4 hours after you last ate.

The first watery stool should appear within 1 hour after you start drinking GaviLyte-N With Flavor Packs. Keep taking the

medicine until you have completed all doses prescribed by your doctor.

Drink plenty of **clear liquids** (water, broth, black coffee, tea, clear soda) before, during, and after you take this medicine. **Avoid foods and beverages that contain pulp or are red or purple in color.** You may also eat popsicles (not fruit bars or fudge bars) or gelatin without fruit pieces or toppings.

Do not eat or drink anything within 2 hours before your colonoscopy or other medical test.

Store the unmixed powder at room temperature away from moisture and heat.

Store the mixed solution in a refrigerator in an upright position.

Throw away any GaviLyte-N With Flavor Packs you have not used within 24 to 48 hours after it was mixed (follow directions for your specific brand of this medicine).

i [Detailed Polyethylene glycol 3350 with electrolytes dosage information](#)

What happens if I miss a dose?

Talk to your doctor if you cannot drink all of the solution prescribed for you. Your test or procedure may need to be rescheduled if your bowel is not completely cleansed.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking GaviLyte-N With Flavor Packs?

Avoid taking other medications, vitamins, or mineral supplements within 1 hour before drinking GaviLyte-N With Flavor Packs. A bowel cleansing can make it harder for your body to absorb other medicines you take by mouth.

Do not use other laxatives while using GaviLyte-N With Flavor Packs unless your doctor has told you to.

What other drugs will affect GaviLyte-N With Flavor Packs?

Tell your doctor about all your other medicines, especially:

- heart or blood pressure medication;
- a diuretic or "water pill";
- medicine to treat anxiety, [depression](#), or mental illness;
- medications to treat kidney problems or low sodium levels ([hyponatremia](#));
- seizure medication; or
- **NSAIDs (nonsteroidal anti-inflammatory drugs)**--aspirin, [ibuprofen](#) (Advil, Motrin), naproxen (Aleve), [celecoxib](#), [diclofenac](#), [indomethacin](#), [meloxicam](#), and others.

This list is not complete. Other drugs may affect GaviLyte-N With Flavor Packs, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

Does polyethylene glycol 3350 with electrolytes interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

polyethylene glycol 3350 with electrolytes

+

Enter a drug name

Add

More about polyethylene glycol 3350 with electrolytes

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(1,121\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: laxatives](#)

Patient resources

- [Polyethylene glycol electrolyte solution drug information](#)
- [Polyethylene glycol 3350, electrolytes, and ascorbic acid \(Advanced Reading\)](#)
- [Polyethylene glycol 3350, sodium and potassium salts \(Advanced Reading\)](#)
- [Polyethylene glycol 3350, sodium ascorbate, sodium sulfate, ascorbic acid, sodium chloride, and potassium chloride \(Advanced Reading\)](#)
- [Polyethylene Glycol-Electrolyte Solution \(MoviPrep, Plenvu\)](#)
- [Polyethylene Glycol-Electrolyte Solution Prep](#)

Other brands

[GoLYTELY](#), [GaviLyte-G](#), [Plenvu](#), [PEG-3350 with Electrolytes](#), ... +7 more

Professional resources

- [PEG-3350, Electrolytes and Ascorbate prescribing information](#)
- [PEG-3350, Sodium Chloride, Sodium Bicarbonate and Potassium Chloride \(FDA\)](#)
- [Polyethylene Glycol \(FDA\)](#)

Other brands

GoLYTELY, GaviLyte-G, Plenvu, Suflave, ... +6 more

Related treatment guides

- [Bowel Preparation](#)
- [Constipation, Chronic](#)
- [Gastrointestinal Decontamination](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.03.

DRUG STATUS

Availability

Rx Prescription only

[Pregnancy & Lactation](#)

 Risk data available

CSA Schedule*

N/A Not a controlled drug



User Reviews & Ratings

6.0 / 10


[1,121 Reviews](#)

Images

Suflave (polyethylene glycol 3350 with electrolytes) 178.7 g polyethylene glycol 3350, 7.3 g sodium sulfate, 1.12 g potassium chloride, 0.9 g magnesium sulfate, 0.5 g sodium chloride for reconstitution

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.