

Home

Multaq

Multaq ⊲»

Generic name: dronedarone [droe-NE-da-rone]

Drug class: Group III antiarrhythmics

Medically reviewed by Philip Thornton, DipPharm. Last updated on May 31, 2024.

Uses Warnings Before taking Dosage Side effects Interactions

What is Multaq?

Multaq is a heart rhythm medicine that helps maintain normal heartbeats in certain people with life-threatening rhythm disorders of the atrium (the upper chambers of the heart that allow blood to flow into the heart).

Multag helps lower your risk of needing to be hospitalized for a heart rhythm disorder called atrial fibrillation.

Multag is for adults who have had this disorder in the past, but now have normal heart rhythm.

Warnings

You should not use Multaq if you have severe liver disease, if you are pregnant or breast-feeding, or if you have ever used amiodarone and then had liver or lung problems. You should not use Multaq if you have a serious heart condition such as very slow heartbeats, "sick sinus syndrome," or "AV block" (unless you have a pacemaker).

Multaq can double your risk of death if you have certain heart conditions. You should not use this medicine if you have severe heart failure, if you were recently hospitalized for worsening heart failure symptoms, or if you have a "permanent" type of atrial fibrillation (this will be determined by your doctor).

Check your pulse often, and tell your doctor right away if you notice an irregular rhythm.

Tell your doctor about all your current medicines and any you start or stop using. Many drugs can interact with dronedarone, and some drugs should not be used together.

Multaq can cause liver problems. Call your doctor at once if you have symptoms such as nausea, loss of appetite, unusual tiredness, dark urine, or yellowing of your skin or eyes.

Before taking this medicine

You should not use Multaq if you are allergic to dronedarone, or if you have:

· severe liver disease;

- a serious heart condition such as "sick sinus syndrome," "AV block" (unless you have a pacemaker), or very slow heartbeats that have caused you to faint;
- if you are pregnant or breastfeeding; or
- if you used a medicine called amiodarone and then had lung problems or liver problems.

Dronedarone can double your risk of death if you have certain heart conditions. You should not use this medicine if:

- you have severe heart failure;
- you were recently hospitalized for worsening heart failure symptoms (shortness of breath, chest tightness, night-time breathing problems, swelling, rapid weight gain); or
- you have a "permanent" atrial fibrillation that cannot be changed back to a normal rhythm (this will be determined by your doctor).

Many drugs can interact and cause dangerous effects. Some drugs should not be used together with Multaq. Your doctor may change your treatment plan if you also use:

- other heart rhythm medicines;
- cyclosporine;
- ritonavir or other antiviral medicine;
- an antibiotic such as clarithromycin or telithromycin;
- an antidepressant such as amitriptyline, clomipramine, doxepin, imipramine, nefazodone, and others;
- antifungal medicine such as itraconazole, ketoconazole, or voriconazole; or
- antipsychotic medicine such as chlorpromazine, fluphenazine, promethazine, thioridazine, and others.

Tell your doctor if you have other heart problems.

Dronedarone may harm an unborn baby. Use effective birth control to prevent pregnancy, and tell your doctor if you become pregnant.

You should not breastfeed while using Multaq.

Multag pregnancy and breastfeeding warnings (more detail)

How should I take Multaq?

Take Multaq exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

Multag is usually taken with morning and evening meals.

Your heart function may need to be checked every 3 months to help determine if you still need Multaq. Do not stop taking Multaq without your doctor's advice.

Check your pulse often, and call your doctor if you notice an irregular rhythm.

Store at room temperature away from heat and moisture.

Dosing information

Usual Adult Dose for Atrial Fibrillation:

400 mg orally 2 times a day with a meal

Use: To reduce the risk of cardiovascular hospitalization in patients with paroxysmal or persistent atrial fibrillation (AF) or atrial flutter (AFL), with a recent episode of AF/AFL and associated cardiovascular risk factors (i.e., age greater than 70 years, hypertension, diabetes, prior cerebrovascular accident, left atrial diameter 50 mm or more, or left ventricular ejection fraction [LVEF] less than 40%), who are in sinus rhythm or who will be cardioverted.

Usual Adult Dose for Atrial Flutter:

400 mg orally 2 times a day with a meal

Use: To reduce the risk of cardiovascular hospitalization in patients with paroxysmal or persistent atrial fibrillation (AF) or atrial flutter (AFL), with a recent episode of AF/AFL and associated cardiovascular risk factors (i.e., age greater than 70 years, hypertension, diabetes, prior cerebrovascular accident, left atrial diameter 50 mm or more, or left ventricular ejection fraction [LVEF] less than 40%), who are in sinus rhythm or who will be cardioverted.

Detailed Multag dosage information

What happens if I miss a dose?

Skip the missed dose and use your next dose at the regular time. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Multaq?

Grapefruit may interact with dronedarone and cause side effects. Avoid consuming grapefruit while taking Multaq.

Multaq side effects

Get emergency medical help if you have **signs of an allergic reaction to Multaq:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- dry cough, shortness of breath;
- little or no urination;
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out);
- heart problems swelling, rapid weight gain, feeling short of breath;
- liver problems loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, jaundice (yellowing

of the skin or eyes); or

• low magnesium or potassium levels - dizziness, numbness or tingling, irregular heartbeats, fluttering in your chest, increased thirst or urination, muscle cramps or limp feeling.

Common Multaq side effects may include:

- stomach pain, indigestion, nausea, vomiting, diarrhea;
- · feeling weak or tired; or
- skin rash, itching, or redness.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Multaq side effects (more detail)

Related/similar drugs

Xarelto

Xarelto (rivaroxaban) is a factor Xa inhibitor used to reduce the risk of blood clots and stroke in ...

Reviews & ratings

5.4 / 10

224 Reviews

View more

Cardizem

Cardizem is used to treat hypertension (high blood pressure), angina (chest pain), and heart rhythm ...

Reviews & ratings

7.3 / 10

12 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Lanoxin

Lanoxin (digoxin) is used to treat congestive heart failure and atrial fibrillation. Includes ...

Reviews & ratings

Add a review

View more

What other drugs will affect Multaq?

Multaq can cause a serious heart problem. Your risk may be higher if you also use certain other medicines for

infections, asthma, heart problems, high blood pressure, depression, mental illness, cancer, malaria, or HIV.

Tell your doctor about all your current medicines. Many drugs can interact with dronedarone, especially:

- digoxin;
- St. John's wort;
- a blood thinner (warfarin, Coumadin, Jantoven);
- a diuretic or "water pill";
- medicines to treat tuberculosis;
- medicine to prevent organ transplant rejection;
- seizure medicine; or
- "statin" cholesterol medication (Lipitor, Zocor, Vytorin, and others).

This list is not complete and many other drugs may interact with dronedarone. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Multaq drug interactions (more detail)

Does Multaq interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



References

1. Multaq Product Label

More about Multaq (dronedarone)

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (43)
- Drug images
- Side effects
- Dosage information
- During pregnancy

- · Generic availability
- · FDA approval history
- Drug class: group III antiarrhythmics
- En español

Professional resources

- Multaq prescribing information
- Dronedarone (AHFS Monograph)

Related treatment guides

- Atrial Fibrillation
- Atrial Flutter

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Multaq only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 13.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ം Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+ years FDA approved 2009

User Reviews & Ratings

6.5 / 10

43 Reviews

Images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.