

Home

2. Degarelix

Degarelix

Generic name: degarelix [*DEG-a-REL-ix*]

Brand name: Firmagon

Drug classes: Gonadotropin-releasing hormone antagonists, Hormones / antineoplastics

Medically reviewed by Philip Thornton, DipPharm. Last updated on Aug 23, 2023.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is degarelix?

Degarelix is a man-made form of a protein that reduces the amount of certain hormones in the body, including testosterone.

Degarelix injection is used to treat prostate cancer. It works by decreasing the amount of testosterone in the body, which the tumor needs to grow.

It is not known if degarelix is safe or effective in children.

Warnings

Although this medicine is not for use by women, degarelix should not be used by a woman who is pregnant or breastfeeding, or who may become pregnant.

Before you receive degarelix, tell your doctor if you have liver or kidney disease, a personal or family history of "Long QT syndrome," congestive heart failure, or an electrolyte imbalance, such as low or high levels of potassium or magnesium in your blood.

Tell your doctor about all other medications you use, especially a heart rhythm medication.

Degarelix is usually given once every 28 days. Follow your doctor's instructions.

To be sure this medication is helping your condition, your blood may need to be tested on a regular basis. Do not miss any follow-up visits to your doctor.

Before taking this medicine

You should not use degarelix if you are allergic to it.

To make sure this medicine is safe for you, tell your doctor if you have:

- · heart problems;
- long QT syndrome (in you or a family member);
- · liver or kidney disease; or
- an electrolyte imbalance, such as low or high levels of potassium, calcium, or magnesium in your blood.

Although degarelix is not for use by women, this medicine can harm an unborn baby or cause miscarriage if a woman is exposed to it during pregnancy. **Degarelix should not be used by a woman who is pregnant.**

This medicine may affect fertility (your ability to have children), whether you are a man or a woman.

Degarelix also should not be used by a woman who is breastfeeding a baby.

Degarelix pregnancy and breastfeeding warnings (more detail)

How is degarelix given?

Degarelix is injected under the skin around your stomach. A healthcare provider will give you this injection.

On the day of your injection, avoid wearing a belt, tight waistband, or tight clothing around your stomach where the injection will be given.

Degarelix is usually given once every 28 days. Follow your doctor's instructions.

The first time you receive this medicine, you will be given two injections. At your monthly follow-up visits you will receive only one injection.

You will need frequent medical tests.

Degarelix can affect the results of certain medical tests. Tell any doctor who treats you that you are using this medicine.

Dosing information

Usual Adult Dose for Prostate Cancer:

-Initial Dose: 240 mg administered as 2 subcutaneous injections of 120 mg each at a concentration of 40 mg/mL.

-Maintenance Dose: 80 mg administered as 1 subcutaneous injection at a concentration of 20 mg/mL every 28 days.

Comments: Administer the first maintenance dose 28 days after the initial dose.

Use: Treatment of advanced prostate cancer

Detailed Degarelix dosage information

What happens if I miss a dose?

Call your doctor for instructions if you miss an appointment for your injection.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while receiving degarelix?

Avoid rubbing or scratching the skin where a degarelix injection was given.

Degarelix can pass into body fluids (urine, feces, vomit). For at least 48 hours after you receive a dose, avoid allowing your body fluids to come into contact with your hands or other surfaces. Caregivers should wear rubber gloves while cleaning up a patient's body fluids, handling contaminated trash or laundry or changing diapers. Wash hands before and after removing gloves. Wash soiled clothing and linens separately from other laundry.

Degarelix side effects

Get emergency medical help if you have **signs of an allergic reaction to degarelix:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- · fast or pounding heartbeats, fluttering in your chest;
- · shortness of breath;
- sudden dizziness (like you might pass out); or
- severe headache, blurred vision, pounding in your neck or ears.

Common degarelix side effects may include:

- · hot flashes;
- · weight gain;
- · abnormal liver function tests; or
- · pain, swelling, redness, itching, or a hard lump where the medicine was injected.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Degarelix side effects (more detail)

Related/similar drugs

Premarin

Premarin is used to treat symptoms of menopause such as hot flashes, and vaginal dryness. Learn ...

Reviews & ratings

7.7 / 10

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Estrace

Estrace (estradiol) is used to treat symptoms of menopause such as hot flashes and vaginal dryness ...

Reviews & ratings

7.3 / 10

12 Reviews

View more

Xtandi

Xtandi (enzalutamide) is an antiandrogen used to treat prostate prostate cancer. Includes Xtandi ...

Reviews & ratings

5.6 / 10

21 Reviews

View more

Zytiga

Zytiga (abiraterone) is used together with prednisone to treat prostate cancer that has spread to ...

Reviews & ratings

8.2 / 10

41 Reviews

View more

Lynparza

Lynparza (olaparib) is used to treat ovarian cancer, breast cancer, pancreatic cancer, and prostate ...

Reviews & ratings

7.4 / 10

17 Reviews

View more

Casodex

Casodex (bicalutamide) is an antiandrogen and is used together with LHRH to treat prostate cancer ...

Reviews & ratings

7 Reviews

View more

Bicalutamide

Bicalutamide is a prescription medicine used to treat prostate cancer; Includes bicalutamide side ...

Reviews & ratings

8.4 / 10

13 Reviews

View more

What other drugs will affect degarelix?

Degarelix can cause a serious heart problem. Your risk may be higher if you also use certain other medicines for infections, asthma, heart problems, high blood pressure, depression, mental illness, cancer, malaria, or HIV.

Other drugs may interact with degarelix, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

Degarelix drug interactions (more detail)

Does degarelix interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

degarelix	
+	
Enter a drug name	Add

Popular FAQ

Is Firmagon (degarelix) the same as Lupron?

How long does it take for Degarelix to work?

References

1. Degarelix Product Label

More about degarelix

- · Check interactions
- Compare alternatives
- Reviews (15)
- Side effects
- Dosage information
- During pregnancy
- Drug class: gonadotropin-releasing hormone antagonists
- En español

Patient resources

Other brands

Firmagon

Professional resources

• Degarelix monograph

Other brands

Firmagon

Related treatment guides

Prostate Cancer

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use degarelix only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 7.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্য Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

8.0 / 10

15 Reviews

Images

Firmagon (degarelix) starting dose: two single-dose vials each delivering 120 mg



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

×