

Home

Flexeril

Flexeril 🖘

Generic name: cyclobenzaprine [sye-kloe-BEN-za-preen]

Brand names: Amrix, Fexmid, Flexeril **Drug class:** Skeletal muscle relaxants

Medically reviewed by <u>Sophia Entringer</u>, <u>PharmD</u>. Last updated on Aug 23, 2023.

Uses Warnings Before taking Dosage Side effects Interactions FAQ



The Flexeril brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be generic equivalents available.

What is Flexeril?

Flexeril is a muscle relaxant. It works by blocking nerve impulses (or pain sensations) that are sent to your brain.

Flexeril is used together with rest and physical therapy to treat skeletal muscle conditions such as pain, injury, or spasms.

Flexeril may also be used for purposes not listed in this medication guide.

Warnings

You should not use Flexeril if you have a thyroid disorder, heart block, congestive heart failure, a heart rhythm disorder, or you have recently had a heart attack.

Do not use cyclobenzaprine if you have taken an MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, or tranylcypromine.

Before taking this medicine

You should not use Flexeril if you are allergic to cyclobenzaprine, or if you have:

- · a thyroid disorder;
- heart block, heart rhythm disorder, congestive heart failure; or
- if you have recently had a heart attack.

Flexeril is not approved for use by anyone younger than 15 years old.

Do not use Flexeril if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

Some medicines can interact with cyclobenzaprine and cause a serious condition called serotonin syndrome. Be sure your doctor knows if you also take stimulant medicine, opioid medicine, herbal products, or medicine for depression, mental illness, Parkinson's disease, migraine headaches, serious infections, or prevention of nausea and vomiting. Ask your doctor before making any changes in how or when you take your medications.

To make sure Flexeril is safe for you, tell your doctor if you have ever had:

- · liver disease:
- glaucoma;
- enlarged prostate; or
- · problems with urination.

It is not known whether cyclobenzaprine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It may not be safe to breast-feed while using this medicine. Ask your doctor about any risk.

Older adults may be more sensitive to the effects of this medicine.

flexeril pregnancy and breastfeeding warnings (more detail)

How should I take Flexeril?

Flexeril is usually taken once daily for only 2 or 3 weeks. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Take the medicine at the same time each day.

Call your doctor if your symptoms do not improve after 3 weeks, or if they get worse.

Store at room temperature away from moisture, heat, and light.

Detailed Flexeril dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of cyclobenzaprine** can be fatal.

Overdose symptoms may include severe drowsiness, vomiting, fast heartbeats, tremors, agitation, or hallucinations.

What to avoid

Avoid driving or hazardous activity until you know how cyclobenzaprine will affect you. Your reactions could be impaired.

Avoid drinking alcohol. Dangerous side effects could occur.

Flexeril side effects

Get emergency medical help if you have **signs of an allergic reaction to Flexeril**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using Flexeril and call your doctor at once if you have:

- · fast or irregular heartbeats;
- chest pain or pressure, pain spreading to your jaw or shoulder; or
- sudden numbness or weakness (especially on one side of the body), slurred speech, balance problems.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Serious side effects may be more likely in older adults.

Common Flexeril side effects may include:

- · drowsiness, tiredness;
- · headache, dizziness;
- dry mouth; or
- upset stomach, nausea, constipation.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

flexeril side effects (more detail)

Related/similar drugs

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-thecounter painkiller and fever ...

Reviews & ratings

6.1 / 10

13 Reviews

View more

Ibuprofen

Ibuprofen (Advil, Motrin) treats minor aches and pains caused by the common cold, headaches ...

Reviews & ratings

7.3 / 10

240 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

Reviews & ratings

5.8 / 10

30 Reviews

View more

Robaxin

Robaxin (methocarbamol) is used to treat skeletal muscle conditions such as pain or injury ...

Reviews & ratings

6.3 / 10

80 Reviews

View more

Advil

Advil (ibuprofen) reduces fever, pain and inflammation caused by

headaches, toothaches, arthritis ...

Reviews & ratings

6.9 / 10

41 Reviews

View more

Valium

Valium is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

Reviews & ratings

8.6 / 10

341 Reviews

View more

Voltaren

Voltaren (diclofenac) is used to treat pain or inflammation caused by arthritis or ankylosing ...

Reviews & ratings

7.3 / 10

165 Reviews

View more

Baclofen

Baclofen is a muscle relaxant used to treat spasticity and muscle pain in conditions like multiple ...

Reviews & ratings

7.0 / 10

447 Reviews

View more

Diclofenac

Diclofenac is a nonsteroidal antiinflammatory drug used to treat pain and inflammation associated ...

- . . .

Reviews & ratings

7.3 / 10

695 Reviews

View more

What other drugs will affect Flexeril?

Using Flexeril with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your other medicines, especially:

- bupropion (Zyban, for smoking cessation);
- · meperidine;
- tramadol;
- · verapamil;
- · narcotics;
- cold or allergy medicine that contains an antihistamine (Benadryl and others);
- medicine to treat Parkinson's disease;
- medicine to treat excess stomach acid, stomach ulcer, motion sickness, or irritable bowel syndrome;
- medicine to treat overactive bladder; or
- bronchodilator asthma medication.

This list is not complete. Other drugs may interact with cyclobenzaprine, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

flexeril drug interactions (more detail)

Does Flexeril interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

Is Flexeril addictive, will it show up on a drug test?	~
Does cyclobenzaprine make you sleepy?	~
How long does it take for cyclobenzaprine to work?	~
How much cyclobenzaprine can I take?	~
Is cyclobenzaprine a controlled substance or addictive?	~

More FAQ

• How long does Flexeril stay in your system?

View more FAQ...

More about Flexeril (cyclobenzaprine)

- · Check interactions
- · Compare alternatives
- Reviews (237)
- Drug images
- Side effects
- Dosage information
- Patient tips
- During pregnancy
- · Generic availability
- Support group
- Drug class: skeletal muscle relaxants
- Breastfeeding

Patient resources

Other brands

Amrix, Fexmid

Professional resources

- Flexeril prescribing information
- Cyclobenzaprine (AHFS Monograph)

Other brands

Related treatment guides

- Sciatica
- Muscle Spasm

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Flexeril only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

DRUG STATUS

Availability



Pregnancy & Lactation

ন্ড Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+ years FDA approved 1977

User Reviews & Ratings

6.1 / 10

237 Reviews

Images

Flexeril 10 mg (FLEXERIL)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





