

Home

2. Doxylamine

Doxylamine do

Generic name: doxylamine [dox-IL-a-meen]

Brand names: Unisom, Nytol Maximum Strength, Aldex AN

Dosage form: oral tablet (25 mg)

Drug class: Miscellaneous anxiolytics, sedatives and hypnotics



Medically reviewed by Drugs.com on Jan 1, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is doxylamine?

Doxylamine is an antihistamine that reduces the effects of natural chemical histamine in the body. Histamine can produce symptoms of sneezing, itching, watery eyes, and runny nose. Antihistamines can cause drowsiness and are sometimes used a sleep aids.

Doxylamine is used to treat sneezing, runny nose, watery eyes, hives, skin rash, itching, and other cold or allergy symptoms.

Doxylamine is also used as a short-term treatment for sleep problems (insomnia).

Doxylamine may also be used for purposes not listed in this medication guide.

Doxylamine side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Doxylamine may cause serious side effects. Call your doctor at once if you have:

- · confusion, hallucinations;
- · severe dizziness or drowsiness; or
- little or no urinating.

Common side effects of doxylamine may include:

- blurred vision;
- dry mouth, nose, or throat;
- · constipation; or

· mild dizziness or drowsiness.

Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



Doxylamine side effects (more detail)

Related/similar drugs

Lorazepam

Lorazepam is in a group of drugs called benzodiazepines and is used to treat anxiety disorders ...

Reviews & ratings

7.9 / 10

1,054 Reviews

View more

Cephalexin

Cephalexin is a cephalosporin antibiotic used to treat bacterial infections like respiratory, skin ...

Reviews & ratings

6.4 / 10

509 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Montelukast

Montelukast is a daily oral medication used to prevent asthma attacks

exercise-induced ...

Reviews & ratings

5.6 / 10

571 Reviews

View more

Azithromycin

Azithromycin is an antibiotic used to treat many different types of infections caused by bacteria ...

Reviews & ratings

6.8 / 10

1,306 Reviews

View more

Doxycycline

Doxycycline is a tetracycline antibiotic used to treat many different bacterial infections such as ...

Reviews & ratings

6.4 / 10

1,635 Reviews

View more

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

1,404 Reviews

View more

Cetirizine

Cetirizine treats cold or allergy symptoms such as sneezing, itching, watery eyes, or runny nose ...

Reviews & ratings

View more

Warnings

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use doxylamine if you are allergic to it.

Ask a doctor or pharmacist if it is safe for you to take doxylamine if you have other medical conditions, especially:

- glaucoma;
- an enlarged prostate;
- · problems with urination; or
- asthma, bronchitis, emphysema, or other chronic lung disease.

Older adults may be more sensitive to the effects of this medicine.

Doxylamine is not expected to be harmful to an unborn baby. **Do not use this medicine without a doctor's advice if you are pregnant.**

Doxylamine may pass into breast milk and may harm a nursing baby. Antihistamines may also slow breast milk production. **Do not use this medicine without a doctor's advice if you are breast-feeding a baby.**

How should I take doxylamine?

Use exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. This medicine is usually taken only for a short time until your symptoms clear up.

Do not use doxylamine to treat insomnia in a child younger than 12 years old. Ask a doctor before using this medicine to treat cold or allergy symptoms in a child younger than 6 years old. **Always ask a doctor before giving a cold or allergy medicine to a child.** Death can occur from the misuse of these medicines in very young children.

Take this medicine with food or milk if it upsets your stomach.

To treat insomnia, it is best to take doxylamine only when you can devote several hours to sleep. Follow your doctor's instructions.

Call your doctor if your **cold or allergy symptoms** do not improve after 7 days of treatment, or if your **sleep problems** do not improve after 2 weeks of treatment.

This medication can cause you to have unusual results with allergy skin tests. Tell any doctor who treats you that you are

taking an antihistamine.

Store at room temperature away from moisture and heat.

Doxylamine dosing information

Usual Adult Dose for Insomnia:

25 mg orally once a day 30 minutes before bedtime.

Duration of Therapy: No more than 2 weeks consecutively.

Use: Helps to reduce difficulty in falling asleep.

Usual Pediatric Dose for Insomnia:

Age 12 Years and Older: 25 mg orally once a day 30 minutes before bedtime.

Duration of Therapy: No more than 2 weeks consecutively.

Use: Helps to reduce difficulty in falling asleep.

Detailed Doxylamine dosage information

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. **Do not** take extra medicine to make up the missed dose.

When treating insomnia, if it is almost your normal waking hour, skip the missed dose and wait until you are ready for bed again.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe forms of some of the side effects listed in this medication guide.

What should I avoid while taking doxylamine?

doxylamine may cause blurred vision and may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert and able to see clearly.

Ask a doctor or pharmacist before using any other cold, cough, allergy, or sleep medicine. Many combination medicines contain antihistamines. Taking certain products together can cause you to get too much of this medicine.

Ask a doctor or pharmacist before using any other cold, cough, allergy, or sleep medicine. Antihistamines are contained in many combination medicines. Taking certain products together can cause you to get too much of a certain drug. Check the label to see if a medicine contains an antihistamine.

Drinking alcohol can increase certain side effects of doxylamine.

Avoid becoming overheated or dehydrated during exercise and in hot weather. Doxylamine can decrease sweating and you may be more prone to heat stroke.

What other drugs will affect doxylamine?

Ask a doctor or pharmacist before using this medicine if you are also using any other drugs, including prescription and over-the-counter medicines, vitamins, and herbal products. Some medicines can cause unwanted or dangerous effects when used together. Not all possible interactions are listed in this medication guide.

Taking this medicine with other drugs that make you sleepy or slow your breathing can worsen these effects. Ask your doctor before taking doxylamine with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

1 Doxylamine drug interactions (more detail)

Does doxylamine interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

What are the strongest sleeping pills?

Does Mucinex help with Covid?

More about doxylamine

- Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (236)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- · Drug class: miscellaneous anxiolytics, sedatives and hypnotics

- Breastfeeding
- En español

Patient resources

Other brands

Unisom SleepTabs, Sleep Aid, Nighttime Sleepaid

Professional resources

• Doxylamine monograph

Other brands

Sleep Aid, Nighttime Sleepaid, Equate Sleep Aid, Equaline Sleep Aid

Related treatment guides

- Insomnia
- Allergic Rhinitis
- Allergies
- Conjunctivitis, Allergic

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 3.03.

DRUG STATUS

Availability RX and/or OTC Pregnancy & Lactation Risk data available CSA Schedule* N/A Not a controlled drug Approval History Drug history at FDA

User Reviews & Ratings

6.5 / 10

236 Reviews

Images

Doxylamine Succinate 25 mg (L 441)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







