

Home

Silver sulfadiazine

Silver sulfadiazine topical

Generic name: silver sulfadiazine topical [SIL-ver-SUL-fa-DYE-a-zeen-TOP-ik-al]

Brand names: Silvadene, SSD, Thermazene, SSD AF

Dosage form: topical cream (1%)
Drug class: Topical antibiotics



Medically reviewed by Drugs.com on May 27, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is silver sulfadiazine topical?

Silver sulfadiazine is an antibiotic. It fights bacteria and yeast on the skin.

Silver sulfadiazine topical (for the skin) is used to treat or prevent serious infection on areas of skin with second- or third-degree burns.

Silver sulfadiazine topical may also be used for purposes not listed in this medication guide.

Silver sulfadiazine topical side effects

Get emergency medical help if you have any of these **signs of an allergic reaction:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Although the risk of serious side effects is low when silver sulfadiazine is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream.

Silver sulfadiazine topical may cause serious side effects. Call your doctor at once if you have:

- sudden weakness or ill feeling, fever, chills, sore throat, mouth sores, red or swollen gums, trouble swallowing;
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin;
- pale or yellowed skin, dark colored urine, fever, confusion or weakness;
- kidney problems--red or pink urine, little or no urinating, swelling, rapid weight gain;
- liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes); or
- severe skin reaction--fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.

Silver sulfadiazine topical side effects (more detail)

Related/similar drugs

Lidoderm

Lidoderm is a lidocaine patch used to relieve nerve pain after shingles by numbing the skin and ...

Reviews & ratings

4.1 / 10

89 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Vancomycin

Vancomycin is used to treat infections of the intestines that cause colitis. Learn about side ...

Reviews & ratings

6.5 / 10

57 Reviews

View more

Warnings

Silver sulfadiazine topical may cause serious medical problems in a newborn if you use silver sulfadiazine topical during late pregnancy (close to your delivery date). This medicine should also not be used on premature babies or any child younger than 2 months old.

Before taking this medicine

You should not use this medicine if you are allergic to silver sulfadiazine.

Silver sulfadiazine topical may cause **serious medical problems** in a newborn if you use this medicine during late pregnancy (close to your delivery date). This medicine should also not be used on premature babies or any child younger than 2 months old.

To make sure silver sulfadiazine topical is safe for you, tell your doctor if you have:

- · liver disease;
- · kidney disease;
- a genetic enzyme deficiency called glucose-6-phosphate dehydrogenase (G6PD) deficiency; or
- an allergy to sulfa drugs.

FDA pregnancy category B. Silver sulfadiazine topical is not expected to harm an unborn baby. However, **this medicine** can cause serious medical problems in a newborn and should not be used during late pregnancy.

It is not known whether silver sulfadiazine topical passes into breast milk or if it could harm a nursing baby. You should not breast-feed while using this medicine.

How should I use silver sulfadiazine topical?

Follow all directions on your prescription label. Do not use silver sulfadiazine topical in larger or smaller amounts or for longer than recommended.

Wash your hands before and after applying silver sulfadiazine cream.

The person applying silver sulfadiazine to burn wounds should wear sterile disposable gloves. Take care to keep the treatment area as clean as possible to prevent further infection.

Clean the area to be treated as directed by your doctor. Apply enough silver sulfadiazine to cover the affected area evenly. This medicine should be applied in a layer about one 16th (1/16) of an inch thick, or 1.6 millimeters.

If needed, apply more cream to replace any medicine that has come off on bandages, clothing, or bed linens. Reapply the cream after bathing or water therapy.

Silver sulfadiazine cream is usually applied 1 or 2 times daily. Burn wounds must be kept covered with this medicine at all times. Treated skin areas can be left uncovered, or you may use a gauze bandage if directed by your doctor.

While using silver sulfadiazine topical, you may need frequent blood tests. Your kidney function may also need to be tested.

Use this medicine for the full prescribed length of time. Your symptoms may improve before the infection is completely cleared. Skipping doses may also increase your risk of further infection that is resistant to antibiotics.

Store at room temperature away from moisture and heat.

Silver sulfadiazine topical dosing information

Usual Adult Dose for Burns - External:

Apply to the affected area 1 to 2 times a day to a thickness of about 1/16 of an inch

Comments:

- -This drug should be reapplied to any areas from which it has been removed by patient activity and immediately after hydrotherapy.
- -Treatment should be continued until satisfactory healing has occurred or until burn site is ready for grafting.
- -This drug should not be withdrawn from the therapeutic regimen while there remains the possibility of infection except if a significant adverse reaction occurs.

Use: As an adjunct for the prevention and treatment of wound sepsis in patients with second and third degree burns

Usual Adult Dose for Wound Sepsis:

Apply to the affected area 1 to 2 times a day to a thickness of about 1/16 of an inch

Comments:

- -This drug should be reapplied to any areas from which it has been removed by patient activity and immediately after hydrotherapy.
- -Treatment should be continued until satisfactory healing has occurred or until burn site is ready for grafting.
- -This drug should not be withdrawn from the therapeutic regimen while there remains the possibility of infection except if a significant adverse reaction occurs.

Use: As an adjunct for the prevention and treatment of wound sepsis in patients with second and third degree burns

1 Detailed Silver sulfadiazine topical dosage information

What happens if I miss a dose?

Apply the missed dose as soon as you remember. Your burn wounds should be kept covered with silver sulfadiazine cream at all times.

What happens if I overdose?

An overdose of silver sulfadiazine topical is not expected to be dangerous. Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 if anyone has accidentally swallowed the medication.

What should I avoid while using silver sulfadiazine topical?

Do not take by mouth. Silver sulfadiazine topical is for use only on the skin. If silver sulfadiazine topical gets in your eyes, nose, or mouth, rinse with water.

What other drugs will affect silver sulfadiazine topical?

Other drugs may interact with silver sulfadiazine topical, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

0

Does silver sulfadiazine topical interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

silver sulfadiazine topical	
+	
Enter a drug name	Add

More about silver sulfadiazine topical

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (75)
- Side effects
- Dosage information
- During pregnancy
- Drug class: topical antibiotics
- En español

Patient resources

- · Silver sulfadiazine advanced reading
- Silver Sulfadiazine

Other brands

Silvadene, SSD, Thermazene, SSD AF

Professional resources

- Silver Sulfadiazine (Topical) monograph
- Silver Sulfadiazine Cream (FDA)

Other brands

Silvadene, SSD

Related treatment guides

· Burns, External

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 3.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্য Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

9.8 / 10

75 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×