

Home

2. Altace

Altace do

Generic name: ramipril [RAM-i-pril]

Drug class: Angiotensin Converting Enzyme Inhibitors

Medically reviewed by Sanjai Sinha, MD. Last updated on Apr 22, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is Altace?

Altace is an ACE inhibitor. ACE stands for angiotensin converting enzyme.

Altace is used to treat high blood pressure (hypertension). Lowering blood pressure may lower your risk of a stroke or heart attack.

Altace is also used to treat congestive heart failure.

Warnings

Do not Altace use if you are pregnant. If you become pregnant, stop taking this medicine and tell your doctor right away.

You should not use Altace if you have ever had angioedema. Do not take ramipril within 36 hours before or after taking medicine that contains sacubitril (such as Entresto).

If you have diabetes, do not use ramipril together with any medication that contains aliskiren (a blood pressure medicine).

Before taking this medicine

You should not use Altace if you are allergic to ramipril, or if:

- · you recently took a heart medicine called sacubitril; or
- you have ever had a severe allergic reaction to any ACE inhibitor (benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, or trandolapril).

Do not take Altace within 36 hours before or after taking medicine that contains sacubitril (such as Entresto).

If you have diabetes, do not use ramipril together with any medication that contains aliskiren (a blood pressure medicine).

You may also need to avoid taking ramipril with aliskiren if you have kidney disease.

To make sure Altace is safe for you, tell your doctor if you have ever had:

- kidney disease (or if you are on dialysis);
- an electrolyte imbalance (such as high levels of potassium in your blood);
- · liver disease:
- a connective tissue disease such as Marfan syndrome, Sjogren's syndrome, lupus, scleroderma, or rheumatoid arthritis;
- if you are on a low-salt diet; or
- if you are also taking telmisartan (Micardis).

Do not use if you are pregnant. If you become pregnant, stop taking this medicine and tell your doctor right away. Altace can cause injury or death to the unborn baby if you take the medicine during your second or third trimester.

You should not breast-feed while you are using ramipril.

Altace is not approved for use by anyone younger than 18 years old.

Altace pregnancy and breastfeeding warnings (more detail)

How should I take Altace?

Take Altace exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Swallow the tablet whole.

Your blood pressure will need to be checked often, and you may need blood tests.

Call your doctor if you have ongoing vomiting or diarrhea, or if you are sweating more than usual. You can easily become dehydrated while taking Altace. This can lead to very low blood pressure, electrolyte disorders, or kidney failure.

If you need surgery, tell the surgeon ahead of time that you are using ramipril.

If you are being treated for high blood pressure, keep using this medicine even if you feel well. High blood pressure often has no symptoms. You may need to use blood pressure medication for the rest of your life.

Store at room temperature away from moisture, heat, and light. Keep the bottle tightly closed when not in use.

Dosing information

Usual Adult Dose for Diabetic Nephropathy:

Initial dose: 2.5 mg orally once a day for patients not receiving a diuretic Maintenance dose: 2.5 to 20 mg/day orally in 1 to 2 divided doses

Usual Adult Dose for Hypertension:

Initial dose: 2.5 mg orally once a day for patients not receiving a diuretic

Maintenance dose: 2.5 to 20 mg/day orally in 1 to 2 divided doses

Usual Adult Dose for Congestive Heart Failure:

Initial dose: 2.5 mg orally twice a day Maintenance dose: 5 mg orally twice a day

Usual Adult Dose for Left Ventricular Dysfunction:

Initial dose: 2.5 mg orally twice a day

Maintenance dose: 5 mg orally twice a day

Usual Adult Dose for Myocardial Infarction:

Initial dose: 2.5 mg orally twice a day

Maintenance dose: 5 mg orally twice a day

Detailed Altace dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What to avoid

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

Do not use potassium supplements or salt substitutes, unless your doctor has told you to.

Altace side effects

Get emergency medical help if you have **signs of an allergic reaction to Altace:** hives; severe stomach pain; difficulty breathing; swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American.

Call your doctor at once if you have:

- a light-headed feeling, like you might pass out;
- jaundice (yellowing of your skin or eyes);
- little or no urination;
- fever, chills, sore throat; or
- high potassium nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement.

Common Altace side effects may include:

- · headache;
- · cough; or
- dizziness, weakness, tired feeling.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Altace side effects (more detail)

Related/similar drugs

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

FEATURED

Wegovy

Wegovy is a once-weekly semaglutide injection used for chronic weight management and to reduce ...

Reviews & ratings

7.6 / 10

478 Reviews

View more

Jardiance

Jardiance (empagliflozin) is used to treat type 2 diabetes, chronic kidney disease and reduce the ...

Reviews & ratings

5.3 / 10

325 Reviews

View more

Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

Reviews & ratings

6.4 / 10

56 Reviews

View more

Propranolol

Propranolol is a beta-blocker that is used to treat tremors, chest pain, high blood pressure, heart ...

Reviews & ratings

8.2 / 10

1,376 Reviews

View more

Hydrochlorothiazide

HCTZ (hydrochlorothiazide) used to treat high blood pressure (hypertension) and edema. Includes ...

Reviews & ratings

5.2 / 10

185 Reviews

View more

Spironolactone

Spironolactone is a potassium-sparing

diuretic that is primarily used to treat heart failure, high ...

Reviews & ratings

6.9 / 10

702 Reviews

View more

Carvedilol

Carvedilol (Coreg) is used to treat heart failure and hypertension (high blood pressure). Includes ...

Reviews & ratings

5.3 / 10

201 Reviews

View more

What other drugs will affect Altace?

Tell your doctor about all your current medicines. Many drugs can interact with Altace, especially:

- a diuretic ("water pill"), or any other blood pressure medications;
- lithium,
- gold injections to treat arthritis;
- insulin or oral diabetes medicine; or
- NSAIDs (nonsteroidal anti-inflammatory drugs) aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others.

This list is not complete and many other drugs may interact with Altace. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Altace drug interactions (more detail)

Does Altace interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

+
Enter a drug name
Add

Frequently asked questions

• Do ACE inhibitors make COVID-19 worse?

More about Altace (ramipril)

- Check interactions
- · Compare alternatives
- · Pricing & coupons
- Reviews (12)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- · Generic availability
- Drug class: Angiotensin Converting Enzyme Inhibitors
- Breastfeeding
- En español

Professional resources

- Altace prescribing information
- Ramipril (AHFS Monograph)

Related treatment guides

- High Blood Pressure
- Cardiovascular Risk Reduction
- Diabetic Kidney Disease
- Heart Attack
- Heart Failure
- Left Ventricular Dysfunction

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Altace only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 15.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্প Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+ years FDA approved 1991

User Reviews & Ratings

6.9 / 10

12 Reviews

Images

Altace 10 mg (ALTACE 10mg MP)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

X