

Home

2. Prednisolone

Prednisolone 40

Generic name: prednisolone [pred-NIS-oh-lone]

Brand names: Flo-Pred, Millipred, Orapred, Pediapred, Veripred 20, ... show all 15 brands **Dosage forms:** oral liquid, oral suspension, oral syrup, oral tablet, oral disintegrating tablet

Drug class: Glucocorticoids

Medically reviewed by Carmen Pope, BPharm. Last updated on Aug 7, 2023.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is prednisolone?

Prednisolone is a corticosteroid that may be used to reduce inflammation and calm down an overactive immune system. It works by mimicking the effects of cortisol, a hormone released by the adrenal glands (located on top of the kidneys) that regulates metabolism and stress. Prednisolone prevents the release of substances in the body that cause inflammation.

Prednisolone is used to treat many different inflammatory conditions such as arthritis, lupus, psoriasis, ulcerative colitis, allergic disorders, gland (endocrine) disorders, and conditions that affect the skin, eyes, lungs, stomach, nervous system, or blood cells.

Prednisolone has predominantly glucocorticoid activity, which means it mainly affects our immune response and reduces inflammation, rather than affecting the body's balance of electrolytes and water (this is called mineralocorticoid activity).

Prednisolone was FDA approved in 1955.

Warnings

You should not use prednisolone if you have a fungal infection anywhere in your body.

Before taking this medicine

You should not use prednisolone if you are allergic to it, or if you have:

• a fungal infection anywhere in your body.

Prednisolone can weaken your immune system, making it easier for you to get an infection. Steroids can also worsen an infection you already have, or reactivate an infection you recently had. Tell your doctor about any illness or infection you have had within the past several weeks.

To make sure prednisolone is safe for you, tell your doctor if you have ever had:

- active tuberculosis;
- a thyroid disorder;
- herpes infection of the eyes;
- stomach ulcers, ulcerative colitis, or diverticulitis;
- depression, mental illness, or psychosis;
- liver disease (especially cirrhosis);
- high blood pressure;
- · osteoporosis;
- a muscle disorder such as myasthenia gravis; or
- multiple sclerosis.

Also tell your doctor if you have diabetes. Steroid medicines may increase the glucose (sugar) levels in your blood or urine. You may also need to adjust the dose of your diabetes medications.

It is not known whether prednisolone will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether prednisolone passes into breast milk or if it could affect the nursing baby. Tell your doctor if you are breast-feeding.

1 Prednisolone pregnancy and breastfeeding warnings (more detail)

How should I take prednisolone?

Take prednisolone exactly as prescribed by your doctor. Your doctor may occasionally change your dose. Do not use prednisolone in larger or smaller amounts or for longer than recommended.

Prednisolone is sometimes taken every other day. Follow your doctor's dosing instructions very carefully.

Measure liquid medicine with the dosing syringe provided, or with a special dose-measuring spoon or medicine cup. If you do not have a dose-measuring device, ask your pharmacist for one.

You may need to shake the **oral suspension** (liquid) well just before you measure a dose. Follow the directions on your medicine label.

Keep prednisolone disintegrating tablets in their blister pack until you are ready to take the medicine. Open the package using dry hands, and peel back the foil from the tablet blister (do not push the tablet through the foil). Remove the tablet and place it in your mouth. Allow the disintegrating tablet to dissolve in your mouth without chewing. Swallow several times as the tablet dissolves. If desired, you may drink liquid to help swallow the dissolved tablet.

Your dose needs may change if you have unusual stress such as a serious illness, fever or infection, or if you have surgery or a medical emergency. Tell your doctor about any such situation that affects you.

Prednisolone can cause unusual results with certain medical tests. Tell any doctor who treats you that you are using prednisolone.

You should not stop using prednisolone suddenly. Follow your doctor's instructions about tapering your dose.

Wear a medical alert tag or carry an ID card stating that you take prednisolone. Any medical care provider who treats you should know that you take steroid medication.

If you need surgery, tell the surgeon ahead of time that you are using prednisolone. You may need to stop using the medicine for a short time.

Store prednisolone at room temperature away from moisture and heat.

Detailed Prednisolone dosage information

What happens if I miss a dose?

Call your doctor for instructions if you miss a dose of prednisolone.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

An overdose of prednisolone is not expected to produce life threatening symptoms. However, long term use of high steroid doses can lead to symptoms such as thinning skin, easy bruising, changes in the shape or location of body fat (especially in your face, neck, back, and waist), increased acne or facial hair, menstrual problems, impotence, or loss of interest in sex.

What should I avoid while taking prednisolone?

Do not receive a "live" vaccine while using prednisolone. The vaccine may not work as well during this time, and may not fully protect you from disease. Live vaccines include measles, mumps, rubella (MMR), polio, rotavirus, typhoid, yellow fever, varicella (chickenpox), zoster (shingles), and nasal flu (influenza) vaccine.

Do not receive a smallpox vaccine or you could develop serious complications.

Avoid being near people who are sick or have infections. Call your doctor for preventive treatment if you are exposed to chickenpox or measles. These conditions can be serious or even fatal in people who are using steroid medication.

Prednisolone side effects

Get emergency medical help if you have **signs of an allergic reaction to prednisolone**.: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Prednisolone may cause serious side effects. Call your doctor at once if you have:

- shortness of breath (even with mild exertion), swelling, rapid weight gain;
- bruising, thinning skin, or any wound that will not heal;
- · severe depression, changes in personality, unusual thoughts or behavior;
- new or unusual pain in an arm or leg or in your back;

- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- · severe pain in your upper stomach spreading to your back, nausea and vomiting;
- a seizure (convulsions); or
- low potassium--leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling.

Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using prednisolone.

Common side effects of prednisolone may include:

- fluid retention (swelling in your hands or ankles);
- · dizziness, spinning sensation;
- · changes in your menstrual periods;
- · headache:
- · muscle pain or weakness; or
- stomach discomfort, bloating.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Prednisolone side effects (more detail)

Related/similar drugs

Kesimpta

Kesimpta (ofatumumab) is used to treat relapsing forms of multiple sclerosis. Includes Kesimpta ...

Reviews & ratings

6.8 / 10

74 Reviews

View more

Stelara

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

Reviews & ratings

7.6 / 10

252 Reviews

View more

FEATURED

Skyrizi

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis, ulcerative colitis ...

Reviews & ratings

7.9 / 10

56 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Loratadine

Loratadine is a non-drowsy antihistamine used to relieve allergy symptoms like sneezing, runny ...

Reviews & ratings

4.7 / 10

139 Reviews

View more

Methylprednisolone

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

Reviews & ratings

7.1 / 10

484 Reviews

View more

Methotrexate

Methotrexate is used to treat certain types of cancer of the breast, skin, head and neck, or lung ...

Reviews & ratings

6.5 / 10

335 Reviews

View more

Triamcinolone

Triamcinolone is used to treat allergies, skin conditions, ulcerative colitis, and arthritis. Learn ...

Reviews & ratings

6.7 / 10

226 Reviews

View more

What other drugs will affect prednisolone?

Other drugs may interact with prednisolone, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

Prednisolone drug interactions (more detail)

Does prednisolone interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

prednisolone
+
Enter a drug name
Add

Popular FAQ

Is prednisolone best taken in the morning and why?

More FAQ

• Prednisone vs Prednisolone - What's the difference?

More about prednisolone

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (48)
- Drug images
- · Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: glucocorticoids
- Breastfeeding
- En español

Patient resources

- Prednisolone Oral Solution patient information
- Prednisolone Orally Disintegrating Tablets

Other brands

Prelone, Millipred, Bubbli-Pred, Flo-Pred, ... +5 more

Professional resources

- PrednisoLONE, prednisoLONE Acetate, prednisoLONE Sodium Phosphate monograph
- Prednisolone ODT (FDA)
- Prednisolone Oral Solution (FDA)
- Prednisolone Sodium Phosphate Oral Solution (FDA)
- Prednisolone Syrup (FDA)
- Prednisolone tablets (FDA)

Other brands

Orapred, Millipred, Flo-Pred, Orapred ODT, ... +2 more

Related treatment guides

- · Asthma, acute
- Bronchopulmonary Dysplasia
- Allergies
- Dermatitis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use prednisolone only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 9.01.

DRUG STATUS

Availability
Rx Prescription only
Pregnancy & Lactation
Risk data available
CSA Schedule*
N/A Not a controlled drug
Approval History
Drug history at FDA

WADA Class ດິດ Anti-Doping Classification

User Reviews & Ratings

6.4 / 10

48 Reviews

Images

Prednisolone Sodium Phosphate (Orally Disintegrating) 10 mg (base) (M P10)



spriate (eraily biomograting) to mg (bace) (in the

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

X