

Home

Radium 223 dichloride

Radium Ra 223 dichloride

Generic name: radium Ra 223 dichloride [RAY-dee-um-Ra-223 dye-KLOR-ide]

Brand name: Xofigo

Dosage form: intravenous solution (-) **Drug class:** Therapeutic radiopharmaceuticals



Medically reviewed by Drugs.com on Mar 7, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is radium Ra 223 dichloride?

Radium Ra 223 dichloride is used to treat prostate cancer that has spread to the bones and cannot be treated with surgery.

Radium Ra 223 dichloride may also be used for purposes not listed in this medication guide.

Radium Ra 223 dichloride side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Radium Ra 223 dichloride may cause serious side effects. Call your doctor at once if you have:

- easy bruising, unusual bleeding, purple or red spots under your skin;
- a light-headed feeling, like you might pass out;
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin;
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath;
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet; or
- low white blood cell counts--fever, mouth sores, skin sores, sore throat, cough, trouble breathing.

Common side effects of radium Ra 223 dichloride may include:

- nausea, vomiting, diarrhea;
- · low blood cell counts; or
- · swelling in your hands or feet.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

• Radium 223 dichloride side effects (more detail)

Related/similar drugs

Premarin

Premarin is used to treat symptoms of menopause such as hot flashes, and vaginal dryness. Learn ...

Reviews & ratings

7.7 / 10

84 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Estrace

Estrace (estradiol) is used to treat symptoms of menopause such as hot flashes and vaginal dryness ...

Reviews & ratings

7.3 / 10

12 Reviews

View more

Xtandi

Xtandi (enzalutamide) is an antiandrogen used to treat prostate prostate cancer. Includes Xtandi ...

Reviews & ratings

View more

Zytiga

Zytiga (abiraterone) is used together with prednisone to treat prostate cancer that has spread to ...

Reviews & ratings

8.2 / 10

41 Reviews

View more

Lynparza

Lynparza (olaparib) is used to treat ovarian cancer, breast cancer, pancreatic cancer, and prostate ...

Reviews & ratings

7.4 / 10

17 Reviews

View more

Casodex

Casodex (bicalutamide) is an antiandrogen and is used together with LHRH to treat prostate cancer ...

Reviews & ratings

7 Reviews

View more

Bicalutamide

Bicalutamide is a prescription medicine used to treat prostate cancer; Includes bicalutamide side ...

Reviews & ratings

8.4 / 10

13 Reviews

View more

Warnings

radium Ra 223 dichloride can increase your risk of bleeding or infection. You will need frequent medical tests.

Before taking this medicine

Radium Ra 223 dichloride can harm an unborn baby or cause birth defects **if you father a child while using this medicine**. Use condoms while you are being treated with this medicine and for at least 6 months after treatment ends.

Your sex partner should also use effective birth control to prevent pregnancy while you are using this medicine and for at least 6 months after your last dose. **Tell your doctor right away if a pregnancy occurs during this time.**

This medicine may affect fertility (your ability to have children). However, it is important to use birth control to prevent pregnancy because radium Ra 223 dichloride can harm an unborn baby.

How is Ra 223 dichloride given?

Radium Ra 223 dichloride is given as an infusion into a vein. A healthcare provider will give you this injection.

Radium Ra 223 dichloride is usually given once every 4 weeks for a total of 6 injections. Your doctor will determine how long to treat you with radium Ra 223 dichloride.

Radium Ra 223 dichloride can lower your blood cell counts. **Your blood will need to be tested often.** Your cancer treatments may be delayed based on the results.

After you are treated with radium Ra 223 dichloride, your body fluids (urine, stools, vomit) will contain radioactive materials. Use a toilet rather than a urinal and sit on the toilet while urinating. Flush the toilet several times with the lid down after use.

Always wash your hands after using the bathroom. **Avoid handling any clean-up of your stools or urine without wearing latex rubber gloves.** If another person is handling your stools or urine, they should wear rubber gloves, disposable medical gown, eye protection, and a mask to cover the nose and mouth.

When cleaning any spills of bodily fluid, use only disposable cleaning cloths that can be flushed down a toilet. Ask your doctor or health department how to dispose of any bodily fluid spills that cannot be flushed down a toilet.

Wash any soiled clothing separately from the laundry of other people in your home.

Radium Ra 223 dichloride doses are based on weight. Your dose needs may change if you gain or lose weight.

Drink plenty of water each day to keep from getting dehydrated and to keep your kidneys working properly.

Radium Ra 223 dichloride dosing information

Usual Adult Dose for Prostate Cancer:

55 kilobecquerel (kBq) or 1.49 microcurie, per kg body weight, IV, every 4 weeks

Duration of therapy: 6 doses

Comments: See manufacturer's product information for complete dosing instructions including decay correction factor table and calculation of correct volume for dosing.

Use: The treatment of patients with castration-resistant prostate cancer, symptomatic bone metastases, and no known visceral metastatic disease.

Detailed Radium 223 dichloride dosage information

What happens if I miss a dose?

Call your doctor for instructions if you miss an appointment for your radium Ra 223 dichloride injection.

What happens if I overdose?

Since radium Ra 223 dichloride is given by a healthcare professional in a medical setting, an overdose is unlikely to occur.

What should I avoid while receiving radium Ra 223 dichloride?

The radioactivity in this medicine can pass into body fluids (urine, feces, vomit). For at least 7 days after you receive a dose, avoid allowing your body fluids to come into contact with your hands or other surfaces. Caregivers should wear rubber gloves and a disposable medical gown while cleaning up a patient's body fluids, handling contaminated trash or laundry or changing diapers. Wash hands before and after removing gloves. Wash soiled clothing and linens separately from other laundry.

What other drugs will affect radium Ra 223 dichloride?

You may have an increased risk of bone fractures or death if you are treated with radium Ra 223 dichloride while you are also taking another medicine for prostate cancer called abiraterone (Yonsa, Zytiga) in combination with prednisone or prednisolone (steroid medicine).

Other drugs may affect radium Ra 223 dichloride, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

Radium 223 dichloride drug interactions (more detail)

Does radium 223 dichloride interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

+ Enter a drug name Add

More about radium 223 dichloride

- · Check interactions
- Compare alternatives
- Reviews (3)
- Side effects
- Dosage information
- During pregnancy
- Drug class: therapeutic radiopharmaceuticals
- Breastfeeding
- En español

Patient resources

• Radium ra 223 dichloride advanced reading

Other brands

Xofigo

Professional resources

Other brands

Xofigo

Related treatment guides

Prostate Cancer

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 1.02.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation Risk data available CSA Schedule* N/A Not a controlled drug Approval History □ Drug history at FDA

User Reviews & Ratings

3 Reviews

Images

Xofigo (radium 223 dichloride) 1100 kBq/mL (30 microcurie/mL) with a total radioactivity of 6,600 kBq/vial (178 microcurie/vial)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









