



[Home](#)

2. [Moderiba](#)

Moderiba

Generic name: [ribavirin](#) (oral) [*RYE-ba-VYE-rin*]

Brand names: [Copegus](#), [Moderiba](#), [RibaPak](#), [RibaTab](#)

Drug class: [Purine nucleosides](#)



[Medically reviewed](#) by Drugs.com on Jul 31, 2023. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)



The [Moderiba](#) brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be [generic equivalents available](#).

What is Moderiba?

Moderiba is an antiviral medication that is used to treat [chronic hepatitis C](#).

Moderiba is not effective when used alone. Moderiba must be used in combination with interferon alfa or peginterferon alfa.

Moderiba is sometimes given to people taking other antiviral medications to treat [hepatitis C](#).

Moderiba may also be used for purposes not listed in this medication guide.

Moderiba side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Moderiba can cause anemia. In rare cases, this can lead to fatal heart problems. **Get emergency medical attention if you have chest pain.**

Moderiba may cause serious side effects. Call your doctor at once if you have:

- problems with your vision;
- severe pain in your upper stomach spreading to your back, nausea, vomiting, diarrhea;
- new or worsening cough, fever, stabbing chest pain, [wheezing](#), feeling short of breath;
- severe [depression](#), thoughts about suicide, or thoughts about hurting someone else;
- **signs of serious anemia**--pale or yellowed skin, dark colored urine, confusion or weakness; or

- **other signs of low blood cell counts**--fever, chills, flu-like symptoms, swollen gums, mouth sores, skin sores, easy bruising, unusual bleeding, feeling light-headed.

Moderiba can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine.

Common side effects of Moderiba may include:

- nausea, vomiting, loss of appetite;
- fever, chills or shaking;
- low blood cell counts, anemia;
- feeling weak or tired;
- [headache](#), [muscle pain](#); or
- mood changes, feeling anxious or irritable.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Moderiba side effects](#) (more detail)

Related/similar drugs

Epclusa

Epclusa treats chronic hepatitis C in adults and children 3+. This once-daily antiviral combines ...

Reviews & ratings

7.5 / 10

[397 Reviews](#)

[View more](#)

Harvoni

Harvoni is used to treat hepatitis C virus (HCV) infections in adults and children aged 3 years and ...

Reviews & ratings

8.4 / 10

[403 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Mavyret

Mavyret (glecaprevir/pibrentasvir) is an oral combination medicine containing fixed doses of two ...

Reviews & ratings

8.3 / 10

[426 Reviews](#)

[View more](#)

Sovaldi

Sovaldi (sofosbuvir) is used to treat chronic hepatitis C virus (HCV) infection. Includes Sovaldi ...

Reviews & ratings

7.6 / 10

[47 Reviews](#)

[View more](#)

Pegasys

Pegasys is used to treat chronic hepatitis B or C. Learn about side effects, interactions and ...

Reviews & ratings

6.6 / 10

[14 Reviews](#)

[View more](#)

Vosevi

Vosevi (sofosbuvir, velpatasvir and voxilaprevir) is used to treat chronic hepatitis C. Includes ...

Reviews & ratings

[2 Reviews](#)

[View more](#)

Glecaprevir/pibrentasvir

Glecaprevir/pibrentasvir (Mavyret) is an oral combination medicine containing fixed doses of 2 ...

Reviews & ratings

8.3 / 10

[428 Reviews](#)

[View more](#)

Sofosbuvir/velpatasvir

Sofosbuvir-velpatasvir (brand name Epclusa) is used to treat chronic hepatitis C (HCV) genotypes 1 ...

Reviews & ratings

7.6 / 10

[417 Reviews](#)

[View more](#)

Warnings

You should not use Moderiba if you have [autoimmune hepatitis](#), moderate to severe kidney disease, a hemoglobin blood cell disorder, or if you take [didanosine](#).

Moderiba can harm an unborn baby if the mother or the father is using Moderiba. **Do not use if you are pregnant, or if you are a man whose sex partner is pregnant.** Both men and women should use effective [birth control](#) while taking this medicine and for at least 6 months after the last dose.

Moderiba is not effective when used alone and must be used in combination with an interferon.

In rare cases, Moderiba can cause fatal heart problems. **Get emergency medical attention if you have chest pain.**

Before taking this medicine

You should not take Moderiba if you are allergic to it, or if you have:

- a hemoglobin blood cell disorder such as sickle-cell anemia or [thalassemia](#);
- autoimmune hepatitis;

- moderate to severe kidney disease;
- if you are also taking didanosine; or
- if you are pregnant, or if you are a man whose sex partner is pregnant.

When you take Moderiba in combination with other medications: There may be other reasons you should not take the combination treatment. Tell your doctor about all your medical conditions.

Tell your doctor if you have ever had:

- a blood cell disorder such as anemia (low red blood cells);
- heart problems, [high blood pressure](#), or [heart attack](#);
- breathing problems;
- vision problems (you may need an eye exam before taking Moderiba);
- diabetes;
- human [immunodeficiency](#) virus (HIV or AIDS) or a weak immune system;
- depression, mental illness, thoughts of hurting yourself or others;
- a thyroid disorder;
- kidney disease;
- an [organ transplant](#);
- liver problems other than hepatitis C; or
- treatment for hepatitis C that did not work well.

Both men and women taking Moderiba should use effective birth control to prevent pregnancy. This medicine can cause birth defects, miscarriage, or death to an unborn baby if the mother or father is using this medicine.

- **If you are a woman**, do not take Moderiba if you are pregnant. You will need to have a negative pregnancy test before taking this medicine. You will also need pregnancy tests during treatment and for 6 months after your last dose.
- **If you are a man**, do not take Moderiba if your sex partner is pregnant.
- Both men and women should keep using effective birth control for at least 6 months after the last dose of Moderiba.
- Tell your doctor right away if a pregnancy occurs while either the mother or the father is taking Moderiba. Your name may be listed on a pregnancy registry to track the effects of Moderiba on the baby.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk.

Moderiba is not approved for use by anyone younger than 3 years old. This medicine **tablets** are not approved for use by anyone younger than 5 years old.

How should I take Moderiba?

Moderiba is not effective when used alone and must be used in combination with other medications to treat hepatitis C.

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Take Moderiba with food.

Swallow the capsule whole and do not crush, chew, break, or open it. **Tell your doctor if you have trouble swallowing the capsule.**

Measure **liquid medicine** carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

Drink extra fluids while you are taking Moderiba to keep from getting dehydrated.

Moderiba can cause [dry mouth](#), which could lead to tooth decay or [gum disease](#). **If you vomit while taking this medicine, rinse your mouth out with water to prevent damage to your teeth or gums.** Be sure to have regular dental exams while you are taking this medication.

You will need frequent medical tests.

Store Moderiba **tablets** or **capsules** at room temperature away from moisture, heat, and light.

Store Moderiba **liquid** (oral solution) at room temperature or in the refrigerator.

 [Detailed Moderiba dosage information](#)

What happens if I miss a dose?

Take the missed dose on the same day you remember it. Skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

Get your prescription refilled before you run out of medicine completely.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Moderiba?

Avoid drinking alcohol. It can make your liver problems worse.

Using Moderiba will not prevent you from passing hepatitis to other people. Follow your doctor's instructions about how to prevent passing the disease to another person.

What other drugs will affect Moderiba?

Tell your doctor about all your other medicines, especially:

- [azathioprine](#); or
- medicines to treat HIV or AIDS.

This list is not complete. Other drugs may affect Moderiba, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

 [Moderiba drug interactions](#) (more detail)

Does Moderiba interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Moderiba

+

Enter a drug name

Add

More about Moderiba (ribavirin)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(1\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: purine nucleosides](#)
- [Breastfeeding](#)

Patient resources

Other brands

[Copegus](#), [Ribasphere](#), [Virazole](#), [Rebetol](#), ... +2 more

Professional resources

- [Moderiba prescribing information](#)
- [Ribavirin \(AHFS Monograph\)](#)

Other brands

[Copegus](#), [Virazole](#), [Rebetol](#)

Related treatment guides

- [Hepatitis C](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.01.

DRUG STATUS

Availability

 Discontinued

Pregnancy & Lactation

 Risk data available

CSA Schedule*

N/A Not a controlled drug



User Reviews & Ratings

[Review this drug](#)

Images

Moderiba 400 mg (3RP 400)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.