

Home

2. Myobloc

Myobloc

Generic name: rimabotulinumtoxinB [RIM-a-BOT-ue-LYE-num-TOX-in-BEE]

Drug class: Skeletal muscle relaxants



<u>Medically reviewed</u> by Drugs.com on Dec 9, 2024. Written by <u>Cerner Multum</u>.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is Myobloc?

Myobloc, also called botulinum toxin type B, is made from the bacteria that causes botulism. Botulinum toxin blocks nerve activity in the muscles, causing a temporary reduction in muscle activity.

Myobloc is used in adults to treat abnormal head position and neck pain due from cervical dystonia.

Myobloc is also used in adults to treat chronic drooling (sialorrhea).

Myobloc may also be used for purposes not listed in this medication guide.

Myobloc side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives, difficult breathing, swelling of your face, lips, tongue, or throat.

The botulinum toxin contained in Myobloc can spread to other body areas beyond where it was injected. This has caused side effects in some people after receiving botulinum toxin injections, even for cosmetic purposes. Swallowing and breathing problems can be serious life-threatening, especially in children with certain muscle problems.

Call your doctor at once if you have any of these side effects (up to several hours or weeks after an injection):

- unusual or severe muscle weakness (especially in a body area that was not injected with the medication);
- trouble breathing, talking, or swallowing;
- hoarse voice, droopy eyelids, blurred vision, double vision, or other vision problems; or
- · loss of bladder control.

Common side effects of Myobloc may include:

- pain or discomfort where the medicine was injected;
- · headache;

- dry mouth; or
- · trouble swallowing.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Myobloc side effects (more detail)

Related/similar drugs

Botox

Botox is used to treat chronic migraines, excessive sweating, bladder conditions, eye muscle ...

Reviews & ratings

5.7 / 10

473 Reviews

View more

Xeomin

Xeomin (incobotulinumtoxinA) is used to treat cervical dystonia, blepharospasm, upper facial lines ...

Reviews & ratings

5 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Dysport

Dysport (abobotulinumtoxinA) is used to treat cervical dystonia, glabellar lines and limb ...

Reviews & ratings

4.7 / 10

9 Reviews

View more

Botox Cosmetic

Botox Cosmetic temporarily improves the look of moderate to severe frown lines, crow's feet ...

Reviews & ratings

4.9 / 10

12 Reviews

View more

Daxxify

Daxxify (daxibotulinumtoxinA-lanm) is used to improve frown lines and for the treatment of cervical ...

Reviews & ratings

3 Reviews

View more

OnabotulinumtoxinA

OnabotulinumtoxinA information from Drugs.com, includes OnabotulinumtoxinA side effects ...

Reviews & ratings

5.7 / 10

493 Reviews

View more

Warnings

The botulinum toxin contained in Myobloc can spread to other body areas beyond where it was injected. This can cause serious life-threatening side effects.

Call your doctor at once if you have a hoarse voice, droopy eyelids, vision problems, severe muscle weakness, loss of bladder control, or trouble breathing, talking, or swallowing. **Some of these effects can occur up to several weeks after a botulinum toxin injection.**

Before taking this medicine

You should not be treated with Myobloc if you are allergic to it or to any other botulinum toxin injection, such as onabotulinumtoxinA, abobotulinumtoxinA, or incobotulinumtoxinA, or if you have an infection in the area where the medicine will be injected.

Tell your doctor if you have ever had a side effect after receiving a botulinum toxin.

Tell your doctor if you have or have ever had:

- a nerve-muscle disorder, such as myasthenia gravis, ALS (Lou Gehrig's disease), or Lambert-Eaton syndrome;
- a breathing disorder or problems such as asthma or emphysema;
- · droopy eyelids;
- problems with swallowing;
- bleeding problems;
- surgery on your face;
- a planned surgery; or
- received other botulinum toxin injection in the past 4 months.

Myobloc is made from donated human plasma and may contain viruses or other infectious agents. Donated plasma is tested and treated to reduce the risk of contamination, but there is still a small possibility it could transmit disease. Ask your doctor about any possible risk.

It is not known if Myobloc will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

Ask a doctor if it is safe to breastfeed while using Myobloc.

How is Myobloc given?

Myobloc is injected into a muscle or into a gland by a healthcare provider. Your doctor will determine when to receive the next dose.

Your injections may be given into more than one area at a time, depending on the condition being treated.

Your doctor may occasionally change your dose until your doctor determines the best dose.

Your doctor may need to check your progress on a regular basis, and your next dose may change based on your response to treatment.

Call your doctor if your symptoms do not improve, or if they get worse.

Do not receive botulinum toxin injections from more than one medical professional at a time. If you switch healthcare providers, be sure to tell your new provider how long it has been since your last botulinum toxin injection.

Using this medicine more often than prescribed will not make it more effective and may result in serious side effects.

What happens if I miss a dose?

Call your doctor for instructions if you miss an appointment for your Myobloc.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may not appear right away, but can include muscle weakness, trouble swallowing, weak or shallow breathing, or loss of movement in any part of your body.

What should I avoid after receiving Myobloc?

Myobloc may impair your reactions and may cause weakness, loss of strength, or vision changes. Avoid driving or hazardous activity until you know how Myobloc will affect you.

What other drugs will affect Myobloc?

Tell your doctor about all your other medicines, especially:

- a muscle relaxer;
- an injectable antibiotic;
- · sleep medicine;
- botulinum toxin such as Dysport, Xeomin, Botox, Botox cosmetic;
- cold or allergy medicine (Benadryl and others);
- medicine to treat Parkinson's disease;
- medicine to treat stomach problems, motion sickness, or irritable bowel syndrome;
- medicine to treat overactive bladder; or
- · bronchodilator asthma medication.

This list is not complete and many other drugs may affect Myobloc. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Myobloc drug interactions (more detail)

Does Myobloc interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Myobloc
+

Enter a drug name

Myobloc Biosimilars

Biosimilar and interchangeable products are biological products that are highly similar to and have no clinically meaningful differences from the reference product.

Reference products

These are biological products that have already been approved by the FDA, against which biosimilar products are compared. There is 1 for Myobloc.

Myobloc (rimabotulinumtoxinB) - Solstice NeuroSciences, LLC

Formulation type	Strength
Single-Dose Vial	10,000 units/2 mL (5,000 units/mL)
Single-Dose Vial	2,500 units/0.5 mL
Single-Dose Vial	5,000 units/mL

Frequently asked questions

• What is botulinum toxin used to treat?

More about Myobloc (rimabotulinumtoxinB)

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (3)
- Side effects
- Dosage information
- During pregnancy
- FDA approval history
- Drug class: skeletal muscle relaxants
- Breastfeeding
- En español

Patient resources

Myobloc advanced reading

Professional resources

- Myobloc prescribing information
- RimabotulinumtoxinB (AHFS Monograph)

Related treatment guides

- Sialorrhea
- Cervical Dystonia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 4.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

3 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

×