



[Home](#)

2. Triamcinolone

# Triamcinolone

**Generic name:** triamcinolone (oral) [ *trye-am-SIN-oh-lone* ]

**Brand name:** [Aristocort](#)

**Drug class:** [Glucocorticoids](#)

Medically reviewed by [Melisa Puckey, BPharm](#). Last updated on Mar 1, 2024.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

## What is triamcinolone?

Triamcinolone belongs to a class of drugs called corticosteroids. It prevents the release of substances in the body that cause inflammation.

Triamcinolone oral (taken by mouth) is used to treat many different conditions such as allergic disorders, skin conditions, ulcerative colitis, arthritis, lupus, psoriasis, or breathing disorders.

Triamcinolone may also be used for purposes not listed in this medication guide.

## Warnings

You should not use this medication if you are allergic to triamcinolone, or if you have a fungal infection anywhere in your body.

Before taking triamcinolone, tell your doctor about all of your medical conditions, and about all other medicines you are using. There are many other disease that can be affected by steroid use, and many other medicines that can interact with steroids.

Your steroid medication needs may change if you have any unusual stress such as a serious illness, fever or infection, or if you have surgery or a medical emergency. Tell your doctor about any such situation that affects you during treatment.

Steroid medications such as triamcinolone can weaken your immune system, making it easier for you to get an infection or worsening an infection you already have or have recently had. Tell your doctor about any illness or infection you have had within the past several weeks.

Avoid being near people who are sick or have infections. Call your doctor for preventive treatment if you are exposed to chicken pox or measles. These conditions can be serious or even fatal in people who are using steroid medication.

Do not receive a "live" vaccine while you are taking triamcinolone. Vaccines may not work as well while you are taking a steroid.

Do not stop using triamcinolone suddenly, or you could have unpleasant withdrawal symptoms. Talk to your doctor about how to avoid withdrawal symptoms when stopping the medication. Wear a medical alert tag or carry an ID card stating that you take triamcinolone. Any medical care provider who treats you should know that you take steroid medication.

## Before taking this medicine

You should not use this medication if you are allergic to triamcinolone, or if you have a fungal infection anywhere in your body.

Steroid medication can weaken your immune system, making it easier for you to get an infection. Steroids can also worsen an infection you already have, or reactivate an infection you recently had. Before taking this medication, tell your doctor about any illness or infection you have had within the past several weeks.

To make sure you can safely take triamcinolone, tell your doctor if you have any of these other conditions:

- liver disease (such as cirrhosis);
- kidney disease;
- a thyroid disorder;
- diabetes;
- a history of malaria;
- tuberculosis;
- osteoporosis;
- a muscle disorder such as myasthenia gravis;
- glaucoma or cataracts;
- herpes infection of the eyes;
- stomach ulcers, ulcerative colitis, or diverticulitis;
- depression or mental illness;
- congestive heart failure; or
- high blood pressure

FDA pregnancy category C. It is not known whether triamcinolone will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medication. Triamcinolone can pass into breast milk and may harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby. Triamcinolone can affect growth in children. Talk with your doctor if you think your child is not growing at a normal rate while using triamcinolone.

 [Triamcinolone pregnancy and breastfeeding warnings](#) (more detail)

## How should I take triamcinolone?

Take triamcinolone exactly as prescribed by your doctor. Do not take in larger or smaller amounts or for longer than recommended. Follow the directions on your prescription label.

Your doctor may occasionally change your dose to make sure you get the best results from this medication.

Take triamcinolone with food to prevent stomach upset.

Your steroid medication needs may change if you have unusual stress such as a serious illness, fever or infection, or if you have surgery or a medical emergency. Tell your doctor about any such situation that affects you.

This medication can cause you to have unusual results with certain medical tests. Tell any doctor who treats you that you are using triamcinolone.

Do not stop using triamcinolone suddenly, or you could have unpleasant withdrawal symptoms. Talk to your doctor about how to avoid withdrawal symptoms when stopping the medication. Carry an ID card or wear a medical alert bracelet stating that you are taking a steroid, in case of emergency. Any doctor, dentist, or emergency medical care provider who treats you should know that you are taking steroid medication.

Store triamcinolone at room temperature away from moisture and heat.

 [Detailed Triamcinolone dosage information](#)

## What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

An overdose of triamcinolone is not expected to produce life threatening symptoms. However, long term use of high steroid doses can lead to symptoms such as thinning skin, easy bruising, changes in the shape or location of body fat (especially in your face, neck, back, and waist), increased acne or facial hair, menstrual problems, impotence, or loss of interest in sex.

## What should I avoid?

Avoid being near people who are sick or have infections. Call your doctor for preventive treatment if you are exposed to chicken pox or measles. These conditions can be serious or even fatal in people who are using triamcinolone.

Do not receive a "live" vaccine while using triamcinolone. The vaccine may not work as well during this time, and may not fully protect you from disease. Avoid drinking alcohol while you are taking triamcinolone.

## Triamcinolone side effects

Get emergency medical help if you have any of these signs of an allergic reaction to triamcinolone: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have a serious side effect such as:

- problems with your vision;

- swelling, rapid weight gain, feeling short of breath;
- severe depression, unusual thoughts or behavior, seizure (convulsions);
- bloody or tarry stools, coughing up blood;
- pancreatitis (severe pain in your upper stomach spreading to your back, nausea and vomiting, fast heart rate);
- low potassium (confusion, uneven heart rate, extreme thirst, increased urination, leg discomfort, muscle weakness or limp feeling); or
- dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, uneven heartbeats, seizure).

Less serious triamcinolone side effects may include:

- sleep problems (insomnia), mood changes;
- acne, dry skin, thinning skin, bruising or discoloration;
- slow wound healing;
- increased sweating;
- headache, dizziness, spinning sensation;
- nausea, stomach pain, bloating;
- muscle weakness; or
- changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist);

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Triamcinolone side effects](#) (more detail)

## Related/similar drugs

### Nplate

Nplate is used to prevent bleeding episodes in people with chronic immune thrombocytopenic purpura ...

#### Reviews & ratings

7.5 / 10

[11 Reviews](#)

[View more](#)

### Otezla

Otezla (apremilast) is used to treat plaque psoriasis, psoriatic arthritis, and oral ulcers ...

## Reviews & ratings

5.2 / 10

[439 Reviews](#)

[View more](#)

### FEATURED

## Benlysta

Benlysta infusion is used to treat active systemic lupus erythematosus (SLE) and active lupus ...

## Reviews & ratings

7.9 / 10

[63 Reviews](#)

[View more](#)

## Saphnelo

Saphnelo is used for moderate to severe SLE (lupus) for adults who are also receiving other lupus ...

## Reviews & ratings

8.4 / 10

[10 Reviews](#)

[View more](#)

## Skyrizi

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis, ulcerative colitis ...

## Reviews & ratings

7.9 / 10

[56 Reviews](#)

[View more](#)

## Xeljanz

Xeljanz (tofacitinib) is an oral Janus kinase (JAK) inhibitor that may be used alone or with other ...

## Reviews & ratings

5.7 / 10

[145 Reviews](#)

[View more](#)

## Stelara

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

### Reviews & ratings

7.6 / 10

[252 Reviews](#)

[View more](#)

## Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

### Reviews & ratings

7.0 / 10

[223 Reviews](#)

[View more](#)

## Xolair

Xolair injection (omalizumab) is used to reduce the risk of severe food allergy reactions ...

### Reviews & ratings

7.5 / 10

[229 Reviews](#)

[View more](#)

## Duloxetine

Duloxetine is a selective serotonin and norepinephrine reuptake inhibitor antidepressant used to ...

### Reviews & ratings

6.2 / 10

[2,456 Reviews](#)

[View more](#)

## What other drugs will affect triamcinolone?

Many drugs can interact with triamcinolone. Below is just a partial list. Tell your doctor if you are using:

- aspirin (taken on a daily basis or at high doses);
- a diuretic (water pill);
- a blood thinner such as warfarin (Coumadin, Jantoven);
- cyclosporine (Gengraf, Neoral, Sandimmune);
- insulin or diabetes medications you take by mouth;
- ketoconazole (Nizoral);
- rifampin (Rifadin, Rifater, Rifamate, Rimactane); or
- seizure medications such as phenytoin (Dilantin) or phenobarbital (Luminal, Solfoton).

This list is not complete and other drugs may interact with triamcinolone. Tell your doctor about all medications you use. This includes prescription, over-the-counter, vitamin, and herbal products. Do not start a new medication without telling your doctor.

 [Triamcinolone drug interactions](#) (more detail)

## Does triamcinolone interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

triamcinolone

+

Enter a drug name

Add

## Popular FAQ

- Where should you not use triamcinolone acetonide cream?

▼
- Is triamcinolone acetonide an antifungal cream?

▼
- Is triamcinolone good for poison ivy?

▼
- Clobetasol vs. triamcinolone - how do they compare?

▼
- What are steroid injections (cortisone shots)?

▼
- Halobetasol vs triamcinolone: which is better?

▼

## More FAQ

- [What causes Plaque Psoriasis?](#)
- [Can triamcinolone acetonide B be used to treat severe diaper rash on a little girl?](#)

[View more FAQ...](#)

## More about triamcinolone

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(226\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Support group](#)
- [Drug class: glucocorticoids](#)
- [Breastfeeding](#)

## Patient resources

- [Triamcinolone \(Intra-articular\) advanced reading](#)

## Other brands

[Kenalog-40](#), [Zilretta](#), [Aristocort](#), [Azmacort](#), ... +5 more

## Professional resources

- [Triamcinolone Acetonide monograph](#)
- [Triamcinolone Injection \(FDA\)](#)

## Other brands

[Kenalog-40](#), [Zilretta](#), [Kenalog-10](#)

## Related treatment guides

- [Allergic Reactions](#)
- [Alopecia](#)
- [Osteoarthritis](#)



- [Adrenocortical Insufficiency](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 7.04.

### DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	Drug history at FDA
WADA Class	
	Anti-Doping Classification

### User Reviews & Ratings

6.7 / 10

[226 Reviews](#)

### Related News

[Do Steroid Shots Help Ease Back Pain? Experts Weigh In](#)

### Images

[Triamcinolone 4 MG \(5388\)](#)



# Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



## About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

## Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

## Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.