

Home

Ezetimibe

Ezetimibe 🕁

Pronunciation: ez-ET-i-mibe Generic name: ezetimibe Brand name: Zetia

Dosage form: oral tablet (10 mg)

Drug class: Cholesterol absorption inhibitors

Medically reviewed by Melisa Puckey, BPharm. Last updated on May 6, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is ezetimibe?

Ezetimibe is a medicine used to lower cholesterol in people who have high blood cholesterol. Ezetimibe works by reducing the amount of cholesterol your body absorbs from your intestines. High cholesterol can cause fatty deposits to form in your blood vessels which increases your risk of heart disease.

Ezetimibe is used in patients who cannot control their cholesterol levels by diet and exercise alone.

You can take ezetimibe as a single ingredient tablet or it is also available in a tablet that contains both ezetimibe and another cholesterol lowering medication called simvastatin.

Warnings

Ezetimibe is only part of a complete program of treatment that also includes diet, exercise, and weight control. Follow your diet, medication, and exercise routines very closely.

Some cholesterol medications should not be taken at the same time. If you take ezetimibe with another cholesterol medicine, follow your doctor's dosing instructions very carefully.

You should not use ezetimibe if you have moderate to severe liver disease. You should not use this medicine with a "statin" cholesterol medicine if you have active liver disease, or if you are pregnant or breast-feeding a baby.

Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine.

Before taking this medicine

You should not use ezetimibe if you are allergic to it, or if you have:

• moderate to severe liver disease.

Ezetimibe is not approved for use by anyone younger than 10 years old.

You should not use ezetimibe with a "statin" cholesterol medicine (Zocor, Lipitor, Crestor, and others) if:

- · you have active liver disease;
- · you are pregnant; or
- you are breastfeeding a baby.

Statin cholesterol medications can cause the breakdown of muscle tissue, which can lead to kidney failure. This happens more often in women, in older adults, or people who have kidney disease or poorly controlled hypothyroidism (underactive thyroid).

To make sure this medicine is safe for you, tell your doctor if you have:

- · liver disease;
- · kidney disease; or
- · unexplained muscle pain or weakness.

Tell your doctor if you are pregnant or breastfeeding.

Use effective birth control to prevent pregnancy while you are using ezetimibe with a statin medicine. Tell your doctor if you plan to become pregnant.

Do not breastfeed if you take ezetimibe with a statin medicine.

Ezetimibe pregnancy and breastfeeding warnings (more detail)

How should I take ezetimibe?

Take ezetimibe exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

Ezetimibe is usually taken once daily at the same time each day.

You may take this medicine with or without food.

Ezetimibe may be taken at the same time with fenofibrate, or with a statin medication such as atorvastatin, lovastatin, simvastatin, pravastatin, or fluvastatin.

Your treatment may also include diet, exercise, weight control, and blood tests.

You may need frequent medical tests, even if you have no symptoms.

Your cholesterol levels may not improve for up to 2 weeks.

Store at room temperature away from moisture and heat.

1 Ezetimibe patient tips (more detail)

Dosing information

Usual Adult Dose for Hyperlipidemia:

10 mg orally once a day.

Usual Adult Dose for Homozygous Familial Hypercholesterolemia:

10 mg orally once a day

Usual Adult Dose for Sitosterolemia:

10 mg orally once a day.

Usual Pediatric Dose for Hyperlipidemia:

10 years or older: 10 mg orally once a day.

Usual Pediatric Dose for Homozygous Familial Hypercholesterolemia:

10 years or older: 10 mg orally once a day.

Usual Pediatric Dose for Sitosterolemia:

10 years or older: 10 mg orally once a day.

Detailed Ezetimibe dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking ezetimibe?

Avoid eating foods high in fat or cholesterol, or this medicine will not be as effective.

Ezetimibe side effects

Get emergency medical help if you have **signs of an allergic reaction to ezetimibe:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Ezetimibe can cause the breakdown of muscle tissue, which can lead to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark urine.

Side effects may be more likely in older adults.

Common ezetimibe side effects may include:

- diarrhea;
- cold symptoms such as stuffy nose, sneezing, sore throat;
- pain in an arm or leg;
- · joint pain; or
- muscle pain while taking ezetimibe with a statin.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Ezetimibe side effects (more detail)

Related/similar drugs

Lipitor

Lipitor is used to treat high cholesterol. Learn about side effects, interactions and indications.

Reviews & ratings

4.3 / 10

102 Reviews

View more

Crestor

Crestor (rosuvastatin) is used to treat high cholesterol and high triglycerides in the blood ...

Reviews & ratings

5.6 / 10

138 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Zetia

Zetia (ezetimibe) is used to treat high cholesterol. Includes Zetia side effects, interactions and ...

Reviews & ratings

4.3 / 10

75 Reviews

View more

Zocor

Zocor (simvastatin) reduces lowdensity lipoprotein cholesterol and total cholesterol in the blood ...

Reviews & ratings

5.0 / 10

22 Reviews

View more

Niacin

Niacin is used to lower cholesterol and triglycerides. Learn about side effects, interactions and ...

Reviews & ratings

7.4 / 10

117 Reviews

View more

Pravastatin

Pravastatin (Pravachol) is a statin medicine used to lower cholesterol and triglycerides in the ...

Reviews & ratings

4.5 / 10

78 Reviews

View more

Simvastatin

Simvastatin (Zocor) is a cholesterollowering medication that blocks the production of cholesterol ...

Reviews & ratings

4.8 / 10

131 Reviews

View more

What other drugs will affect ezetimibe?

If you also take other cholesterol medication: Take ezetimibe 2 hours before or 4 hours after taking cholestyramine, colestipol, or colesevelam.

Tell your doctor about all your other medicines, especially:

- cyclosporine;
- · fenofibric acid;
- gemfibrozil; or
- a blood thinner warfarin, Coumadin, Jantoven.

This list is not complete. Other drugs may interact with ezetimibe, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

f Ezetimibe drug interactions (more detail)

Does ezetimibe interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

ezetimibe
+
Enter a drug name
Add

Popular FAQ

Does Zetia (ezetimibe) cause memory loss?

Can Zetia (ezetimibe) cause liver damage?

How does Zetia (ezetimibe) work to lower cholesterol?

More FAQ

- Does Zetia (ezetimibe) cause weight gain/loss or muscle aches/pains?
- What happens if you stop taking Ezetimibe?
- When should you take ezetimibe?
- Is ezetimibe a statin drug?
- When is the best time to take ezetimibe?
- Does ezetimibe lower triglycerides?

View more FAQ...

More about ezetimibe

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (172)
- Drug images
- Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: cholesterol absorption inhibitors
- Breastfeeding
- En español

Patient resources

Other brands

Zetia

Professional resources

- Ezetimibe monograph
- Ezetimibe (FDA)

Other brands

Zetia

Related treatment guides

- High Cholesterol, Familial Heterozygous
- High Cholesterol
- Sitosterolemia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use ezetimibe only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 7.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

প্ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

3.9 / 10

172 Reviews

Related News

Two-Drug Combo of a Statin Plus Ezetimibe, Lowers Cholesterol Better Than Statins Alone

Images

Ezetimibe 10 mg (K 31)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.