

[Home](#)

2. Advil Cold and Sinus Liqui-Gels

Advil Cold and Sinus Liqui-Gel

Generic name: [ibuprofen and pseudoephedrine](#) [*EYE-bue-pro-fen-and-SOO-doe-ee-FED-rin*]**Brand names:** Advil Cold & Sinus, Advil Cold and Sinus Liqui-Gel**Drug class:** [Upper respiratory combinations](#)[Medically reviewed](#) by Drugs.com on Jun 4, 2024. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is Advil Cold and Sinus Liqui-Gel?

Ibuprofen is a nonsteroidal anti-inflammatory drug (NSAID). [Pseudoephedrine](#) is a decongestant.

[Advil Cold and Sinus](#) Liqui-Gel is a combination medicine used to treat stuffy nose, sinus congestion, cough, and pain or fever caused by the common cold or flu.

Advil Cold and Sinus Liqui-Gel may also be used for purposes not listed in this medication guide.

Advil Cold and Sinus Liqui-Gel side effects

Get emergency medical help if you have **signs of an allergic reaction** (sneezing, runny or stuffy nose, hives, wheezing or trouble breathing, swelling in your face or throat) **or a severe skin reaction** (fever, [sore throat](#), burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Get emergency medical help if you have **signs of a heart attack or stroke**: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath.

Advil Cold and Sinus Liqui-Gel may cause serious side effects. Stop using Advil Cold and Sinus Liqui-Gel and call your doctor at once if you have:

- confusion, severe drowsiness, ringing in your ears, severe dizziness, feeling like you might pass out;
- fast, pounding, or uneven heartbeat;
- easy bruising or bleeding (nosebleeds, bleeding gums);
- a skin rash, no matter how mild;
- **signs of stomach bleeding**--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- **liver problems**--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- **kidney problems**--little or no urination, swelling in your feet or ankles, feeling tired or short of breath; or

- **nerve problems**--fever, headache, neck stiffness, chills, increased sensitivity to light, seizure (convulsions).

Common side effects of Advil Cold and Sinus Liqui-Gel may include:

- diarrhea, [constipation](#);
- headache, dizziness, drowsiness, weakness;
- [upset stomach](#), nausea, vomiting, loss of appetite;
- feeling anxious or excited;
- sleep problems (insomnia); or
- flushing (warmth, redness, or tingly feeling).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Advil Cold and Sinus Liqui-Gels side effects](#) (more detail)

Related/similar drugs

Nasonex 24HR Allergy

Nasonex 24HR Allergy (mometasone)
is used to treat nasal allergies such as
congestion, sneezing ...

Reviews & ratings

[Add a review](#)

[View more](#)

FEATURED

Omvox

Omvox is used to treat moderate to
severe ulcerative colitis or Crohn's
disease in adults. This ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Warnings

Ibuprofen can increase your risk of fatal heart attack or stroke. Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG). **Ibuprofen may also cause stomach or intestinal bleeding, which can be fatal.**

Before taking this medicine

Ibuprofen can increase your risk of fatal heart attack or stroke, even if you don't have any risk factors. Do not use Advil Cold and Sinus Liqui-Gel just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Ibuprofen may also cause stomach or intestinal bleeding, which can be fatal. These conditions can occur without warning while you are using Advil Cold and Sinus Liqui-Gel, especially in older adults.

You should not use this medicine if you are allergic to ibuprofen or pseudoephedrine, or if you have ever had an asthma attack or severe allergic reaction after taking aspirin or an NSAID.

Do not use Advil Cold and Sinus Liqui-Gel if you have used an MAO inhibitor in the past 14 days. A dangerous [drug interaction](#) could occur. MAO inhibitors include [isocarboxazid](#), linezolid, [methylene blue](#) injection, phenelzine, [tranylcypromine](#), and others.

Do not give ibuprofen to a child younger than 12 years old.

Ask a doctor or pharmacist if this medicine is safe to use if you have ever had:

- [heart disease](#), [high blood pressure](#), [high cholesterol](#), diabetes, or if you smoke;
- a [heart attack](#), stroke, or blood clot;
- stomach ulcers or bleeding;
- asthma;
- liver or kidney disease;
- a thyroid disorder;
- diabetes; or
- [enlarged prostate](#), urination problems.

If you are pregnant, you should not take Advil Cold and Sinus Liqui-Gel unless your doctor tells you to. Taking an NSAID during the last 20 weeks of pregnancy can cause serious heart or kidney problems in the unborn baby and possible complications with your pregnancy.

Ask a doctor before using this medicine if you are breastfeeding.

How should I take Advil Cold and Sinus Liqui-Gel?

Use exactly as directed on the label, or as prescribed by your doctor. **An overdose of ibuprofen can damage your stomach or intestines.**

Take this medicine with food or milk to lessen [stomach upset](#).

Call your doctor if you have a fever lasting longer than 3 days, if you have new symptoms, or if your condition does not improve after taking this medication for 7 days.

If you need surgery, tell the surgeon ahead of time if you have taken this medicine within the past few days.

Store at room temperature away from moisture and heat.

 [Detailed Advil Cold and Sinus Liqui-Gels dosage information](#)

What happens if I miss a dose?

Since Advil Cold and Sinus Liqui-Gel is used when needed, you may not be on a dosing schedule. Skip any missed dose if it's almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include vomiting, stomach pain, diarrhea, ringing in your ears, severe drowsiness, agitation, sweating, coughing up blood, weak or shallow breathing, fainting, or seizure (convulsions).

What should I avoid while taking Advil Cold and Sinus Liqui-Gel?

Avoid drinking alcohol. It may increase your risk of stomach bleeding.

Ask a doctor or pharmacist before using other cough or cold medicines that may contain similar ingredients.

What other drugs will affect Advil Cold and Sinus Liqui-Gel?

Ask your doctor before using Advil Cold and Sinus Liqui-Gel if you take an antidepressant. Taking certain [antidepressants](#) with an NSAID may cause you to bruise or bleed easily.

Ask a doctor or pharmacist before using Advil Cold and Sinus Liqui-Gel with any other medications, especially:

- lithium;
- [methotrexate](#);
- a blood thinner (warfarin, Coumadin, Jantoven);
- heart or blood pressure medication, including a diuretic or "water pill"; or
- steroid medicine (such as prednisone).

This list is not complete. Other drugs may affect Advil Cold and Sinus Liqui-Gel, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

i [Advil Cold and Sinus Liqui-Gels drug interactions](#) (more detail)

Does Advil Cold and Sinus Liqui-Gels interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Advil Cold and Sinus Liqui-Gels

+

Enter a drug name

Add

More about Advil Cold and Sinus Liqui-Gels (ibuprofen / pseudoephedrine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(2\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: upper respiratory combinations](#)

Professional resources

Other brands

[Motrin Childrens Cold](#)

Other formulations

- [Advil](#)

Related treatment guides

- [Sinus Symptoms](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 9.02.

Advil Pediatric Dosage Guide

Use these dosage charts for infants and children under 12 years old:

[Advil Dosage Charts \(Infants & Children\)](#)

DRUG STATUS

Availability

OTC Over the counter

Pregnancy & Lactation

 Risk data available

CSA Schedule*

N/A Not a controlled drug

WADA Class

 Anti-Doping Classification



User Reviews & Ratings

[Review this drug](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.