

Home

Mirabegron

Mirabegron ⊲[∞]

Generic name: mirabegron [*MIR-a-BEG-ron*] **Brand names:** Myrbetriq, Myrbetriq Granules

Dosage forms: oral granule for reconstitution, extended release (8 mg/mL), oral tablet, extended release (25 mg; 50 mg)

Drug class: Urinary antispasmodics



Medically reviewed by Drugs.com on Mar 26, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is mirabegron?

Mirabegron is used **in adults** to treat overactive bladder with symptoms of frequent or urgent urination and urinary incontinence. Mirabegron is sometimes used together with another medicine called solifenacin (Vesicare).

Mirabegron is used **in children** to treat neurogenic detrusor overactivity (NDO). Urinary incontinence caused by NDO is related to permanent nerve damage from conditions such as multiple sclerosis or spinal injury.

Mirabegron **tablets** may be used in children at least 3 years old. Mirabegron **granules** are for use in children at least 3 years old who also weigh at least 77 pounds (35 kilograms).

Mirabegron may also be used for purposes not listed in this medication guide.

Mirabegron side effects

Stop using mirabegron and get emergency medical help if you have **signs of an allergic reaction:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Mirabegron may cause serious side effects. Call your doctor at once if you have:

- pain or burning when you urinate; or
- dangerously high blood pressure--severe headache, blurred vision, pounding in your neck or ears.

Common side effects of mirabegron may include:

- painful urination;
- · increased blood pressure;
- sinus pain, sore throat;
- · constipation;

- · headache; or
- dry mouth or fast heartbeats (when taken with solifenacin).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Mirabegron side effects (more detail)

Related/similar drugs

Botox

Botox is used to treat chronic migraines, excessive sweating, bladder conditions, eye muscle ...

Reviews & ratings

5.7 / 10

473 Reviews

View more

Myrbetriq

Myrbetriq (mirabegron) is used to treat overactive bladder with symptoms of frequent or urgent ...

Reviews & ratings

4.8 / 10

276 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Gemtesa

Gemtesa (vibegron) is used to treat overactive bladder symptoms (OAB) in

adults or men taking ...

Reviews & ratings

7.6 / 10

200 Reviews

View more

Estrace

Estrace (estradiol) is used to treat symptoms of menopause such as hot flashes and vaginal dryness ...

Reviews & ratings

7.3 / 10

12 Reviews

View more

Oxybutynin

Oxybutynin is used to treat symptoms of overactive bladder such as incontinence. Learn about side ...

Reviews & ratings

6.7 / 10

458 Reviews

View more

Detrol

Detrol is used to treat overactive bladder symptoms of urinary frequency, urgency, and ...

Reviews & ratings

7.5 / 10

17 Reviews

View more

Warnings

You should not use mirabegron if you have uncontrolled hypertension (high blood pressure).

Before taking this medicine

You should not use mirabegron if you are allergic to it.

Tell your doctor if you have ever had:

- high blood pressure;
- a bladder obstruction;
- trouble emptying your bladder (very little urine or a weak stream of urine);
- · kidney disease; or
- liver disease.

Tell your doctor if you are pregnant or breastfeeding.

Mirabegron should not be given to a child younger than 3 years old.

How should I take mirabegron?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

If you take mirabegron with solifenacin, take both medicines at the same time each day.

Take the mirabegron tablet with a full glass of water. Swallow the tablet whole and do not crush, chew, or break it.

An adult may take a mirabegron tablet with or without food. However, a child should take the tablet with food.

Both adults and children should take mirabegron granules with food.

A pharmacist will mix mirabegron granules into a suspension (liquid) before you receive the medicine.

Shake the **suspension** before you measure a dose. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

If the **suspension** will not be used for 2 days or longer, shake the bottle vigorously for 1 minute each day to keep the granules thoroughly mixed.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions.

Your dose needs may change if you switch from using mirabegron tablets to using the suspension. **Avoid medication** errors by using only the form and strength your doctor prescribes.

Your blood pressure will need to be checked often.

It may take up to 8 weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

Store at room temperature away from moisture and heat. Throw away any unused mirabegron **suspension** after the **expiration** date on the medicine label has passed.

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if you are more than 12 hours late for the dose. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

An overdose may cause pounding heartbeats or fluttering in your chest.

What should I avoid while taking mirabegron?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

What other drugs will affect mirabegron?

Tell your doctor about all your other medicines, especially:

- · digoxin;
- flecainide:
- propafenone;
- · solifenacin; or
- thioridazine.

This list is not complete. Other drugs may affect mirabegron, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Mirabegron drug interactions (more detail)

Does mirabegron interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

mirabegron			
+			
Enter a drug r	ame		Add

More about mirabegron

Check interactions



- · Compare alternatives
- Pricing & coupons
- Reviews (352)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: urinary antispasmodics
- Breastfeeding
- En español

Patient resources

· Mirabegron advanced reading

Other brands

Myrbetriq

Professional resources

- Mirabegron monograph
- Mirabegron (FDA)

Other brands

Myrbetriq

Related treatment guides

- Overactive Bladder
- Urinary Frequency
- Urinary Incontinence
- Neurogenic Detrusor Overactivity

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ত Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

User Reviews & Ratings

4.7 / 10

352 Reviews

Images

Mirabegron Extended-Release 50 mg (1160)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×