

[Home](#)2. [Promethazine DM](#)

Promethazine DM

Generic name: [dextromethorphan and promethazine](#) [*dex-troe-me-THOR-fan-and-pro-METH-a-zeen*]**Drug class:** [Upper respiratory combinations](#)[Medically reviewed](#) by Drugs.com on Aug 11, 2023. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is Promethazine DM?

[Dextromethorphan](#) is a cough suppressant. [Promethazine](#) is an antihistamine.

Promethazine DM is a combination medicine used to treat cough, stuffy nose, runny nose, and sneezing caused by colds or allergies.

Promethazine DM is for use in adults and children at least 2 years old.

Promethazine DM may also be used for purposes not listed in this medication guide.

Promethazine DM side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have:

- weak or shallow breathing, breathing that stops during sleep;
- unusual restlessness or excitement;
- uncontrolled muscle movements in your neck, tongue, or eyes;
- fast, slow, or uneven heart rate;
- hallucinations, ringing in your ears;
- a seizure;
- fever, chills, weakness;
- easy bruising, unusual bleeding;
- yellowing of your skin or eyes; or
- **severe nervous system reaction**--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven

heartbeats, tremors, feeling like you might pass out.

Common side effects of Promethazine DM may include:

- drowsiness, dizziness;
- dry mouth, nose, or throat;
- blurred vision;
- nausea, vomiting; or
- loss of coordination.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Promethazine DM side effects](#) (more detail)

Related/similar drugs

Guaifenesin

Guaifenesin is an expectorant. It helps loosen congestion in your chest and throat. Includes ...

Reviews & ratings

7.1 / 10

[305 Reviews](#)

[View more](#)

Zyrtec

Zyrtec (cetirizine) is used to treat allergy symptoms such as sneezing, itching, watery eyes, or ...

Reviews & ratings

5.8 / 10

[197 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Diphenhydramine

Diphenhydramine is an antihistamine used to treat sneezing, runny nose, itching, hives and other ...

Reviews & ratings

5.5 / 10

[541 Reviews](#)

[View more](#)

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

[162 Reviews](#)

[View more](#)

Promethazine

Promethazine is a phenothiazine and antihistamine used to treat allergies, motion sickness, nausea ...

Reviews & ratings

7.2 / 10

[388 Reviews](#)

[View more](#)

Methylprednisolone

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

Reviews & ratings

7.1 / 10

[484 Reviews](#)

[View more](#)

Loratadine

Loratadine is a non-drowsy antihistamine used to relieve allergy symptoms like sneezing, runny ...

Reviews & ratings

4.7 / 10

[139 Reviews](#)

[View more](#)

Cetirizine

Cetirizine treats cold or allergy symptoms such as sneezing, itching, watery eyes, or runny nose ...

Reviews & ratings

5.8 / 10

[294 Reviews](#)

[View more](#)

Warnings

Promethazine can slow the breathing, which has caused death in very young children. Do not use Promethazine DM in a child younger than 2 years old. Follow all directions when giving the medicine to a child 2 years or older.

Do not use Promethazine DM if you have used an MAO inhibitor in the past 14 days, such as [isocarboxazid](#), linezolid, [methylene blue](#) injection, phenelzine, rasagiline, selegiline, or [tranylcypromine](#).

Before taking this medicine

You should not use this medicine if you are allergic to dextromethorphan or promethazine, or if you have:

- asthma or other breathing problems; or
- an allergy to medicines similar to promethazine ([chlorpromazine](#), [fluphenazine](#), [prochlorperazine](#), thioridazine, and others).

Do not use this medicine if you have used an MAO inhibitor in the past 14 days. A dangerous [drug interaction](#) could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, tranylcypromine, and others.

Tell your doctor if you have ever had:

- chronic obstructive pulmonary disease (COPD), sleep apnea, or other breathing disorder;

- a seizure;
- eczema;
- bone marrow depression;
- glaucoma;
- a [stomach ulcer](#);
- blockage in your stomach or intestines;
- a bladder obstruction or other urination problems;
- an [enlarged prostate](#);
- [liver disease](#);
- heart problems; or
- if you are on bed rest or otherwise debilitated.

Tell your doctor if you are pregnant or breastfeeding.

Taking this medicine during the last 2 weeks of pregnancy may increase the risk of bleeding in the newborn baby.

Ask a doctor before giving this medicine to a child younger than 12 years old.

How should I take Promethazine DM?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Do not use Promethazine DM in a child younger than 2 years old, and carefully follow all dosing directions about use in a child 2 years or older. **Promethazine can slow the breathing, which has caused death in very young children.**

Measure **liquid medicine** with the supplied syringe or a dose-measuring device (not a kitchen spoon).

This medicine can affect the results of certain medical tests, and may cause false results on a pregnancy test. Tell the doctor or laboratory staff if you have taken Promethazine DM in the past few days.

Call your doctor if your symptoms do not improve, or if they get worse.

Store at room temperature away from moisture and heat. Keep the bottle tightly closed when not in use.

 [Detailed Promethazine DM dosage information](#)

What happens if I miss a dose?

Promethazine DM is used when needed. If you are on a dosing schedule, skip any missed dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include overactive reflexes, feeling restless or confused, slow breathing, dilated pupils, or fainting.

What should I avoid while taking Promethazine DM?

Avoid driving, riding a bike, or any other hazardous activity until you know how Promethazine DM will affect you. Your reactions could be impaired.

Ask a doctor or pharmacist before using other cough or cold medicines that may contain dextromethorphan.

Avoid drinking alcohol.

Avoid sunlight, and wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

What other drugs will affect Promethazine DM?

Using Promethazine DM with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your other medicines, especially:

- an antidepressant;
- a barbiturate such as [phenobarbital](#);
- bronchodilator asthma medication;
- cold or allergy medicine (Benadryl and others);
- medicine to treat [Parkinson's disease](#);
- medicine to treat [overactive bladder](#); or
- medicine to treat stomach problems, [motion sickness](#), or [irritable bowel syndrome](#).

This list is not complete. Other drugs may affect Promethazine DM, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

 [Promethazine DM drug interactions](#) (more detail)

Does Promethazine DM interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Promethazine DM

+

Enter a drug name

Add

More about Promethazine DM (dextromethorphan / promethazine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(43\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: upper respiratory combinations](#)

Professional resources

- [Promethazine DM prescribing information](#)

Related treatment guides

- [Cough](#)
- [Allergic Rhinitis](#)
- [Cold Symptoms](#)


Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

DRUG STATUS	
Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	Drug history at FDA

User Reviews & Ratings

5.7 / 10

[43 Reviews](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.