

Home

2. Chlor-Trimeton

Chlor-Trimeton s

Generic name: chlorpheniramine [KLOR-fen-IR-a-meen]

Brand names: Aller-Chlor, Allergy Relief (Chlorpheniramine), Ed-Chlor Ped Jr, Wal-finate, Allergy Relief, ... show all 14 brands

Drug class: Antihistamines



Medically reviewed by Drugs.com on Jul 26, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is Chlor-Trimeton?

Chlor-Trimeton is an antihistamine that reduces the effects of natural chemical histamine in the body. Histamine can produce symptoms of sneezing, itching, watery eyes, and runny nose.

Chlor-Trimeton is used to treat runny nose, sneezing, itching, and watery eyes caused by allergies, the common cold, or the flu.

Chlor-Trimeton may also be used for purposes not listed in this medication guide.

Chlor-Trimeton side effects

Get emergency medical help if you have any of these **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Chlor-Trimeton may cause serious side effects. Stop using Chlor-Trimeton and call your doctor at once if you have:

- · fast or uneven heart rate;
- · mood changes;
- tremor, seizure (convulsions);
- easy bruising or bleeding, unusual weakness;
- · feeling short of breath; or
- little or no urinating.

Common side effects of Chlor-Trimeton may include:

- dizziness, drowsiness;
- dry mouth, nose, or throat;

- constipation;
- · blurred vision; or
- feeling nervous or restless.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Chlor-Trimeton side effects (more detail)

Related/similar drugs

Diphenhydramine

Diphenhydramine is an antihistamine used to treat sneezing, runny nose, itching, hives and other ...

Reviews & ratings

5.5 / 10

541 Reviews

View more

Triamcinolone

Triamcinolone is used to treat allergies, skin conditions, ulcerative colitis, and arthritis. Learn ...

Reviews & ratings

6.7 / 10

226 Reviews

View more

FEATURED

Xolair

Xolair injection (omalizumab) is used to reduce the risk of severe food allergy reactions ...

Reviews & ratings

7.5 / 10

229 Reviews

View more

Promethazine

Prometnazine is a pnenotniazine and antihistamine used to treat allergies, motion sickness, nausea ...

Reviews & ratings

7.2 / 10

388 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Loratadine

Loratadine is a non-drowsy antihistamine used to relieve allergy symptoms like sneezing, runny ...

Reviews & ratings

4.7 / 10

139 Reviews

View more

Methylprednisolone

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

Reviews & ratings

7.1 / 10

484 Reviews

View more

Cetirizine

Cetirizine treats cold or allergy symptoms such as sneezing, itching, watery eyes, or runny nose ...

Reviews & ratings

5.8 / 10

294 Reviews

View more

Montelukast

Montelukast is a daily oral medication used to prevent asthma attacks, exercise-induced ...

Reviews & ratings

5.6 / 10

571 Reviews

View more

Warnings

Always ask a doctor before giving a cough or cold medicine to a child. Death can occur from the misuse of cough and cold medicines in very young children.

You should not use Chlor-Trimeton if you have narrow-angle glaucoma, a blockage in your stomach or intestines, an enlarged prostate, if you are unable to urinate, or if you are having an asthma attack.

Before taking this medicine

You should not use Chlor-Trimeton if you are allergic to it, or if you have:

- narrow-angle glaucoma;
- a blockage in your stomach or intestines;
- an enlarged prostate;
- if you are unable to urinate; or
- if you are having an asthma attack.

Ask a doctor or pharmacist if it is safe for you to take Chlor-Trimeton if you have:

- a blockage in your digestive tract (stomach or intestines), a colostomy or ileostomy;
- liver or kidney disease;
- asthma or COPD, cough with mucus, or cough caused by smoking, emphysema, or chronic bronchitis;
- high blood pressure, heart disease, coronary artery disease, or recent heart attack;
- epilepsy or other seizure disorder;
- · urination problems;

- pheochromocytoma (an adrenal gland tumor); or
- · overactive thyroid.

FDA pregnancy category B. Chlor-Trimeton is not expected to harm an unborn baby. **Do not use this medicine without** your doctor's advice if you are pregnant.

Chlorpheniramine may pass into breast milk and may harm a nursing baby. Antihistamines may also slow breast milk production. Do not use this medicine without your doctor's advice if you are breast-feeding a baby.

Artificially sweetened cold medicine may contain phenylalanine. If you have phenylketonuria (PKU), check the medication label to see if the product contains phenylalanine.

How should I take Chlor-Trimeton?

Use exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. Chlor-Trimeton is usually taken only for a short time until your symptoms clear up.

Do not take for longer than 7 days in a row. Talk with your doctor if your symptoms do not improve after 7 days of treatment, or if you have a fever with a headache or skin rash.

Do not crush, chew, or break an extended-release tablet. Swallow it whole. Breaking the pill may cause too much of the drug to be released at one time.

Measure liquid medicine with a special dose-measuring spoon or medicine cup, not with a regular table spoon. If you do not have a dose-measuring device, ask your pharmacist for one.

Always ask a doctor before giving a cough or cold medicine to a child. Death can occur from the misuse of cough and cold medicines in very young children.

If you need surgery or medical tests, tell the surgeon or doctor ahead of time if you have taken this medicine within the past few days.

Store at room temperature away from moisture and heat. Do not allow the liquid form of this medicine to freeze.

Detailed Chlor-Trimeton dosage information

What happens if I miss a dose?

Since this medicine is taken when needed, you may not be on a dosing schedule. If you are taking the medication regularly, take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. **Do not** take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe forms of some of the side effects listed in this medication guide.

What should I avoid while taking Chlor-Trimeton?

Chlor-Trimeton may cause blurred vision or impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert and able to see clearly.

Drinking alcohol can increase certain side effects of chlorpheniramine.

Ask a doctor or pharmacist before using any other cold, cough, allergy, or sleep medicine. Antihistamines are contained in many combination medicines. Taking certain products together can cause you to get too much of a certain drug. Check the label to see if a medicine contains an antihistamine.

Avoid becoming overheated or dehydrated during exercise and in hot weather. Chlor-Trimeton can decrease sweating and you may be more prone to heat stroke.

What other drugs will affect Chlor-Trimeton?

Taking Chlor-Trimeton with other drugs that make you sleepy or slow your breathing can worsen these effects. Ask your doctor before taking this medicine with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

Ask a doctor or pharmacist before using this medicine if you are also using any other drugs, including prescription and over-the-counter medicines, vitamins, and herbal products. **Some medicines can cause unwanted or dangerous effects when used together.** Not all possible interactions are listed in this medication guide.

1 Chlor-Trimeton drug interactions (more detail)

Does Chlor-Trimeton interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Chlor-Trimeton	
+	
Enter a drug name	Add

Frequently asked questions

Can you take antihistamines when pregnant?

More about Chlor-Trimeton (chlorpheniramine)

- · Check interactions
- Compare alternatives
- Reviews (18)
- Drug images
- Side effects
- · Dosage information

- During pregnancy
- Drug class: antihistamines
- Breastfeeding

Patient resources

- Chlor-Trimeton (Chlorpheniramine Syrup) patient information
- Chlor-Trimeton (Chlorpheniramine Tablets)

Other brands

Allergy Relief, Aller-Chlor, C.P.M., Chlo-Amine, ... +7 more

Professional resources

• Chlor-Trimeton 12 Hour prescribing information

Related treatment guides

- Allergic Reactions
- Allergic Rhinitis
- Cold Symptoms
- Urticaria

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.


Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 10.01.

DRUG STATUS

Availability



Pregnancy & Lactation

ഴെ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

The Drug history at FDA

User Reviews & Ratings

8.7 / 10

18 Reviews

Images

Chlor-Trimeton 4 mg (080 SCHERING logo)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





