

#### Home

Paclitaxel

# Paclitaxel do

Generic name: paclitaxel [ PAK-li-TAK-sel ]

Brand names: Onxol, Taxol

**Dosage form:** intravenous solution (6 mg/mL)

Drug class: Mitotic inhibitors



Medically reviewed by Drugs.com on Aug 14, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

## What is paclitaxel?

Paclitaxel is used to treat breast cancer, ovarian cancer, and lung cancer. It is also used to treat AIDS-related Kaposi's sarcoma.

Paclitaxel is sometimes given after other treatments have failed.

Paclitaxel may also be used for purposes not listed in this medication guide.

## Paclitaxel side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; feeling like you might pass out; swelling of your face, lips, tongue, or throat.

Paclitaxel may cause serious side effects. Call your doctor at once if you have:

- · severe stomach pain or diarrhea;
- cold symptoms such as stuffy nose, sneezing, sore throat;
- flushing (warmth, redness, or tingly feeling);
- numbness, tingling, or burning pain in your hands or feet;
- severe redness or irritation, swelling or a hard lump, or other skin changes where the injection was given (may occur
  7 to 10 days after an injection);
- pain or burning when you urinate;
- cough with mucus, chest pain, feeling short of breath;
- chest pain, shortness of breath, fast or slow heartbeats;
- a light-headed feeling, like you might pass out;

- severe headache, blurred vision, pounding in your neck or ears;
- a seizure;
- · easy bruising, unusual bleeding, purple or red spots under your skin;
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet; or
- low white blood cell counts--fever, mouth sores, skin sores, sore throat, cough, trouble breathing.

Your cancer treatments may be delayed or permanently discontinued if you have certain side effects.

#### Common side effects of paclitaxel may include:

- fever, chills, or other signs of infection;
- anemia;
- feeling weak, tired, or light-headed;
- · bleeding;
- trouble breathing or swallowing;
- hair loss, skin rash, hives;
- numbness, tingling, or burning;
- swelling in your face, hands, or feet;
- sores or white patches in or around your mouth;
- joint or muscle pain;
- · nausea, vomiting, diarrhea; or
- tenderness or irritation where the medicine was injected.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Paclitaxel side effects (more detail)

# Related/similar drugs

#### **Trodelvy**

Trodelvy is a targeted anticancer medication used to treat types of breast cancer and bladder ...

#### **Reviews & ratings**

4 Reviews

View more

#### **Upuivu**

Opdivo is used to treat many cancers, such as melanoma, non-small cell lung cancer, malignant ...

#### **Reviews & ratings**

5.8 / 10

93 Reviews

#### View more

#### FEATURED

#### **Verzenio**

Verzenio is used to treat certain HRpositive and HER2-negative breast cancers in adults. It is an ...

#### **Reviews & ratings**

8.0 / 10

30 Reviews

#### View more

## Loqtorzi

Loqtorzi (toripalimab-tpzi) is a PD-1 inhibitor medicine used for the treatment of advanced ...

#### **Reviews & ratings**

Add a review

#### View more

## **Opdivo Qvantig**

Opdivo Qvantig is used to treat adults with melanoma, non-small cell lung cancer, renal cell ...

#### **Reviews & ratings**

Add a review

#### View more

## **Rybrevant**

Rybrevant is used to treat adults with non-small cell lung cancer (NSCLC) that has spread to other ...

#### **Reviews & ratings**

4 Reviews

#### View more

## **Keytruda**

Keytruda is used to treat multiple types of cancer such as melanoma, nonsmall cell lung cancer ...

#### **Reviews & ratings**

4.9 / 10

252 Reviews

View more

#### **Pembrolizumab**

Pembrolizumab (Keytruda) is a monoclonal antibody cancer medicine used to treat

#### **Reviews & ratings**

5.2 / 10

279 Reviews

View more

### **Tamoxifen**

Tamoxifen blocks the actions of estrogen and is used to treat and prevent some types of breast ...

#### **Reviews & ratings**

5.6 / 10

96 Reviews

View more

# **Warnings**

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; feeling like you might pass out; swelling of your face, lips, tongue, or throat.

Paclitaxel can lower blood cells that help your body fight infections and help your blood to clot. You may get an infection or bleed more easily. Call your doctor if you have unusual bruising or bleeding, or signs of infection (fever, sore throat, mouth sores, cough).

# Before taking this medicine

You should not be treated with paclitaxel if you are allergic to it, or if you have:

- low white blood cell (WBC) counts; or
- an allergy to castor oil (contained in paclitaxel and other medicines such as cyclosporine or teniposide).

Tell your doctor if you have ever had:

- · heart problems; or
- · liver disease.

Paclitaxel may harm an unborn baby. Use effective birth control to prevent pregnancy, and tell your doctor if you become pregnant.

You should not breastfeed while you are using paclitaxel.

# How is paclitaxel given?

Paclitaxel is given as an infusion into a vein. A healthcare provider will give you this injection.

paclitaxel must be given slowly, and the infusion can take 3 to 24 hours to complete.

Paclitaxel is usually given once every 2 to 3 weeks. Follow your doctor's dosing instructions very carefully.

You may be given other medications to prevent an allergic reaction while you are receiving paclitaxel.

Your breathing, blood pressure, oxygen levels, kidney function, and other vital signs will be watched closely while you are receiving paclitaxel.

Tell your caregivers if you feel any burning, pain, or swelling around the IV needle when paclitaxel is injected.

Paclitaxel can lower blood cells that help your body fight infections and help your blood to clot. **Your blood will need to be tested often.** Your cancer treatments may be delayed based on the results of these tests.

Detailed Paclitaxel dosage information

# What happens if I miss a dose?

Call your doctor for instructions if you miss an appointment for your paclitaxel injection.

# What happens if I overdose?

Since paclitaxel is given by a healthcare professional in a medical setting, an overdose is unlikely to occur.

# What should I avoid while using paclitaxel?

Paclitaxel contains alcohol and may cause a drunken feeling when the medicine is injected into your vein. Avoid drinking alcohol on the day of your paclitaxel injection.

Avoid being near people who are sick or have infections. Tell your doctor at once if you develop signs of infection.

Avoid activities that may increase your risk of bleeding or injury. Use extra care to prevent bleeding while shaving or brushing your teeth.

## What other drugs will affect paclitaxel?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Many drugs can affect paclitaxel. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here. Tell your doctor about all your current medicines and any medicine you start or stop using.

1 Paclitaxel drug interactions (more detail)

## Does paclitaxel interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

paclitaxel
+
Enter a drug name
Add

# **Popular FAQ**

Why give Taxol (Paxel) before carboplatin?

What is Paxel called in the USA?

# More about paclitaxel

- · Check interactions
- · Compare alternatives
- · Pricing & coupons
- Reviews (10)
- Latest FDA alerts (2)
- Side effects
- Dosage information
- During pregnancy
- · Drug class: mitotic inhibitors

- Breastfeeding
- En español

#### **Patient resources**

- · Paclitaxel advanced reading
- Paclitaxel (Conventional)

#### Other brands

Taxol, Onxol

## **Professional resources**

- · Paclitaxel monograph
- Paclitaxel (FDA)
- Paclitaxel Concentrate Injection (FDA)
- Paclitaxel Injection (FDA)

#### Other brands

**Taxol** 

## Related treatment guides

- Breast Cancer
- Breast Cancer, Adjuvant
- · Breast Cancer, Metastatic
- Kaposi's Sarcoma

## **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

## **DRUG STATUS**

**Availability** 

Rx Prescription only

## Pregnancy & Lactation ৪ Risk data available

CSA Schedule\*
N/A Not a controlled drug

Approval History

Drug history at FDA

### **User Reviews & Ratings**

7.2 / 10

10 Reviews

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com
Advertising policy
Attribution & citations

#### **Terms & privacy**

Terms of use

**Editorial policy** 

Privacy policy

#### **Support**

Help center

Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







