



[Home](#)

2. [Orap](#)

Orap

Generic name: [pimozide](#) [*PIM-oh-zide*]

Drug class: [Miscellaneous antipsychotic agents](#)



[Medically reviewed](#) by Drugs.com on Aug 9, 2023. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)



The Orap brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be [generic equivalents available](#).

What is Orap?

Orap works by changing the actions of chemicals in the brain.

Orap is used in people with [Tourette's syndrome](#). This medicine suppresses the physical (motor) and vocal (phonic) symptoms of tics when these symptoms interfere with daily life function.

Orap is not for use in treating motor tics that are not caused by Tourette's syndrome.

Orap may also be used for purposes not listed in this medication guide.

Orap side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

High doses or long-term use of pimozide can cause a serious movement disorder that may not be reversible.

The longer you use Orap, the more likely you are to develop this disorder, especially if you are an older adult.

Orap may cause serious side effects. Call your doctor at once if you have:

- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement);
- fast or pounding heartbeats, fluttering in your chest, [shortness of breath](#), and sudden dizziness (like you might pass out);
- fever, chills, tiredness, mouth sores, skin sores;
- a seizure (convulsions); or

- **severe nervous system reaction**--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out.

Common side effects of Orap may include:

- problems with speech or vision;
- [drowsiness](#), trouble sleeping;
- feeling restless;
- tight muscles;
- [constipation](#); or
- [dry mouth](#).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Orap side effects](#) (more detail)

Related/similar drugs

Abilify

Abilify is an antipsychotic medicine used to treat the symptoms of schizophrenia and bipolar ...

Reviews & ratings

6.1 / 10

[1,187 Reviews](#)

[View more](#)

FEATURED

Omvox

Omvox is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Abilify Maintena

Abilify Maintena (aripiprazole) is an atypical antipsychotic used to treat schizophrenia and ...

Reviews & ratings

[View more](#)

Aristada

Aristada (aripiprazole) is an antipsychotic medication. It works by changing the actions of ...

Reviews & ratings

4.5 / 10

[17 Reviews](#)

[View more](#)

Abilify Asimtufii

Abilify Asimtufii is used for the treatment of schizophrenia and bipolar I disorder, Abilify ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Warnings

You should not use Orap if you have [long QT syndrome](#), low blood levels of potassium or magnesium, or if you take any medication that can cause tics (such as a stimulant or ADHD medication).

Tell your doctor about all your current medicines and any you start or stop using. **Many drugs can cause serious medical problems if you take them together with Orap.**

Before taking this medicine

You should not use this medicine if you are allergic to Orap or other antipsychotic medicines, or if:

- you have low levels of potassium or magnesium in your blood;
- you have long QT syndrome (in you or a family member); or
- you take medicines that can cause tics, such as a stimulant or ADHD medication (Adderall, Ritalin, and others).

Some medicines can cause unwanted or dangerous effects when used with Orap. Your doctor may change your treatment plan if you also use:

- ADHD medication;

- an [antibiotic](#) or antifungal medication;
- anti-psychotic medication;
- an antidepressant;
- antiviral medicines to treat [hepatitis C](#) or HIV;
- anti-malaria medication;
- heart rhythm medication;
- medicine to prevent or treat nausea and vomiting (such as Zofran);
- "triptan" [migraine headache](#) medicine (such as Imitrex or Maxalt); or
- opioid medication.

Tell your doctor if you have ever had:

- [liver disease](#);
- kidney disease;
- seizures or [epilepsy](#);
- heart problems, [high blood pressure](#), or a [heart attack](#);
- an [enlarged prostate](#) or urination problems; or
- [glaucoma](#).

Taking Orap during the last 3 months of pregnancy may cause problems in the newborn, such as withdrawal symptoms, breathing problems, feeding problems, fussiness, tremors, and limp or stiff muscles. However, you may have withdrawal symptoms or other problems if you stop taking your medicine during pregnancy. **Do not stop taking this medicine without your doctor's advice.**

You should not breast-feed while using Orap.

Orap is not approved for use by anyone younger than 12 years old.

How should I take Orap?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Call your doctor if you are sick with vomiting or diarrhea. You may develop an electrolyte imbalance, which could cause heart rhythm problems while you are taking Orap.

It may take several weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve, or if they get worse.

Your doctor will need to check your progress on a regular basis. Your heart function may need to be checked using an electrocardiograph or ECG (sometimes called an EKG).

Store at room temperature away from moisture, heat, and light.

Do not stop using Orap suddenly, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using Orap.

i [Detailed Orap dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include muscle stiffness, shallow breathing, or feeling light-headed.

What should I avoid while taking Orap?

Avoid driving or hazardous activity until you know how Orap will affect you. Your reactions could be impaired.

Drinking alcohol can increase certain side effects of [pimozide](#).

Grapefruit may interact with Orap and lead to unwanted side effects. **Avoid the use of grapefruit products.**

What other drugs will affect Orap?

Orap can cause a serious heart problem. Your risk may be higher if you also use certain other medicines for infections, asthma, heart problems, high blood pressure, [depression](#), mental illness, cancer, malaria, or HIV.

Using Orap with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Many drugs can affect Orap, and some drugs should not be used at the same time. Tell your doctor about all your current medicines and any medicine you start or stop using. This includes prescription and over-the-counter medicines, vitamins, and [herbal products](#). **Not all possible interactions are listed here.**

i [Orap drug interactions](#) (more detail)

Does Orap interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Orap

+

Enter a drug name

Add

Frequently asked questions

- [What drugs cause tardive dyskinesia?](#)

More about Orap (pimozide)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(17\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: miscellaneous antipsychotic agents](#)
- [Breastfeeding](#)

Patient resources

- [Orap advanced reading](#)

Professional resources

- [Orap prescribing information](#)
- [Pimozide \(AHFS Monograph\)](#)

Related treatment guides

- [Tourette's Syndrome](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.


Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 13.01.

DRUG STATUS

Availability

 Discontinued

[Pregnancy & Lactation](#)

 Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+
years FDA approved 1984



User Reviews & Ratings

6.6 / 10

[17 Reviews](#)

Images

[Orap 2 mg \(ORAP 2 LEMMON\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)





[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.