

[Home](#)

2. [Microgestin Fe 1.5/30](#)
3. [Dosage](#)

## Microgestin Fe 1.5/30 Dosage

**Generic name:** [NORETHINDRONE ACETATE 1.5mg, ETHINYL ESTRADIOL 30ug](#);

**Dosage form:** tablets

**Drug class:** [Contraceptives](#)

[Medically reviewed](#) by Drugs.com. Last updated on Apr 16, 2024.

The compact tablet dispenser has been designed to make oral contraceptive dosing as easy and as convenient as possible. The tablets are arranged in four rows of seven tablets each, with the days of the week appearing on the compact tablet dispenser above the first row of tablets.

**Note:** Each compact tablet dispenser has been preprinted with the days of the week, starting with Sunday, to facilitate a Sunday-Start regimen. Six different day label stickers have been provided with the Detailed Patient & Brief Summary Patient Package Insert in order to accommodate a Day-1 Start regimen. If the patient is using the Day-1 Start regimen, she should place the self-adhesive day label sticker that corresponds to her starting day over the preprinted days.

**Important:** The patient should be instructed to use an additional method of protection until after the first week of administration in the initial cycle when utilizing the Sunday-Start regimen.

The possibility of ovulation and conception prior to initiation of use should be considered.

To achieve maximum contraceptive effectiveness, Microgestin Fe 1.5/30 should be taken exactly as directed and at intervals not exceeding 24 hours.

Microgestin Fe 1.5/30 provides a continuous administration regimen consisting of 21 yellow tablets of norethindrone acetate and ethinyl estradiol and 7 brown non-hormone containing tablets of ferrous fumarate. The ferrous fumarate tablets are present to facilitate ease of drug administration via a 28-day regimen and do not serve any therapeutic purpose. There is no need for the patient to count days between cycles because there are no “off-tablet days.”

**A. Sunday-Start Regimen:** The patient begins taking the first yellow tablet from the top row of the dispenser (labeled Sunday) on the first Sunday after menstrual flow begins. When the menstrual flow begins on Sunday, the first yellow tablet is taken on the same day. The patient takes one yellow tablet daily for 21 days. The last yellow tablet in the dispenser will be taken on a Saturday. Upon completion of all 21 yellow tablets, and without interruption, the patient takes one brown tablet daily for 7 days. Upon completion of this first course of tablets, the patient begins a second course of 28-day tablets, without interruption, the next day (Sunday), starting with the Sunday yellow tablet in the top row. Adhering to this regimen of one yellow tablet daily for 21 days, followed without interruption by one brown tablet daily for seven days, the patient will start all subsequent cycles on a Sunday.

**B. Day-1 Start Regimen:** The first day of menstrual flow is Day 1. The patient places the self-adhesive day label sticker that corresponds to her starting day over the preprinted days on the compact tablet dispenser. She starts taking one yellow tablet daily, beginning with the first yellow tablet in the top row. After the last yellow tablet (at the end of the third row) has been taken, the patient will then take the brown tablets for a week (7 days). For all subsequent cycles, the patient begins a new 28 tablet regimen on the eighth day after taking her last yellow tablet, again starting with the first

tablet in the top row after placing the appropriate day label sticker over the preprinted days on the compact tablet dispenser. Following this regimen of 21 yellow tablets and 7 brown tablets, the patient will start all subsequent cycles on the same day of the week as the first course.

Tablets should be taken regularly with a meal or at bedtime. It should be stressed that efficacy of medication depends on strict adherence to the dosage schedule.

## **Special Notes on Administration**

Menstruation usually begins two or three days, but may begin as late as the fourth or fifth day, after the brown tablets have been started. In any event, the next course of tablets should be started without interruption. If spotting occurs while the patient is taking yellow tablets, continue medication without interruption.

If the patient forgets to take one or more yellow tablets, the following is suggested:

One tablet is missed

- take tablet as soon as remembered
- take next tablet at the regular time

Two consecutive tablets are missed (week 1 or week 2)

- take two tablets as soon as remembered
- take two tablets the next day
- use another birth control method for seven days following the missed tablets

Two consecutive tablets are missed (week 3)

### **Sunday-Start Regimen:**

- take one tablet daily until Sunday
- discard remaining tablets
- start new pack of tablets immediately (Sunday)
- use another birth control method for seven days following the missed tablets

### **Day-1 Start Regimen:**

- discard remaining tablets
- start new pack of tablets that same day
- use another birth control method for seven days following the missed tablets

Three (or more) consecutive tablets are missed

### **Sunday-Start Regimen:**

- take one tablet daily until Sunday
- discard remaining tablets
- start new pack of tablets immediately (Sunday)

- use another birth control method for seven days following the missed tablets

#### **Day-1 Start Regimen:**

- discard remaining tablets
- start new pack of tablets that same day
- use another birth control method for seven days following the missed tablets

The possibility of ovulation occurring increases with each successive day that scheduled yellow tablets are missed. While there is little likelihood of ovulation occurring if only one yellow tablet is missed, the possibility of spotting or bleeding is increased. This is particularly likely to occur if two or more consecutive yellow tablets are missed.

If the patient forgets to take any of the seven brown tablets in week four, those brown tablets that were missed are discarded and one brown tablet is taken each day until the pack is empty. A back-up birth control method is not required during this time. A new pack of tablets should be started no later than the eighth day after the last yellow tablet was taken.

In the rare case of bleeding which resembles menstruation, the patient should be advised to discontinue medication and then begin taking tablets from a new compact tablet dispenser on the next Sunday or the first day (Day-1), depending on her regimen. Persistent bleeding which is not controlled by this method indicates the need for reexamination of the patient, at which time nonfunctional causes should be considered.

### **Use of Oral Contraceptives in the Event of a Missed Menstrual Period**

1. If the patient has not adhered to the prescribed dosage regimen, the possibility of pregnancy should be considered after the first missed period and oral contraceptives should be withheld until pregnancy has been ruled out.
2. If the patient has adhered to the prescribed regimen and misses two consecutive periods, pregnancy should be ruled out before continuing the contraceptive regimen.

After several months on treatment, bleeding may be reduced to a point of virtual absence. This reduced flow may occur as a result of medication, in which event it is not indicative of pregnancy.

### **More about Microgestin Fe 1.5 / 30 (ethinyl estradiol / norethindrone)**

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(131\)](#)
- [Drug images](#)
- [Side effects](#)
- [During pregnancy](#)
- [Drug class: contraceptives](#)
- [En español](#)

### **Patient resources**

- [Microgestin FE 1.5/30 birth control drug information](#)
- [Microgestin Fe 1.5/30](#)

## Other brands

[Lo Loestrin Fe](#), [Junel Fe 1/20](#), [Blisovi Fe 1/20](#), [Loestrin 21 1/20](#), ... +77 more

## Professional resources

- [Microgestin Fe 1.5/30 prescribing information](#)

## Other brands

[Lo Loestrin Fe](#), [Blisovi Fe 1/20](#), [Loestrin 21 1/20](#), [Hailey Fe 1/20](#), ... +81 more

## Related treatment guides

- [Birth Control](#)

## Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

### DRUG STATUS

#### Availability

**Rx** Prescription only

#### [Pregnancy & Lactation](#)


 Risk data available

#### CSA Schedule\*

**N/A** Not a controlled drug



#### Approval History

 Drug history at FDA



## User Reviews & Ratings

5.6 / 10

[131 Reviews](#)

## Related Drugs

[Mirena](#), [Provera](#), [Depo Provera](#), [Nexplanon](#), [Sprintec](#), [Yaz](#), [Slynd](#), [drospirenone](#), [levonorgestrel](#), [medroxyprogesterone](#)

## Related News

[Certain Birth Control Pills, Devices Linked to Higher Heart Risks](#)

## Images

Microgestin Fe 1.5/30 ferrous fumarate 75 mg (F N)



[View larger images](#)

## Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



### About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

### Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

### Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT

Copyright © 2000-2025 Drugs.com. All rights reserved.

