

#### Home

2. Lubiprostone

# **Lubiprostone** 49

**Generic name:** lubiprostone [ *loo-bee-PROS-tone* ]

Brand name: Amitiza

**Dosage form:** oral capsule (24 mcg; 8 mcg) **Drug class:** Chloride channel activators



Medically reviewed by Drugs.com on Oct 9, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

## What is lubiprostone?

Lubiprostone is used to treat chronic constipation, or constipation caused by opioid pain medicine.

Lubiprostone may also be used to treat irritable bowel syndrome in women with constipation as the main symptom.

Lubiprostone may also be used for purposes not listed in this medication guide.

## Lubiprostone side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Lubiprostone may cause serious side effects. Call your doctor at once if you have:

- severe or ongoing nausea or diarrhea;
- trouble breathing; or
- a light-headed feeling, like you might pass out.

You may have low blood pressure and feel faint shortly after taking lubiprostone, especially if you've been sick with vomiting or diarrhea. Stop taking lubiprostone and call your doctor if you feel light-headed after every dose or if you have severe diarrhea.

#### Common side effects of lubiprostone may include:

- nausea, stomach pain;
- · diarrhea, gas, bloating; or
- headache.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.



1 <u>Lubiprostone side effects</u> (more detail)

# Related/similar drugs

### **MiraLAX**

MiraLAX (polyethylene glycol) is used as a laxative to treat occasional constipation or irregular ...

### **Reviews & ratings**

7.2 / 10

262 Reviews

View more

#### Colace

Colace makes bowel movements softer and is used to treat or prevent constipation. Includes Colace ...

### **Reviews & ratings**

6.3 / 10

18 Reviews

### View more

#### FEATURED

### **Ozempic**

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

### **Reviews & ratings**

6.6 / 10

1,504 Reviews

### View more

### **Linzess**

Linzess (linaclotide) is used to treat chronic idiopathic constipation, irritable bowel syndrome ...

### **Reviews & ratings**

6.7 / 10

884 Reviews

----

#### View more

### **Bentyl**

Bentyl is used to treat functional bowel or irritable bowel syndrome. Bentyl may also be used for ...

### **Reviews & ratings**

7.3 / 10

138 Reviews

View more

### **Xifaxan**

Xifaxan (rifaximin) is used to treat travelers' diarrhea, irritable bowel syndrome with diarrhea ...

### Reviews & ratings

6.5 / 10

95 Reviews

View more

### **Dulcolax**

Dulcolax (bisacodyl) is a laxative used to treat constipation. Includes Dulcolax side effects ...

### **Reviews & ratings**

5.3 / 10

1,158 Reviews

View more

## **Warnings**

Stop taking lubiprostone and call your doctor if you feel light-headed every time you take lubiprostone or if you have severe diarrhea.

# Before taking this medicine

You should not use lubiprostone if you have a blockage in your digestive tract, or if you have severe diarrhea.

Tell your doctor if you have ever had liver disease.

Tell your doctor if you are pregnant or breastfeeding.

If you are breastfeeding, tell your doctor if you notice diarrhea in the nursing baby.

Not approved for use by anyone younger than 18 years old.

### How should I take Iubiprostone?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Take with food and water.

Swallow the capsule whole and do not crush, chew, break, or open it.

You may have tightness in your chest or feel short of breath within 1 hour after taking lubiprostone. This side effect should go away within 3 hours, but it may occur again when you take your next dose. Talk with your doctor if this side effect becomes bothersome.

Do not take lubiprostone if you have severe diarrhea. Call your doctor for instructions.

Your doctor will need to check your progress on a regular basis.

Store at room temperature away from heat and light.

## Lubiprostone dosing information

**Usual Adult Dose for Constipation -- Chronic:** 

24 mcg orally 2 times a day

Comments: The effectiveness of this drug in the treatment of opioid-induced constipation in patients taking diphenylheptane opioids (e.g., methadone) has not been established.

Use: Treatment of chronic idiopathic constipation and of opioid-induced constipation in adults with chronic non-cancer pain.

### Usual Adult Dose for Irritable Bowel Syndrome:

8 mcg orally 2 times a day

Use: Treatment of irritable bowel syndrome with constipation (IBS-C) in women age 18 years and older.

Detailed Lubiprostone dosage information

## What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

### What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

## What should I avoid while taking lubiprostone?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

### What other drugs will affect lubiprostone?

Make sure your doctor knows if you are using an opioid pain medicine.

Other drugs may affect lubiprostone, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all other medicines you use.

1 <u>Lubiprostone drug interactions</u> (more detail)

### Does lubiprostone interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Lubiprostone
+
Enter a drug name
Add

## **Popular FAQ**

Which drugs cause opioid-induced constipation?

## More about lubiprostone

- Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (280)
- Drug images
- Side effects
- Dosage information
- During pregnancy

- Drug class: chloride channel activators
- Breastfeeding
- En español

### **Patient resources**

Lubiprostone advanced reading

### Other brands

**Amitiza** 

### **Professional resources**

- Lubiprostone monograph
- Lubiprostone (FDA)

### Other brands

**Amitiza** 

### Related treatment guides

- Irritable Bowel Syndrome with Constipation
- Chronic Idiopathic Constipation
- Opioid-Induced Constipation
- Constipation
- Irritable Bowel Syndrome

### **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 4.01.


### **DRUG STATUS**

**Availability** 

Rx Prescription only

**Pregnancy & Lactation** 

൞ Risk data available



### **User Reviews & Ratings**

5.6 / 10

280 Reviews

### **Images**

Lubiprostone 24 mcg (A04)





View larger images

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





### **About**

About Drugs.com
Advertising policy
Attribution & citations

### Terms & privacy

Terms of use

**Editorial policy** 

Privacy policy

### **Support**

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×