

Home

Montelukast

Montelukast do

Generic name: montelukast [mon-te-LOO-kast]

Brand name: Singulair

Dosage forms: oral granule, oral tablet, oral chewable tablet

Drug class: Leukotriene modifiers

Medically reviewed by Carmen Pope, BPharm. Last updated on Oct 4, 2023.

Uses Warnings Before taking Dosage Side effects Interactions

What is montelukast?

Montelukast helps to reduce inflammation and may be used to prevent asthma attacks in adults and children at least 2 years old.

Montelukast is also used to prevent exercise-induced bronchoconstriction (a narrowing of the air passages in the lungs due to exercise that is also known as **exercise-induced asthma**) in adults and children who are at least 6 years old.

For adults and children at least 2 years old with symptoms of seasonal or year-round (perennial) allergies, montelukast may be considered when other treatments have not worked.

If you already take montelukast regularly to prevent asthma or allergy symptoms, do not take an extra dose to treat exercise-induced bronchoconstriction.

Montelukast is not a fast-acting rescue medicine for asthma attacks and needs to be taken daily to work properly.

Montelukast was FDA approved in 1998.

Warnings

Stop taking montelukast and call your doctor right away if you have any unusual changes in mood or behavior (such as agitation, confusion, depression, sleep problems, compulsive behaviors, hallucinations, or suicidal thoughts or actions).

Tell your doctor right away if you have signs of blood vessel inflammation: flu-like symptoms, severe sinus pain, a skin rash, numbness or a "pins and needles" feeling in your arms or legs.

Before taking this medicine

You should not use montelukast if you are allergic to it.

Tell your doctor if you have ever had:

- · mental illness or psychosis; or
- asthma, or a history of severe allergic reaction (sneezing, runny or stuffy nose, wheezing, shortness of breath) after taking aspirin or another NSAID.

The chewable montelukast tablet may contain phenylalanine and could be harmful if you have phenylketonuria (PKU).

Tell your doctor if you are pregnant or breastfeeding.

Do not give this medicine to a child without a doctor's advice.

Montelukast pregnancy and breastfeeding warnings (more detail)

How should I take montelukast?

Take montelukast exactly as prescribed by your doctor and read all medication guides or instruction sheets about montelukast.

Montelukast is not a fast-acting rescue medicine for asthma attacks. Seek medical attention if your breathing problems get worse quickly, or if you think your medications are not working.

Montelukast is usually taken once every evening, with or without food.

For exercise-induced bronchoconstriction, take a single dose at least 2 hours before exercise, and do not take another dose for at least 24 hours.

Swallow the regular tablet whole, with a glass of water.

You must chew the chewable tablet before you swallow it.

Place the oral granules directly in your mouth and swallow, or mix them with applesauce, mashed carrots, rice, or ice cream. The granules may also be mixed with baby formula or breast milk. Do not use any other type of liquid. Use the mixture within 15 minutes. Do not save for later use.

If you also use an oral steroid medication, you should not stop using it suddenly when you start taking montelukast. Follow your doctor's instructions about tapering your dose.

Do not change your dose or stop using asthma medication without your doctor's advice.

Store montelukast at room temperature away from moisture and heat. Do not open a packet of montelukast oral granules until you are ready to use them.

1 Detailed Montelukast dosage information

What happens if I miss a dose of montelukast?

If you miss a dose of montelukast, skip the missed dose and use your next dose at the regular time. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking montelukast?

Avoid situations or activities that may trigger an asthma attack.

If your asthma symptoms get worse when you take aspirin, avoid taking aspirin or other NSAIDs (nonsteroidal anti-inflammatory drugs) such as ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others.

Montelukast side effects

Get emergency medical help if you have **signs of an allergic reaction to montelukast:** hives, blisters, severe itching; difficult breathing; swelling of your face, lips, tongue, or throat.

Tell your doctor right away if you have signs of blood vessel inflammation: flu-like symptoms, severe sinus pain, a skin rash, numbness or a "pins and needles" feeling in your arms or legs.

Some people using montelukast have had new or worsening mental problems. Stop taking montelukast and call your doctor right away if you have unusual changes in mood or behavior, such as:

- · agitation, aggression, feeling restless or irritable;
- anxiety, depression, confusion, problems with memory or attention;
- stuttering, tremors, uncontrolled muscle movements;
- · suicidal thoughts or actions;
- hallucinations, sleep problems, vivid, dreams, sleep-walking; or
- compulsive or repetitive behaviors.

Common side effects of montelukast may include:

- stomach pain, diarrhea;
- fever or other flu symptoms;
- ear pain or full feeling, trouble hearing;
- · headache; or
- cold symptoms such as runny or stuffy nose, sinus pain, cough, sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Montelukast side effects (more detail)

Related/similar drugs

Breztri Aerosphere

Breztri

(budesonide/glycopyrrolate/formoterol fumarate) is a combination inhaler that may be used ...

Reviews & ratings

6.9 / 10

119 Reviews

View more

Xolair

Xolair injection (omalizumab) is used to reduce the risk of severe food allergy reactions ...

Reviews & ratings

7.5 / 10

229 Reviews

View more

FEATURED

Dupixent

Dupixent may be used to treat eczema, eosinophilic or oral-corticosteroid-dependent asthma, chronic ...

Reviews & ratings

7.2 / 10

499 Reviews

View more

Symbicort

Symbicort (budesonide and formoterol) is used to prevent bronchospasm in people with asthma or ...

Reviews & ratings

6.6 / 10

226 Reviews

View more

Zyrtec

Zyrtec (cetirizine) is used to treat allergy symptoms such as sneezing, itching, watery eyes, or ...

Reviews & ratings

5.8 / 10

197 Reviews

View more

Promethazine

Promethazine is a phenothiazine and antihistamine used to treat allergies, motion sickness, nausea ...

Reviews & ratings

7.2 / 10

388 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Loratadine

Loratadine is a non-drowsy antihistamine used to relieve allergy symptoms like sneezing, runny ...

Reviews & ratings

4.7 / 10

139 Reviews

View more

Cetirizine

Cetirizine treats cold or allergy symptoms such as sneezing, itching,

watery eyes, or runny nose ...

Reviews & ratings
5.8 / 10
294 Reviews

View more

What other drugs will affect montelukast?

Other drugs may affect montelukast, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all other medicines you use.

1 Montelukast drug interactions (more detail)

Does montelukast interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

montelukast
+
Enter a drug name
Add

More about montelukast

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (571)
- Drug images
- · Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: leukotriene modifiers
- Breastfeeding
- En español

Patient resources

- Montelukast Chewable Tablets patient information
- Montelukast Oral Granules

Other brands

Singulair

Professional resources

- Montelukast monograph
- Montelukast Chewable Tablets (FDA)
- Montelukast Oral Granules (FDA)
- Montelukast Tablets (FDA)

Other brands

Singulair

Related treatment guides

- Allergic Rhinitis
- COPD
- Allergies
- Bronchospasm Prophylaxis
- Asthma, Maintenance
- Asthma

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer


Copyright 1996-2025 Cerner Multum, Inc. Version: 18.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation



Approval HistoryDrug history at FDA

User Reviews & Ratings

5.6 / 10

571 Reviews

Images

Montelukast Sodium 10 mg (base) (I 114)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

V