

Home

Conjugated estrogens/medroxyprogesterone

Conjugated estrogens and medroxyprogesterone

Generic name: conjugated estrogens and medroxyprogesterone [KON-joo-GAY-ted-ES-troe-jenz-and-me-DOX-ee-proe-JES-ter-one]

Brand names: Premphase, Prempro

Dosage form: oral tablet (0.3 mg-1.5 mg; 0.45 mg-1.5 mg; 0.625 mg-2.5 mg; 0.625 mg-5 mg; biphasic)

Drug class: Sex hormone combinations



Medically reviewed by Drugs.com on Feb 6, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is conjugated estrogens and medroxyprogesterone?

Estrogen is a female sex hormone produced by the ovaries. Estrogen is necessary for many processes in the body. Conjugated estrogens are a mixture of estrogen hormones.

Medroxyprogesterone is a form of progesterone, a female hormone important for regulating ovulation and menstruation.

Conjugated estrogens and medroxyprogesterone is a combination medicine used to treat menopause symptoms such as hot flashes and vaginal changes, and to prevent osteoporosis (bone loss) in menopausal women.

Conjugated estrogens and medroxyprogesterone may also be used for purposes not listed in this medication guide.

Conjugated estrogens and medroxyprogesterone side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Conjugated estrogens and medroxyprogesterone may cause serious side effects. Call your doctor at once if you have:

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance;
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood, pain or warmth in one or both legs;
- swelling or tenderness in your stomach;
- jaundice (yellowing of the skin or eyes);
- memory problems, confusion, unusual behavior;

- · unusual vaginal bleeding, pelvic pain;
- a lump in your breast; or
- high levels of calcium in your blood--nausea, vomiting, constipation, increased thirst or urination, muscle weakness, bone pain, lack of energy.

Common side effects of conjugated estrogens and medroxyprogesterone may include:

- · nausea, gas, stomach pain;
- · headache, back pain;
- · depression, weakness;
- · breast pain;
- · itching, thinning scalp hair; or
- vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

① Conjugated estrogens / medroxyprogesterone side effects (more detail)

Related/similar drugs

Prolia

Prolia (denosumab) is an injection that is administered subcutaneously (under the skin) once every ...

Reviews & ratings

2.9 / 10

374 Reviews

View more

Fosamax

Fosamax (alendronate) is used to treat or prevent postmenopausal osteoporosis and steroid-induced ...

Reviews & ratings

2.8 / 10

76 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Premarin

Premarin is used to treat symptoms of menopause such as hot flashes, and vaginal dryness. Learn ...

Reviews & ratings

7.7 / 10

84 Reviews

View more

Reclast

Reclast (zoledronic acid) is used to treat osteoporosis in women after menopause and also to treat ...

Reviews & ratings

5.1 / 10

275 Reviews

View more

Estrace

Estrace (estradiol) is used to treat symptoms of menopause such as hot flashes and vaginal dryness ...

Reviews & ratings

7.3 / 10

12 Reviews

View more

Zometa

Zometa is used to treat high blood levels of calcium caused by cancer and multiple myeloma. Learn ...

Reviews & ratings

43 Reviews

View more

Denosumab

Denosumab (Prolia, Conexxence, Jubbonti, Ospomyv, and Stoboclo brands/biosimilars) is used to treat ...

Reviews & ratings

3.0 / 10

396 Reviews

View more

Warnings

You should not use conjugated estrogens and medroxyprogesterone if you have: undiagnosed vaginal bleeding, liver disease, a bleeding disorder, if you will have major surgery, or if you have ever had a heart attack, a stroke, a blood clot, or cancer of the breast, uterus/cervix, or vagina.

Do not use if you are pregnant.

Conjugated estrogens and medroxyprogesterone may increase your risk of developing a condition that may lead to uterine cancer. Report any unusual vaginal bleeding right away.

Using this medicine can increase your risk of blood clots, stroke, heart attack, or cancer of the breast, uterus, or ovaries. Estrogens should not be used to prevent heart disease, stroke, or dementia.

Before taking this medicine

You should not use this medicine if you have:

- unusual vaginal bleeding that has not been checked by a doctor;
- · liver disease;
- · a history of heart attack, stroke, or blood clot;
- an increased risk of having blood clots due to a heart problem or a hereditary blood disorder; or
- a history of hormone-related cancer, or cancer of the breast, uterus/cervix, or vagina.

Do not use conjugated estrogens and medroxyprogesterone if you are pregnant. Tell your doctor right away if you become pregnant during treatment.

Using this medicine can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have high blood pressure, diabetes, high cholesterol, if you are overweight, or if you smoke.

Conjugated estrogens and medroxyprogesterone should not be used to prevent heart disease, stroke, or dementia, because this medicine may actually increase your risk of developing these conditions.

Tell your doctor if you have ever had:

- a hysterectomy;
- · heart disease:
- liver problems, or jaundice caused by pregnancy or taking hormones;
- · kidney disease;
- gallbladder disease;
- asthma;
- · epilepsy or other seizure disorder;
- · migraines;
- · lupus;
- endometriosis or uterine fibroid tumors;
- hereditary angioedema;
- porphyria (a genetic enzyme disorder that causes symptoms affecting the skin or nervous system);
- a thyroid disorder; or
- · high levels of calcium in your blood.

Using this medicine may increase your risk of cancer of the breast, uterus, or ovaries. Talk with your doctor about this risk.

Estrogen lowers the hormone needed to produce breast milk and can slow breast milk production. Tell your doctor if you are breast-feeding.

How should I take conjugated estrogens and medroxyprogesterone?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Take the medicine at the same time each day.

Conjugated estrogens and medroxyprogesterone may increase your risk of developing a condition that may lead to uterine cancer. Your doctor may prescribe a progestin to take while you are using conjugated estrogens and medroxyprogesterone, to help lower this risk. **Report any unusual vaginal bleeding right away.**

The 28-day **Premphase** treatment consists of two differently colored pills, 14 of each color. Carefully follow the patient instructions about which pills to take on days 1 through 14 and which pills to take on days 15 through 28.

Your doctor should check your progress on a regular basis to determine whether you should continue this treatment. Self-examine your breasts for lumps on a monthly basis, and have regular mammograms.

If you need major surgery or will be on long-term bed rest, you may need to stop using this medicine for a short

time. Any doctor or surgeon who treats you should know that you are using conjugated estrogens and medroxyprogesterone.

Store at room temperature away from moisture, heat, and light.

Detailed Conjugated estrogens / medroxyprogesterone dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking conjugated estrogens and medroxyprogesterone?

Avoid smoking. It can greatly increase your risk of blood clots, stroke, or heart attack while using conjugated estrogens and medroxyprogesterone.

Grapefruit may interact with conjugated estrogens and medroxyprogesterone and lead to unwanted side effects. **Avoid** the use of grapefruit products.

What other drugs will affect conjugated estrogens and medroxyprogesterone?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Many drugs can affect conjugated estrogens and medroxyprogesterone. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here. Tell your doctor about all your current medicines and any medicine you start or stop using.

Conjugated estrogens / medroxyprogesterone drug interactions (more detail)

Does conjugated estrogens/medroxyprogesterone interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

conjugated estrogens/medroxyprogesterone
+
Enter a drug name
Add

More about conjugated estrogens / medroxyprogesterone

- · Check interactions
- Compare alternatives
- Reviews (70)
- Side effects
- Dosage information
- During pregnancy
- Drug class: sex hormone combinations
- En español

Patient resources

- · Conjugated estrogens and medroxyprogesterone advanced reading
- Estrogens (Conjugated/Equine) and Medroxyprogesterone

Other brands

Prempro, Premphase

Professional resources

Premphase Prempro prescribing information

Related treatment guides

- · Atrophic Urethritis
- Atrophic Vaginitis
- Oophorectomy
- Osteoporosis
- Postmenopausal Symptoms
- Primary Ovarian Failure

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 12.03.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ড Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

The Drug history at FDA

User Reviews & Ratings

7.0 / 10

70 Reviews

Images

Prempro (conjugated estrogens / medroxyprogesterone) 0.625 mg / 2.5 mg (PREMPRO)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×