

#### Home

2. Lysteda

# Lysteda 🖘

Generic name: tranexamic acid (Lysteda) [ tran-ex-AM-ik-AS-id ]

Drug class: Miscellaneous coagulation modifiers

Medically reviewed by Philip Thornton, DipPharm. Last updated on Sep 25, 2023.

Uses Warnings Before taking Dosage Side effects Interactions

## What is Lysteda?

Lysteda is a man-made form of an amino acid (protein) called lysine. Tranexamic acid prevents enzymes in the body from breaking down blood clots.

Lysteda is used to treat heavy menstrual bleeding. This medication will not treat premenstrual syndrome (PMS).

Lysteda may also be used for purposes not listed in this medication guide.

# **Warnings**

You should not use Lysteda if you are allergic to tranexamic acid, if you have problems with the blood vessels in your eyes, or if you have ever had a stroke, blood clot, or bleeding in your brain.

Before using Lysteda, tell your doctor if you have kidney disease, leukemia, a history of endometriosis, or if your menstrual cycles are less than 21 days apart or longer than 35 days apart.

Do not start taking Lysteda until your period has started. Do not take it for longer than 5 days in a row. Talk with your doctor if your symptoms do not improve after 2 cycles of treatment.

Do not take more than 6 tablets in one 24-hour period.

Use a barrier form of birth control (such as a condom or diaphragm with spermicide). Hormonal contraception (such as birth control pills, injections, implants, skin patches, and vaginal rings) may increase your risk of stroke, blood clot, or heart attack if they are used at the same time as Lysteda. Do not share Lysteda with another person, even if they have the same symptoms you have.

# Before taking this medicine

You should not take Lysteda if you are allergic to tranexamic acid, or if you have:

• problems with the blood vessels in your eyes;

- color blindness (only if you are receiving the injectable form of tranexamic acid);
- · a history of stroke;
- if you have a history of bleeding in your brain; or
- if you have recently had a blood clot.

To make sure you can safely take Lysteda, tell your doctor if you have any of these other conditions:

- leukemia;
- · kidney disease;
- · a history of endometriosis; or
- if your menstrual cycles are less than 21 days apart or longer than 35 days apart.

It is not known whether Lysteda will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using Lysteda. Tranexamic acid can pass into breast milk and may harm a nursing baby. Do not use Lysteda without telling your doctor if you are breast-feeding a baby. Do not give this medication to anyone under 18 years old without your doctor's advice. Do not share Lysteda with another person, even if they have the same symptoms you have.

1 Lysteda pregnancy and breastfeeding warnings (more detail)

## How should I use Lysteda?

Take Lysteda exactly as prescribed by your doctor. Do not take in larger or smaller amounts or for longer than recommended. Follow the directions on your prescription label.

Do not start taking Lysteda until your period has started. Do not take it for longer than 5 days in a row.

Lysteda is usually taken three times per day for up to 5 days during your menstrual period. Do not take more than 6 tablets in one 24-hour period.

Take Lysteda with a full glass of water.

You may take tranexamic acid with or without food.

Do not crush, chew, or break a Lysteda tablet. Swallow it whole.

Talk with your doctor if your symptoms do not improve after 2 cycles of treatment.

Store Lysteda at room temperature away from moisture and heat.

Detailed Lysteda dosage information

# What happens if I miss a dose?

Take the missed dose as soon as you remember. Then take your next dose at least 6 hours later. Do not take extra medicine to make up the missed dose.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe forms of some of the side effects listed in this medication guide.

## What should I avoid?

Avoid using a hormonal form of birth control such as birth control pills, injections, implants, skin patches, and vaginal rings. Hormonal contraception may increase your risk of stroke, blood clot, or heart attack while you are taking Lysteda. If you need to use birth control during treatment with Lysteda, use a barrier form (such as a condom or diaphragm with spermicide).

# Lysteda side effects

Get emergency medical help if you have any of these signs of an allergic reaction to Lysteda: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Stop taking Lysteda and call your doctor at once if you have a serious side effect such as:

- problems with your vision (including color vision);
- sudden numbness or weakness, especially on one side of the body;
- sudden headache, confusion, problems with vision, speech, or balance;
- sudden chest pain or trouble breathing;
- pain or swelling in one or both legs;
- migraine headache;
- pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating; or
- · feeling like you might pass out.

Less serious Lysteda side effects include:

- · mild headache;
- runny or stuffy nose;
- · stomach pain;
- · tired feeling;
- · eye redness; or
- back pain, joint or muscle pain.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**1** Lysteda side effects (more detail)

# Related/similar drugs

#### Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

## **Reviews & ratings**

6.1 / 10

2,624 Reviews

View more

## **Kyleena**

The Kyleena IUD is placed in your uterus by your healthcare provider to prevent pregnancy for up to ...

## **Reviews & ratings**

6.6 / 10

1,637 Reviews

View more

#### FEATURED

## **Ozempic**

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

## **Reviews & ratings**

6.6 / 10

1,504 Reviews

View more

## Liletta

Liletta (levonorgestrel) is an intrauterine device (IUD) used to prevent pregnancy and also to ...

## **Reviews & ratings**

5.5 / 10

647 Reviews

View more

## **Skyla**

Skyla (levonorgestrel intrauterine system) is flexible plastic T-shaped device used for prevention ...

#### **Reviews & ratings**

6.9 / 10

1,537 Reviews

View more

## Plan B One-Step

Plan B (levonorgestrel) is a morningafter pill (emergency contraceptive pill) that is used to ...

## **Reviews & ratings**

8.4 / 10

2.799 Reviews

View more

#### **Natazia**

Natazia (dienogest and estradiol) is an oral contraceptive tablet used for birth control. Includes ...

#### **Reviews & ratings**

6.5 / 10

67 Reviews

View more

## What other drugs will affect Lysteda?

Tell your doctor about all other medicines you use, especially:

- hormonal birth control (such as birth control pills, injections, implants, skin patches, and vaginal rings);
- tretinoin (Vesanoid);
- chemotherapy medication to treat leukemia;
- any type of medication to treat a bleeding episode or a blood clot; or
- factor IX (Bebulin VH, Konyne 80, Profilnine SD, Proplex T, and others).

This list is not complete and other drugs may interact with Lysteda. Tell your doctor about all medications you use. This includes prescription, over-the-counter, vitamin, and herbal products. Do not start a new medication without telling your

Lysteda drug interactions (more detail)

## Does Lysteda interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Lysteda	
+	
Enter a drug name	Add

# More about Lysteda (tranexamic acid)

- · Check interactions
- Compare alternatives
- Reviews (70)
- Drug images
- Latest FDA alerts (1)
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- FDA approval history
- Drug class: miscellaneous coagulation modifiers
- Breastfeeding

## **Patient resources**

## Other brands

Cyklokapron

## **Professional resources**

- Lysteda prescribing information
- Tranexamic Acid (AHFS Monograph)

#### Other brands

Cyklokapron

## Related treatment guides

- · Heavy Menstrual Bleeding
- Menstrual Disorders

## **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Lysteda only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

#### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

#### **DRUG STATUS**

#### **Availability**

Rx Prescription only

### **Pregnancy & Lactation**

৵ Risk data available

**CSA Schedule\*** 

N/A Not a controlled drug

#### **Approval History**

Drug history at FDA

## **User Reviews & Ratings**

7.9 / 10

70 Reviews

## **Images**

### Lysteda 650 mg (FP650)





View larger images

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



#### **About**

About Drugs.com
Advertising policy
Attribution & citations

## Terms & privacy

Terms of use Editorial policy Privacy policy

#### **Support**

Help center Sitemap Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×