

Home

2. BuSpar

Generic name: buspirone [byoo-SPYE-rone] **Brand names:** BuSpar, BuSpar Dividose

Drug class: Miscellaneous anxiolytics, sedatives and hypnotics

Medically reviewed by Kaci Durbin, MD. Last updated on Aug 13, 2023.

Uses Warnings Before taking Dosage Side effects Interactions

What is BuSpar?

BuSpar is an anti-anxiety medicine that affects chemicals in the brain that may be unbalanced in people with anxiety.

BuSpar is used to treat anxiety disorders or the symptoms of anxiety, such as fear, tension, irritability, dizziness, pounding heartbeat, and other physical symptoms.

BuSpar is not an anti-psychotic medication and should not be used in place of medication prescribed by your doctor for mental illness.

BuSpar may also be used for purposes not listed in this medication guide.

Warnings

Do not use BuSpar if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, and tranylcypromine.

Before taking this medicine

You should not use BuSpar if you are allergic to buspirone.

Do not use BuSpar if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, and tranylcypromine.

To make sure BuSpar is safe for you, tell your doctor if you have any of these conditions:

- · kidney disease; or
- · liver disease.

BuSpar is not expected to harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during treatment.

It is not known whether buspirone passes into breast milk or if it could harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

Buspirone is not FDA-approved for use by anyone younger than 18 years old, but has been used in children in adolescents under close medical supervision. Do not give buspirone to anyone younger than 18 years of age without consulting with a doctor.

BuSpar pregnancy and breastfeeding warnings (more detail)

How should I take BuSpar?

Take BuSpar exactly as it was prescribed for you. Follow all directions on your prescription label. Your doctor may occasionally change your dose to make sure you get the best results. Do not take this medicine in larger or smaller amounts or for longer than recommended.

You may take BuSpar with or without food but take it the same way each time.

Some BuSpar tablets are scored so you can break the tablet into 2 or 3 pieces in order to take a smaller amount of the medicine at each dose. Do not use a tablet if it has not been broken correctly and the piece is too big or too small. Follow your doctor's instructions about how much of the tablet to take.

If you have switched to BuSpar from another anxiety medication, you may need to slowly decrease your dose of the other medication rather than stopping suddenly. Some anxiety medications can cause withdrawal symptoms when you stop taking them suddenly after long-term use.

Buspirone can cause false positive results with certain medical tests. You may need to stop using the medicine for at least 48 hours before your test. Tell any doctor who treats you that you are using BuSpar.

Store at room temperature away from moisture, heat, and light.

Detailed BuSpar dosage information

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Although no deaths have been reported following BuSpar overdose, symptoms may include nausea, vomiting, dizziness, drowsiness or sleepiness, and stomach upset.

What should I avoid?

This medication may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be

alert.

Drinking alcohol may increase certain side effects of BuSpar.

Grapefruit and grapefruit juice may interact with buspirone and lead to unwanted side effects. Discuss the use of grapefruit products with your doctor.

BuSpar side effects

Get emergency medical help if you have any **signs of an allergic reaction to BuSpar**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- · chest pain;
- · shortness of breath; or
- a light-headed feeling, like you might pass out.

Common BuSpar side effects may include:

- · headache;
- · dizziness, drowsiness;
- sleep problems (insomnia);
- · nausea, upset stomach; or
- · feeling nervous or excited.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

BuSpar side effects (more detail)

Related/similar drugs

Prozac

Prozac (fluoxetine) is an SSRI antidepressant used to treat depression, OCD, panic disorder ...

Reviews & ratings

7.2 / 10

1,239 Reviews

View more

Cymbalta

Cymbalta (duloxetine) is used to treat

major depressive disorder, general anxiety disorder and ...

Reviews & ratings

6.3 / 10

1,899 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Paroxetine

Paroxetine is used treat depression, obsessive-compulsive disorder and anxiety disorders. Learn ...

Reviews & ratings

7.1 / 10

1,548 Reviews

View more

Diazepam

Diazepam is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

Reviews & ratings

8.4 / 10

884 Reviews

View more

Xanax

Xanax (alprazolam) is a benzodiazepine medication used to treat anxiety and panic disorders ...

Reviews & ratings

915 Reviews

View more

Venlafaxine

Venlafaxine is used to treat major depressive disorder, anxiety, and panic disorder. Learn about ...

Reviews & ratings

6.5 / 10

2,786 Reviews

View more

Zoloft

Zoloft is an antidepressant used to treat major depression, obsessivecompulsive disorder, panic ...

Reviews & ratings

7.3 / 10

1,972 Reviews

View more

Lorazepam

Lorazepam is in a group of drugs called benzodiazepines and is used to treat anxiety disorders ...

Reviews & ratings

7.9 / 10

1,054 Reviews

View more

Lexapro

Lexapro is used to treat anxiety and major depressive disorder. Learn about side effects ...

Reviews & ratings

7.6 / 10

2,376 Reviews

What other drugs will affect BuSpar?

Taking this medicine with other drugs that make you sleepy or slow your breathing can worsen these effects. The following medications can interact with BuSpar;

- MAO Inihibitors (including isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine as well as methylene blue injection);
- · Amitriptyline;
- Diazepam;
- · Haloperidol;
- Nefazodone;
- Trazodone:
- · Triazolam or flurazepam;
- · Diltiazem or verapamil;
- · Antibiotics, including erythromycin and rifampin;
- · Antifungals, including itraconazole; or
- · Cimetidine.

In addition, sleeping pills, narcotic pain medicine, muscle relaxers, and other medications for anxiety, depression, or seizures may interact with Buspar.

Other drugs not mentioned here may interact with buspirone, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

BuSpar drug interactions (more detail)

Does BuSpar interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

BuSpar	
+	
Enter a drug name	Add

More about BuSpar (buspirone)

· Check interactions

- · Compare alternatives
- Reviews (636)
- Drug images
- Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: miscellaneous anxiolytics, sedatives and hypnotics
- Breastfeeding

Patient resources

• Buspar Dividose advanced reading

Other brands

Vanspar

Professional resources

- BuSpar prescribing information
- BusPIRone (AHFS Monograph)

Related treatment guides

- Anxiety
- Panic Disorder

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use BuSpar only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

Availability

Rx Prescription only
Pregnancy & Lactation

& Risk data available

CSA Schedule*
N/A Not a controlled drug

Approval History
10+
years FDA approved 1986

User Reviews & Ratings

6.0 / 10

636 Reviews

Images

BuSpar 10 mg (BUSPAR MJ 10)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

 \times