



[Home](#)

2. Byetta

Byetta

Generic name: [exenatide](#) [*ex-EN-a-tide*]

Brand name: Byetta Prefilled Pen

Drug class: [GLP-1 Agonists \(Incretin Mimetics\)](#)

Medically reviewed by [Philip Thornton, DipPharm](#). Last updated on May 20, 2024.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

What is Byetta?

Byetta is an injectable diabetes medicine that helps control blood sugar levels. This medication helps your pancreas produce insulin more efficiently. Byetta is a short-acting form of exenatide.

Byetta is used together with diet and exercise to improve blood sugar control in adults with [type 2 diabetes](#). Byetta is not for treating type 1 diabetes.

This medication guide provides information about the Byetta, an immediate-release brand of exenatide. Bydureon is an extended-release brand of exenatide that is not covered in this medication guide.

Warnings

You should not use Byetta if you have severe kidney disease (or you are on dialysis), or if you have slowed digestion or [diabetic ketoacidosis](#).

You should not use this medicine if Byetta has ever caused you to have low platelet levels in your blood.

Stop using this medicine and call your doctor at once if you have [nausea and vomiting](#) with severe pain in your upper stomach spreading to your back.

You should not use Byetta together with fast-acting or short-acting insulins. Do not use this medicine together with Bydureon.

Before taking this medicine

You should not use Byetta if you are allergic to exenatide, or if you have:

- severe kidney disease (or if you are on dialysis);
- a severe stomach disorder that causes slow digestion;
- a history of low levels of platelets (blood cells that help your blood to clot) while using exenatide; or

- diabetic ketoacidosis (call your doctor for treatment with [insulin](#)).

To make sure Byetta is safe for you, tell your doctor if you have ever had:

- kidney disease or a history of kidney transplant;
- a stomach disorder that causes slow digestion;
- [pancreatitis](#); or
- gall stones.

Follow your doctor's instructions about using this medicine **if you are pregnant or you become pregnant**. Controlling diabetes is very important during pregnancy, and having high blood sugar may cause complications in both the mother and the baby.

Exenatide can make [birth control pills](#) less effective. If you take birth control pills, take your pill at least 1 hour before your Byetta injection.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk.

Byetta is not approved for use by anyone younger than 18 years old.

 [Byetta pregnancy and breastfeeding warnings](#) (more detail)

How should I use Byetta?

Use Byetta exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Byetta is injected under the skin. A healthcare provider may teach you how to properly use the medication by yourself.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you don't understand all instructions.

Prepare your injection only when you are ready to give it. You must give the injection right away after mixing.

Byetta is usually injected twice a day, before the morning and evening meals. Do not use after eating a meal. You must use Byetta within 60 minutes (1 hour) before eating. Your injections should be given at least 6 hours apart.

You may have [low blood sugar \(hypoglycemia\)](#) and feel very hungry, dizzy, irritable, confused, anxious, or shaky. To quickly treat [hypoglycemia](#), eat or drink a fast-acting source of sugar (fruit juice, hard candy, crackers, raisins, or non-diet soda).

Your doctor may prescribe a [glucagon](#) injection kit in case you have severe hypoglycemia. Be sure your family or close friends know how to give you this injection in an emergency.

Also watch for signs of high blood sugar (hyperglycemia) such as increased thirst or urination.

Blood sugar levels can be affected by [stress](#), illness, surgery, exercise, alcohol use, or skipping meals. **Ask your doctor before changing your dose or medication schedule.**

Use a needle and syringe only once and then place them in a puncture-proof "sharps" container. Follow state or local

laws about how to dispose of this container. Keep it out of the reach of children and pets.

Never share an injection pen or syringe with another person, even if the needle has been changed. Sharing these devices can allow infections or disease to pass from one person to another.

Exenatide can lower your platelet counts for up to 10 weeks after you stop using Byetta. You may need frequent medical tests while using this medicine and for a short time after you stop.

Byetta is only part of a complete treatment program that may also include diet, exercise, weight control, regular blood sugar testing, and special medical care. Follow your doctor's instructions very closely.

Storing unopened (not in use) pens: Keep the medicine in its original container. Refrigerate and use until [expiration date](#). Protect from light.

Do not freeze. Throw away the medicine if it has been frozen.

Storing opened (in use) pens: Store at room temperature and use within 30 days. Protect from heat and bright light. Do not store the injection pen with a needle attached.

Dosing information

Usual Adult Dose for Diabetes Type 2:

Immediate-release (Byetta):

Initial dose: 5 mcg subcutaneously twice a day within the 60-minute period before the morning and evening meals

-Maintenance dose: After 1 month, may increase dose to 10 mcg subcutaneously twice a day based on clinical response

-Maximum dose: 10 mcg twice a day

-Alternatively, may give before the two main meals of the day provided dosing is 6 hours or more apart

Comments:

-This drug has not been studied in patients with a history of pancreatitis and therefore, other therapies should be considered in these patients.

-Extended-release exenatide is not recommended as first-line therapy because of the uncertainty of the risk of thyroid C-cell tumors.

-Patients may experience transient (approximately 2 to 4 weeks) elevation in blood sugar when switching from immediate-release exenatide to extended-release.

Use: As an adjunct to diet and exercise to improve glycemic control in adults with type 2 [diabetes mellitus](#).

 [Detailed Byetta dosage information](#)

What happens if I miss a dose?

Skip the missed dose and use your next dose at the regular time. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose can cause severe [nausea](#) and [vomiting](#), or signs of low blood sugar ([headache](#), hunger, irritability, [dizziness](#),

feeling shaky).

What should I avoid while using Byetta?

You should not mix **Byetta** with insulin in the same syringe. **Do not use Byetta together with Bydureon.**

Avoid drinking alcohol. It can lower your blood sugar.

Byetta side effects

Get emergency medical help if you have **signs of an allergic reaction to Byetta**: [hives](#), itching; rapid heartbeats; difficult breathing; feeling light-headed; swelling of your face, lips, tongue, or throat.

Some people using exenatide have had serious or fatal bleeding caused by low levels of platelets (blood cells that help your blood to clot). **Stop using Byetta and call your doctor right away if you have unusual bleeding or bruising.**

Stop using Byetta and call your doctor at once if you have:

- **pancreas or gallbladder problems** - pain in your upper stomach spreading to your back, nausea and vomiting, fever, [fast heart rate](#), yellowing of your skin or eyes;
- **low blood sugar** - headache, hunger, sweating, irritability, dizziness, nausea, fast heart rate, and feeling anxious or shaky; or
- **kidney problems** - little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath.

Common Byetta side effects include:

- [indigestion](#), nausea, vomiting;
- [diarrhea](#), [constipation](#); or
- itching or a small bump where the medicine was injected.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Byetta side effects](#) (more detail)

Related/similar drugs

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Rybelsus

Rybelsus tablets are used to improve blood sugar control in adults with type 2 diabetes, and may ...

Reviews & ratings

6.4 / 10

[282 Reviews](#)

[View more](#)

FEATURED

Mounjaro

Mounjaro (tirzepatide) is FDA-approved to manage type 2 diabetes in adults. It lowers blood sugar ...

Reviews & ratings

8.5 / 10

[1,278 Reviews](#)

[View more](#)

Trulicity

Trulicity is an injectable diabetes medicine that is used together with diet and exercise to ...

Reviews & ratings

5.3 / 10

[970 Reviews](#)

[View more](#)

Lantus

Lantus is a long acting form of insulin used to treat type 1 or type 2 diabetes. Learn about side ...

Reviews & ratings

6.5 / 10

[41 Reviews](#)

[View more](#)

Tresiba

Tresiba (insulin degludec) is used to treat diabetes mellitus. Includes Tresiba side effects ...

Reviews & ratings

4.3 / 10

[152 Reviews](#)

[View more](#)

Victoza

Victoza helps control blood sugar levels and reduce the risk of serious heart problems in people ...

Reviews & ratings

7.6 / 10

[766 Reviews](#)

[View more](#)

Basaglar

Basaglar (insulin glargine) is a long-acting insulin that is used to improve blood sugar control in ...

Reviews & ratings

2.7 / 10

[148 Reviews](#)

[View more](#)

Levemir

Levemir (insulin detemir) is a long acting insulin used to treat diabetes in adults and children ...

Reviews & ratings

4.6 / 10

[70 Reviews](#)

[View more](#)

Novolog

NovoLog is a fast-acting insulin used to treat diabetes to control blood sugar

levels in adults and ...

Reviews & ratings

5.5 / 10

[32 Reviews](#)

[View more](#)

What other drugs will affect Byetta?

If you take the following medicines, take them at least 1 hour before you use Byetta:

- an [antibiotic](#); or
- birth control pills.

If you take these medicines with food, take them with a meal or snack separate from the meal times when you use Byetta.

Exenatide can make it harder for your body to absorb other medicines you take by mouth. Tell your doctor about all your current medicines and any you start or stop using, especially:

- blood pressure medicine;
- a diuretic or "water pill";
- insulin or oral diabetes medicine;
- pain medicine;
- [lovastatin](#) (Advicor, Altoprev, [Mevacor](#)); or
- [warfarin](#) ([Coumadin](#), [Jantoven](#)).

This list is not complete. Other drugs may interact with exenatide, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible drug interactions are listed here.

 [Byetta drug interactions](#) (more detail)

Does Byetta interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Byetta

+

Enter a drug name

Add

Popular FAQ

How long does it take for Bydureon Bcise to take effect?



How long can Bydureon be unrefrigerated?



How do you use the Bydureon pen?



How does Bydureon work?



How do you use the Bydureon Bcise autoinjector?



More FAQ

- [Is there a list of drugs that require cold storage?](#)

References

1. [Byetta Product Label](#)

More about Byetta (exenatide)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(82\)](#)
- [Latest FDA alerts \(4\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [FDA approval history](#)
- [Drug class: GLP-1 Agonists \(Incretin Mimetics\)](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

Other brands

[Bydureon](#), [Bydureon BCise](#)

Professional resources

- [Byetta prescribing information](#)
- [Exenatide \(AHFS Monograph\)](#)

Other brands

[Bydureon](#), [Bydureon BCise](#)

Related treatment guides

- [Type 2 Diabetes](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Byetta only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 14.01.

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	FDA approved 2005

User Reviews & Ratings

7.8 / 10

[82 Reviews](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.