

Home

Coreg

Coreg 🖘

Generic name: carvedilol [KAR-ve-dil-ole]

Brand names: Coreg, Coreg CR

Drug class: Non-cardioselective beta blockers

Medically reviewed by Philip Thornton, DipPharm. Last updated on Jun 4, 2024.

Uses Warnings Before taking Dosage Side effects Interactions

What is Coreg?

Coreg is a beta-blocker. Beta-blockers affect the heart and circulation (blood flow through arteries and veins).

Coreg is used to treat heart failure and hypertension (high blood pressure).

Coreg is also used after a heart attack that has caused your heart not to pump as well.

Warnings

You should not take Coreg if you have asthma, bronchitis, emphysema, severe liver disease, or a serious heart condition such as heart block, "sick sinus syndrome," or slow heart rate (unless you have a pacemaker).

Avoid drinking alcohol within 2 hours before or after taking extended-release Coreg CR capsules. Also avoid taking medicines or other products that might contain alcohol. Alcohol may cause the carvedilol in the controlled release (CR) capsule to be released too quickly into the body.

If you are being treated for high blood pressure, keep using Coreg even if you feel well. High blood pressure often has no symptoms. You may need to use blood pressure medication for the rest of your I

Before taking this medicine

You should not take Coreg if you are allergic to carvedilol, or if you have:

- · asthma, bronchitis, emphysema;
- severe liver disease; or
- a serious heart condition such as severe heart failure, heart block, "sick sinus syndrome," or slow heart rate (unless you have a pacemaker).

To make sure Coreg is safe for you, tell your doctor if you have:

- coronary artery disease (clogged arteries);
- · slow heartbeats that have caused you to faint;
- · fluid retention;
- · asthma or other lung problems;
- angina (chest pain);
- diabetes (taking carvedilol can make it harder for you to tell when you have low blood sugar);
- a thyroid disorder;
- · kidney disease;
- · circulation problems (such as Raynaud's syndrome); or
- pheochromocytoma (tumor of the adrenal gland).

Tell your doctor if you are pregnant or breastfeeding.

Coreg is not approved for use by anyone younger than 18 years old.

1 Coreg pregnancy and breastfeeding warnings (more detail)

How should I take Coreg?

Take Coreg exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

Coreg works best if you take it with food, at the same time every day.

Swallow the extended-release capsule whole and do not crush, chew, break, or open it.

If you cannot swallow a capsule whole, open it and sprinkle the medicine into a spoonful of cold applesauce. Swallow the mixture right away without chewing. Do not save it for later use.

If you are switched from the tablets to Coreg CR extended-release capsules, your daily total dose of this medicine may be higher or lower than before. Older adults may be more likely to become dizzy or feel faint when switching from tablets to extended-release capsules. Follow your doctor's instructions.

Your blood pressure will need to be checked often.

If you need surgery (including cataract surgery), tell your surgeon you currently use this medicine. You may need to stop for a short time.

You should not stop using Coreg suddenly. Stopping suddenly may cause chest pain or a heart attack. Follow your doctor's instructions about tapering your dose.

If you are being treated for high blood pressure, keep using this medication even if you feel well. High blood pressure often has no symptoms. You may need to use blood pressure medication for the rest of your life.

Coreg is only part of a complete treatment program that may also include diet, exercise, and weight control. Follow your doctor's instructions very closely.

Store at room temperature away from moisture and heat.

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include uneven heartbeats, shortness of breath, bluish-colored fingernails, dizziness, weakness, fainting, and seizure (convulsions).

What to avoid

Avoid driving or hazardous activity until you know how this medicine will affect you. Your reactions could be impaired. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

Coreg side effects

Get emergency medical help if you have **signs of an allergic reaction to Coreg:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- · a light-headed feeling, like you might pass out;
- slow or uneven heartbeats;
- cold feeling or numbness in your fingers or toes;
- chest pain, dry cough, wheezing, chest tightness;
- heart problems swelling, rapid weight gain, feeling short of breath; or
- high blood sugar increased thirst, increased urination, dry mouth, fruity breath odor.

Common Coreg side effects may include:

- · dizziness;
- slow heartbeats;
- diarrhea;
- · weight gain;
- dry eyes; or
- problems wearing contact lenses.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.



1 Coreg side effects (more detail)

Related/similar drugs

Hydrochlorothiazide

HCTZ (hydrochlorothiazide) used to treat high blood pressure (hypertension) and edema. Includes ...

Reviews & ratings

5.2 / 10

185 Reviews

View more

Propranolol

Propranolol is a beta-blocker that is used to treat tremors, chest pain, high blood pressure, heart ...

Reviews & ratings

8.2 / 10

1,376 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Aspirin

Aspirin is used to treat mild to moderate pain and to reduce fever or inflammation. Learn about ...

Reviews & ratings

7.8 / 10

51 Reviews

View more

Spironolactone

Spironolactone is a potassium-sparing diuretic that is primarily used to treat heart failure, high ...

Reviews & ratings

6.9 / 10

702 Reviews

View more

Furosemide

Furosemide is a loop diuretic used to treat fluid retention and high blood pressure by increasing ...

Reviews & ratings

6.8 / 10

121 Reviews

View more

Losartan

Losartan is used to treat high blood pressure (hypertension). It is also used to lower the risk of ...

Reviews & ratings

4.5 / 10

578 Reviews

View more

Metoprolol

Metoprolol is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

Reviews & ratings

5.8 / 10

653 Reviews

View more

Jardiance

Jardiance (empagliflozin) is used to treat type 2 diabetes, chronic kidney disease and reduce the ...

Reviews & ratings

5.3 / 10

325 Reviews

View more

Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

Reviews & ratings

6.4 / 10

56 Reviews

View more

What other drugs will affect Coreg?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Other drugs may interact with carvedilol, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

1 Coreg drug interactions (more detail)

Does Coreg interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Coreg
+
Enter a drug name
Add

More about Coreg (carvedilol)

· Check interactions

- Compare alternatives
- · Pricing & coupons
- Reviews (44)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- Support group
- Drug class: non-cardioselective beta blockers
- Breastfeeding
- En español

Professional resources

- Coreg prescribing information
- Carvedilol (AHFS Monograph)

Other formulations

Coreg CR

Related treatment guides

- Atrial Fibrillation
- Left Ventricular Dysfunction
- High Blood Pressure
- Angina
- Heart Failure

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Coreg only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 16.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

্প Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+ years FDA approved 1995

WADA Class

്റ്റ Anti-Doping Classification

User Reviews & Ratings

7.2 / 10

44 Reviews

Images

Coreg 6.25 mg (SB 4140)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us



<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright $\ensuremath{@}$ 2000-2025 Drugs.com. All rights reserved.