

Home

2. Ceftriaxone

Ceftriaxone (injection) ⊲ ¬

Generic name: ceftriaxone (injection) [*SEF-trye-AX-one*] **Brand names:** Rocephin, Rocephin ADD-Vantage

Dosage forms: injectable powder for injection (1 g; 10 g; 2 g; 250 mg; 500 mg), intravenous solution (1 g/50 mL-iso-osmotic dextrose; 2

g/50 mL-iso-osmotic dextrose)

Drug class: Third generation cephalosporins



Medically reviewed by Drugs.com on Aug 8, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is ceftriaxone?

Ceftriaxone is a cephalosporin (SEF a low spor in) antibiotic that is used to treat many kinds of bacterial infections, including severe or life-threatening forms such as E. coli, pneumonia, or meningitis. Ceftriaxone is also used to prevent infection in people having certain types of surgery.

Ceftriaxone may also be used for purposes not listed in this medication guide.

Ceftriaxone side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, sore throat, burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling).

Ceftriaxone may cause serious side effects. Call your doctor at once if you have:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose);
- new signs of infection (fever, chills, sweating);
- nausea, vomiting, pain in your upper stomach that spreads to your back;
- pale or yellowed skin, dark colored urine;
- new or worsening breathing problems (wheezing, feeling short of breath);
- a blood cell disorder--headache, chest pain, dizziness, weakness, severe tingling or numbness; or
- kidney or bladder problems--pain in your side or lower back spreading to your groin, blood in your urine, painful
 or difficult urination, little or no urine.

Common side effects of ceftriaxone may include:

- · symptoms of a blood cell disorder;
- diarrhea;
- · vaginal itching or discharge;
- warmth, tight feeling, or a hard lump where the injection was given;
- · rash; or
- · abnormal liver function tests.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i Ceftriaxone side effects (more detail)

Related/similar drugs

Levofloxacin

Levofloxacin is a fluoroquinolone antibiotic used to treat serious bacterial infections and prevent ...

Reviews & ratings

5.1 / 10

741 Reviews

View more

Clindamycin

Clindamycin (Cleocin) is used to treat serious infections caused by bacteria. Includes clindamycin ...

Reviews & ratings

5.7 / 10

806 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Augmentin

Augmentin is a prescription antibiotic combining amoxicillin and clavulanate to treat bacterial ...

Reviews & ratings

5.4 / 10

447 Reviews

View more

Hydroxychloroquine

Hydroxychloroquine is a quinoline drug used to treat or prevent malaria. It's also used to treat ...

Reviews & ratings

6.4 / 10

356 Reviews

View more

Ciprofloxacin

Ciprofloxacin is an antibiotic belong to a group of drugs called fluoroquinolones. Learn about side ...

Reviews & ratings

5.0 / 10

935 Reviews

View more

Metronidazole

Metronidazole is an antibiotic used to fight bacteria in your body. Learn about side effects ...

Reviews & ratings

6.0 / 10

2,431 Reviews

View more

Cephalexin

Cephalexin is a cephalosporin antibiotic used to treat bacterial infections like respiratory, skin ...

Reviews & ratings

6.4 / 10

509 Reviews

View more

Warnings

You should not use ceftriaxone if you have ever had a severe allergic reaction to any type of cephalosporin antibiotic (Omnicef, Keflex, and others).

Do not use ceftriaxone in a child without a doctor's advice, and never give more than the child's prescribed dose.

Before taking this medicine

You should not take ceftriaxone if you are allergic to cefazolin or any other cephalosporin antibiotic (cefdinir, cefalexin, Keflex, Omnicef, and others).

Do not use ceftriaxone in a child without a doctor's advice, and never give more than the child's prescribed dose. Ceftriaxone injection could be dangerous when given to a newborn baby with any intravenous medicines that contain calcium, including total parental nutrition (TPN).

Tell your doctor if you have ever had:

- liver or kidney disease;
- gallbladder disease;
- · diabetes; or
- bleeding problems.

Tell your doctor if you are pregnant or breastfeeding.

How is ceftriaxone given?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Ceftriaxone is injected into a muscle or as an infusion into a vein (IV). A healthcare provider will give you this injection when ceftriaxone is used **to prevent** infection from surgery. You may be shown how to use the injection at home **to treat** an infection.

Read and carefully follow any Instructions for Use provided with your medicine. You may need to mix ceftriaxone with a liquid (diluent) before using it. When using injections by yourself, be sure you understand how to properly mix and

store the medicine.

Prepare an injection only when you are ready to give it. **Do not use if the medicine has changed colors or has particles in it.** Call your pharmacist for new medicine.

Do not mix ceftriaxone in the same injection with other antibiotics, or with any diluent that contains calcium, including a TPN (total parenteral nutrition) solution.

If you use other injectable medications, be sure to flush your intravenous catheter between injections of each medication.

Drink plenty of liquids to keep your kidneys working properly while using ceftriaxone.

Use this medicine for the full prescribed length of time, even if your symptoms quickly improve. Skipping doses can increase your risk of infection that is resistant to medication. Ceftriaxone will not treat a viral infection such as the flu or a common cold.

Ceftriaxone can affect the results of certain medical tests. Tell any doctor who treats you that you are using ceftriaxone.

After mixing ceftriaxone with a diluent, you must use the medicine within a certain number of hours or days. This will depend on the diluent and how you store the mixture (at room temperature, in a refrigerator, or frozen). Carefully follow all mixing and storage instructions provided with your medicine. Ask your pharmacist if you have questions.

If your medicine was provided in a frozen form, thaw it in a refrigerator or at room temperature. Do not warm in a microwave or boiling water. Use the medicine as soon as possible after thawing it. Do not refreeze.

Use a needle and syringe only once and then place them in a puncture-proof "sharps" container. Follow state or local laws about how to dispose of this container. Keep it out of the reach of children and pets.

Detailed Ceftriaxone dosage information

What happens if I miss a dose?

Call your doctor for instructions if you miss a dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while using ceftriaxone?

Antibiotic medicines can cause diarrhea, which may be a sign of a new infection. If you have diarrhea that is watery or bloody, call your doctor. Do not use anti-diarrhea medicine unless your doctor tells you to.

What other drugs will affect ceftriaxone?

Tell your doctor about all your other medicines, especially:

fluconazole:

- · vancomycin; or
- other injected (IV) antibiotics.

This list is not complete. Other drugs may affect ceftriaxone, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

<u>Ceftriaxone drug interactions</u> (more detail)

Does ceftriaxone interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Frequently asked questions

• Pregnancy - is it safe to take ceftriaxone during pregnancy?

More about ceftriaxone

- Check interactions
- · Compare alternatives
- · Pricing & coupons
- Reviews (175)
- Side effects
- Dosage information
- During pregnancy
- Drug class: third generation cephalosporins
- Breastfeeding
- En español

Patient resources

Other brands

Rocephin

Professional resources

CefTRIAXone monograph

- Ceftriaxone (FDA)
- Ceftriaxone and Dextrose (FDA)

Other brands

Rocephin

Related treatment guides

- Bacterial Infection
- Bone infection
- Bacterial Endocarditis Prevention
- Bacteremia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 13.01.

Antibiotics 101

List of Common Antibiotics & Types

Antibiotics and Alcohol

Probiotics and Antibiotics

Antibiotics for Strep Throat

Antibiotics for a Tooth Infection

Antibiotics for UTI's

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

൞ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

8.5 / 10

175 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.