

Home

Dextromethorphan

Dextromethorphan ⊲_n

Generic name: dextromethorphan [dex-troe-meth-OR-fan]

Brand names: Babee Cof, Benylin DM Pediatric, Buckleys Mixture, Creomulsion, Creo-Terpin, ... show all 17 brands

Drug class: Antitussives

Medically reviewed by Kaci Durbin, MD. Last updated on Feb 24, 2025.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is dextromethorphan?

Dextromethorphan is a cough suppressant. It affects the signals in the brain that trigger cough reflex.

Dextromethorphan is used to treat a cough. It is available over-the-counter alone and is also present in many over-the-counter and prescription combination medications.

Dextromethorphan will not treat a cough that is caused by smoking, asthma, or emphysema.

Warnings

Do not give dextromethorphan to a child younger than 4 years old.

Always ask a doctor before giving a cough or cold medicine to a child. Death can occur from the misuse of cough and cold medicines in very young children.

Do not use dextromethorphan if you have used an MAO inhibitor such as isocarboxazid (Marplan), phenelzine (Nardil), rasagiline (Azilect), selegiline (Eldepryl, Emsam), tranylcypromine (Parnate), or methylene blue injection within the past 14 days. Serious, life-threatening side effects can occur if you take dextromethorphan before the MAO inhibitor has cleared from your body.

Do not use any other over-the-counter cough, cold, or allergy medication without first asking your doctor or pharmacist. If you take certain products together you may accidentally take too much of one or more types of medicine. Read the label of any other medicine you are using to see if it contains dextromethorphan. Dextromethorphan will not treat a cough that is caused by smoking, asthma, or emphysema.

Before taking this medicine

Do not use dextromethorphan if you have used an MAO inhibitor such as isocarboxazid (Marplan), phenelzine (Nardil), rasagiline (Azilect), selegiline (Eldepryl, Emsam), tranylcypromine (Parnate), or methylene blue injection within the past 14 days. Serious, life-threatening side effects can occur if you take dextromethorphan before the MAO inhibitor has

cleared from your body.

Ask a doctor or pharmacist about using dextromethorphan if you have emphysema or chronic bronchitis.

It is unknown whether dextromethorphan is harmful to an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during treatment. Dextromethorphan may pass into breast milk. Do not use this medication without telling your doctor if you are breast-feeding a baby.

Artificially-sweetened liquid forms of cough medicine may contain phenylalanine. This would be important to know if you have phenylketonuria (PKU). Check the ingredients and warnings on the medication label if you are concerned about phenylalanine.

1 Dextromethorphan pregnancy and breastfeeding warnings (more detail)

How should I take dextromethorphan?

Use dextromethorphan exactly as directed on the label, or as it has been prescribed by your doctor. Do not use the medication in larger amounts, or use it for longer than recommended. Cough medicine is usually taken only for a short time until your symptoms clear up.

Do not give dextromethorphan to a child younger than 4 years old.

Always ask a doctor before giving cough or cold medicine to a child. Death can occur from the misuse of cough or cold medicine in very young children.

Measure the liquid form of dextromethorphan with a special dose-measuring spoon or cup, not a regular table spoon. If you do not have a dose-measuring device, ask your pharmacist for one.

Allow the dextromethorphan lozenge or disintegrating strip to dissolve in your mouth.

Drink extra fluids to help loosen the congestion and lubricate your throat while you are taking this medication. Talk with your doctor if your symptoms do not improve after 7 days of treatment, or if you have a fever with a headache, cough, or skin rash.

If you need to have any type of surgery, tell the surgeon ahead of time if you have taken dextromethorphan within the past few days.

Store dextromethorphan at room temperature, away from heat, light, and moisture.

Detailed Dextromethorphan dosage information

What happens if I miss a dose?

Since cough medicine is usually taken only as needed, you may not be on a dosing schedule. If you are taking the medication regularly, take the missed dose as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at your next regularly scheduled time. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention if you think you have used too much of this medicine.

Overdose symptoms may include nausea, vomiting, drowsiness, dizziness, difficulty breathing, fast heartbeat, or seizures (convulsions). Overdose can also result in coma or death.

What should I avoid?

Avoid drinking alcohol. It can increase some of the side effects of dextromethorphan. This medication can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert.

Avoid taking diet pills, caffeine pills, or other stimulants (such as ADHD medications) without your doctor's advice. Taking a stimulant together with cough medicine can increase your risk of unpleasant side effects.

Do not use any other over-the-counter cough, cold, or allergy medication without first asking your doctor or pharmacist. Dextromethorphan is contained in many combination medicines available over the counter. If you take certain products together you may accidentally take too much of this medicine. Read the label of any other medicine you are using to see if it contains dextromethorphan.

Dextromethorphan side effects

Get emergency medical help if you have any of these signs of an allergic reaction: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Stop using dextromethorphan and call your doctor at once if you have any of these serious side effects:

- · severe dizziness, anxiety, restless feeling, or nervousness;
- · seizures or convulsions;
- · confusion, hallucinations; or
- · slow, shallow breathing.

Less serious side effects are more likely, such as stomach upset.

This is not a complete list of dextromethorphan side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Dextromethorphan side effects (more detail)

Related/similar drugs

Benadryl

Benadryl is an antihistamine and is used to treat sneezing, runny nose, itching and other ...

Reviews & ratings

7.0 / 10

79 Reviews

View more

Mucinex

Mucinex (guaifenesin) is used to relieve the symptoms of cough and loosen mucus in the chest ...

Reviews & ratings

6.2 / 10

155 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Norco

Norco (acetaminophen and hydrocodone) is used to relieve moderate to severe pain. Includes Norco ...

Reviews & ratings

6.3 / 10

367 Reviews

View more

Dilaudid

Dilaudid (hydromorphone) is a narcotic pain reliever used to treat moderate to severe pain ...

Reviews & ratings

8.2 / 10

329 Reviews

View more

Vicodin

Vicodin (acetaminophen and hydrocodone) is used to relieve moderate to severe pain. Includes ...

Reviews & ratings

7.7 / 10

146 Reviews

View more

Codeine

Codeine is a opioid pain medicine used to treat mild to moderate pain. Includes codeine side ...

Reviews & ratings

7.8 / 10

70 Reviews

View more

Guaifenesin

Guaifenesin is an expectorant. It helps loosen congestion in your chest and throat. Includes ...

Reviews & ratings

7.1 / 10

305 Reviews

View more

Diphenhydramine

Diphenhydramine is an antihistamine used to treat sneezing, runny nose, itching, hives and other ...

Reviews & ratings

5.5 / 10

541 Reviews

View more

Before taking dextromethorphan, tell your doctor if you are using any of the following drugs:

- celecoxib (Celebrex);
- cinacalcet (Sensipar);
- · darifenacin (Enablex);
- imatinib (Gleevec);
- quinidine (Quinaglute, Quinidex);
- ranolazine (Ranexa)
- ritonavir (Norvir);
- sibutramine (Meridia);
- · terbinafine (Lamisil);
- medicines to treat high blood pressure; or
- antidepressant medications such as amitriptyline (Elavil, Etrafon), bupropion (Wellbutrin, Zyban), fluoxetine (Prozac, Sarafem), fluoxamine (Luvox), imipramine (Janimine, Tofranil), paroxetine (Paxil), sertraline (Zoloft), and others.

This list is not complete and there are numerous other drugs that can interact with dextromethorphan. Tell your doctor about all the prescription and over-the-counter medications you use. This includes vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start using a new medication without telling your doctor.

Dextromethorphan drug interactions (more detail)

Does dextromethorphan interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

dextromethorphan
+
Enter a drug name
Add

Popular FAQ

Does Mucinex help with Covid?

More FAQ

• Dextromethorphan - can you give TusQ Dx to a child and what is the dose?

More about dextromethorphan

Check interactions

- · Compare alternatives
- Pricing & coupons
- Reviews (386)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Support group
- Drug class: antitussives
- Breastfeeding

Patient resources

Other brands

DexAlone, Robitussin CoughGels Long-Acting, Vicks Formula 44, Creomulsion, ... +6 more

Professional resources

• Dextromethorphan monograph

Other brands

Delsym 12 Hour Cough Relief

Related treatment guides

Cough

Further information

What does my medication look like?

Dextromethorphan is available over the counter under many different brand and generic names. Ask your pharmacist any questions you have about this medication, especially if it is new to you.

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.04.

DRUG STATUS

Availability

or Rx and/or OTC

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

The Drug history at FDA

User Reviews & Ratings

3.7 / 10

386 Reviews

Images

Dextromethorphan Hydrobromide 15 mg (778)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×