

[Home](#)

2. Guanfacine

# Guanfacine

**Generic name:** guanfacine [ *GWAHN-fa-seen* ]**Brand names:** [Intuniv](#), [Tenex](#)**Dosage forms:** oral tablet (1 mg; 2 mg), oral tablet, extended release (1 mg; 2 mg; 3 mg; 4 mg)**Drug class:** [Antiadrenergic agents, centrally acting](#)[Medically reviewed](#) by Drugs.com on May 6, 2024. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

## What is guanfacine?

Guanfacine is used to treat [high blood pressure](#) (hypertension). It is sometimes given with other blood pressure medications.

The **Intuniv** brand of guanfacine is used to treat attention deficit hyperactivity disorder (ADHD).

Guanfacine may also be used for purposes not listed in this medication guide.

## Guanfacine side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

**Guanfacine may cause serious side effects. Call your doctor at once if you have:**

- [anxiety](#), nervousness;
- hallucinations (especially in children);
- severe [drowsiness](#);
- slow heartbeats; or
- a light-headed feeling, like you might pass out;

**If you stop taking guanfacine**, tell your doctor if you have headaches, confusion, rapid heartbeats, tremors, increased blood pressure, or if you feel nervous or agitated. If left untreated, these symptoms could lead to very high blood pressure, vision problems, or [seizures](#).

**Common side effects of guanfacine may include:**

- dizziness, drowsiness;

- [low blood pressure](#), slow heartbeats;
- feeling tired or irritable;
- trouble sleeping;
- [dry mouth](#); or
- stomach pain, nausea, [constipation](#).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Guanfacine side effects](#) (more detail)

## Related/similar drugs

### Adderall

Adderall (amphetamine and dextroamphetamine) is used to treat narcolepsy and ADHD. Includes side ...

#### Reviews & ratings

7.2 / 10

[505 Reviews](#)

[View more](#)

### Hydrochlorothiazide

HCTZ (hydrochlorothiazide) used to treat high blood pressure (hypertension) and edema. Includes ...

#### Reviews & ratings

5.2 / 10

[185 Reviews](#)

[View more](#)

#### FEATURED

### Omvox

Omvox is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

#### Reviews & ratings

[1 Reviews](#)

[View more](#)

## Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

### Reviews & ratings

6.4 / 10

[56 Reviews](#)

[View more](#)

## Furosemide

Furosemide is a loop diuretic used to treat fluid retention and high blood pressure by increasing ...

### Reviews & ratings

6.8 / 10

[121 Reviews](#)

[View more](#)

## Losartan

Losartan is used to treat high blood pressure (hypertension). It is also used to lower the risk of ...

### Reviews & ratings

4.5 / 10

[578 Reviews](#)

[View more](#)

## Metoprolol

Metoprolol is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

### Reviews & ratings

5.8 / 10

[653 Reviews](#)

[View more](#)

## Atenolol

Atenolol is used to treat angina (chest

... is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

#### Reviews & ratings

7.2 / 10

[276 Reviews](#)

[View more](#)

### Valsartan

Valsartan is an angiotensin II receptor blocker (ARB) that may be used to treat high blood pressure ...

#### Reviews & ratings

5.7 / 10

[152 Reviews](#)

[View more](#)

### Vyvanse

Vyvanse is a prescription stimulant used to treat ADHD in children and adults, and binge eating ...

#### Reviews & ratings

7.4 / 10

[973 Reviews](#)

[View more](#)

## Warnings

Use only as directed. Tell your doctor if you use other medicines or have other medical conditions or allergies.

## Before taking this medicine

You should not use guanfacine if you are allergic to it.

**Intuniv** is not approved for use by anyone younger than 6 years old. **Guanfacine** is not approved for use by anyone younger than 12 years old.

Tell your doctor if you have ever had:

- heart problems, [coronary artery disease](#) (clogged arteries);
- a heart rhythm disorder;

- a [heart attack](#) or stroke;
- high or low blood pressure;
- [liver disease](#); or
- kidney disease.

It is not known whether guanfacine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It may not be safe to breastfeed a baby while you are using this medicine. Ask your doctor about any risks.

## How should I take guanfacine?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Avoid taking guanfacine with high-fat foods, or your body could absorb the medicine too quickly.

Take **Intuniv** with a full glass of water, milk, or other liquid.

Take **Guanfacine** at bedtime to ease drowsiness.

Swallow the **Intuniv** tablet whole and do not crush, chew, or break it.

If a child is using this medicine, tell your doctor if the child has any changes in weight. **Intuniv** doses are based on weight in children, and any changes may affect your child's dose.

**You should not stop using guanfacine suddenly.** Stopping suddenly can raise your blood pressure and cause unpleasant symptoms.

Call your doctor if you are sick with [vomiting](#) and cannot take your medicine as usual.

Your dose needs may change if you switch to a different brand, strength, or form of this medicine. **Avoid medication errors by using only the form and strength your doctor prescribes.**

Your doctor will need to check your progress on a regular basis. Your blood pressure and heart rate may also need to be checked.

If you have high blood pressure, **keep using guanfacine even if you feel well.** High blood pressure often has no symptoms. You may need to use blood pressure medicine for the rest of your life.

Store at room temperature away from moisture, heat, and light.

 [Guanfacine patient tips](#) (more detail)

## Guanfacine dosing information

### Usual Adult Dose for Hypertension:

Initial dose: 1 mg orally once a day at bedtime; may increase to 2 mg once a day if satisfactory result not achieved after 3 to 4 weeks.

Comments: Adverse reactions increase significantly with doses above 3 mg/day.

#### **Usual Pediatric Dose for Attention Deficit Disorder:**

##### **6 YEARS TO LESS THAN 18 YEARS:**

Initial dose: 1 mg orally once a day, either in the morning or evening, at approximately the same time each day; may adjust in increments of no more than 1 mg/week.

Recommended target dose: 0.05 to 0.12 mg/kg/day (total daily dose between 1 and 7 mg) once a day, depending on clinical response and tolerability

Maximum dose: 6 to 12 years: Doses above 4 mg/day have not been evaluated; 13 to 17 years: Doses above 7 mg/day have not been evaluated.

Comments: Attention Deficit Hyperactivity Disorder (ADHD) may require treatment for an extended period of time; healthcare providers should periodically reevaluate treatment and adjust dose as needed.

Use: Treatment of ADHD as monotherapy and as adjunctive therapy to stimulant medications.

#### **Usual Pediatric Dose for Hypertension:**

##### **12 YEARS OR OLDER:**

Initial dose: 1 mg orally once a day at bedtime; may increase to 2 mg once a day if satisfactory result not achieved after 3 to 4 weeks.

Comments: Adverse reactions increase significantly with doses above 3 mg/day.

 [Detailed Guanfacine dosage information](#)

## **What happens if I miss a dose?**

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

Call your doctor for instructions if you miss more than 2 doses in a row.

## **What happens if I overdose?**

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe drowsiness, [slow heart rate](#), and feeling like you might pass out.

## **What should I avoid while taking guanfacine?**

Avoid driving or hazardous activity until you know how guanfacine will affect you. Your reactions could be impaired.

Drinking alcohol can increase certain side effects of guanfacine.

## What other drugs will affect guanfacine?

**Sometimes it is not safe to use certain medications at the same time.** Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Using guanfacine with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety, [depression](#), or seizures.

Tell your doctor about all your other medicines. Some may affect guanfacine, especially:

- [ketoconazole](#);
- a barbiturate, such as [phenobarbital](#);
- blood pressure medications;
- medicine to treat mental illness; or
- a sedative, such as [Valium](#) or Xanax.

This list is not complete. Other drugs may affect guanfacine, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible [drug interactions](#) are listed here.

 [Guanfacine drug interactions](#) (more detail)

## Does guanfacine interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

guanfacine

+

Enter a drug name

Add

## More about guanfacine

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(317\)](#)
- [Drug images](#)
- [Latest FDA alerts \(1\)](#)
- [Side effects](#)
- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)

- [Support group](#)
- [Drug class: antiadrenergic agents, centrally acting](#)
- [Breastfeeding](#)
- [En español](#)

## Patient resources

- [Guanfacine Extended-Release Tablets patient information](#)
- [Guanfacine Tablets](#)

## Other brands

[Intuniv](#), [Tenex](#)

## Professional resources

- [Guanfacine monograph](#)
- [Guanfacine \(FDA\)](#)
- [Guanfacine ER \(FDA\)](#)

## Other brands

[Intuniv](#), [Tenex](#)

## Related treatment guides

- [ADHD](#)
- [Tourette's Syndrome](#)
- [High Blood Pressure](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 13.01.

### DRUG STATUS

#### Availability

**Rx** Prescription only



## Pregnancy & Lactation



Risk data available

## CSA Schedule\*

N/A Not a controlled drug

## Approval History



Drug history at FDA



## User Reviews & Ratings

6.1 / 10

[317 Reviews](#)

## Images

[Guanfacine Hydrochloride Extended-Release 1 mg \(APO GU1\)](#)



[View larger images](#)

# Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



## About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

## Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

## Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.