

Home

Yuvafem

Yuvafem (local)

Generic name: estradiol vaginal (local) [ES-tra-DYE-ole-VA-jin-ul]

Brand names: Estrace Vaginal, Estring, Imvexxy Maintenance Pack, Imvexxy Starter Pack, Vagifem, Yuvafem, Imvexxy local

Drug classes: Estrogens, Miscellaneous vaginal agents



Medically reviewed by Drugs.com on Dec 19, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Yuvafem (local)?

Estradiol is a form of estrogen, a female sex hormone that regulates many processes in the body.

Some estradiol products placed directly into the vagina are used for "local" treatment of vaginal menopause symptoms (such as dryness, burning, and irritation). Other vaginal estradiol products are used for treating vaginal menopause symptoms and symptoms that affect other parts of the body (such as hot flashes). This type of vaginal estradiol has "systemic" effects, meaning that it can affect parts of the body other than where the medicine is directly applied.

This medication guide provides information about Yuvafem for local treatment of vaginal symptoms of menopause (such as dryness, itching, irritation, and pain during sexual intercourse).

Yuvafem (local) may also be used for purposes not listed in this medication guide.

Yuvafem (local) side effects

Get emergency medical help if you have any of these **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Remove the vaginal ring and seek emergency medical attention if you have: fever with nausea, vomiting, diarrhea, muscle pain, dizziness, fainting, and/or sunburn-like skin rash. These may be signs of a life-threatening disease called toxic shock syndrome.

Yuvafem may cause serious side effects. Call your doctor at once if you have:

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance;
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood, pain or warmth in one or both legs;

- swelling or tenderness in your stomach;
- jaundice (yellowing of the skin or eyes);
- memory problems, confusion, unusual behavior;
- unusual vaginal bleeding, pelvic pain;
- a lump in your breast; or
- high levels of calcium in your blood--nausea, vomiting, constipation, increased thirst or urination, muscle weakness, bone pain, lack of energy.

Common side effects of Yuvafem may include:

- nausea, vomiting, bloating, stomach cramps;
- · headache:
- swelling in your hands or feet, weight gain;
- · breast pain; or
- vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i Yuvafem side effects (more detail)

Related/similar drugs

Premarin

Premarin is used to treat symptoms of menopause such as hot flashes, and vaginal dryness. Learn ...

Reviews & ratings

7.7 / 10

84 Reviews

View more

Estrace

Estrace (estradiol) is used to treat symptoms of menopause such as hot flashes and vaginal dryness ...

Reviews & ratings

7.3 / 10

12 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Prempro

Prempro (conjugated estrogens and medroxyprogesterone) is used to treat the symptoms of menopause ...

Reviews & ratings

6.9 / 10

66 Reviews

View more

Vagifem

Vagifem vaginal inserts are used treat menopause symptoms such as dryness, burning, and irritation ...

Reviews & ratings

6.0 / 10

102 Reviews

View more

Estrogel

Estrogel treats symptoms of menopause such as hot flashes, vaginal dryness, burning, and ...

Reviews & ratings

Add a review

View more

Estradiol

Estradiol is used to treat symptoms of

menopause sucn as not tiasnes, and vaginal dryness, burning ...

Reviews & ratings

5.8 / 10

587 Reviews

View more

Warnings

You should not use estradiol if you have: undiagnosed vaginal bleeding, liver disease, a bleeding disorder, if you will have major surgery, or if you have ever had a heart attack, a stroke, a blood clot, or cancer of the breast, uterus/cervix, or vagina.

Do not use if you are pregnant.

Estradiol may increase your risk of developing a condition that may lead to uterine cancer. Report any unusual vaginal bleeding right away.

Using Yuvafem can increase your risk of blood clots, stroke, heart attack, or cancer of the breast, uterus, or ovaries. Estradiol should not be used to prevent heart disease, stroke, or dementia.

Before taking this medicine

You should not use estradiol if you are allergic to it, or if you have:

- unusual vaginal bleeding that has not been checked by a doctor;
- · liver disease:
- · a history of heart attack, stroke, or blood clot;
- an increased risk of having blood clots due to a heart problem or a hereditary blood disorder; or
- a history of hormone-related cancer, or cancer of the breast, uterus/cervix, or vagina.

Do not use estradiol if you are pregnant. Tell your doctor right away if you become pregnant during treatment.

Using this medicine can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have high blood pressure, diabetes, high cholesterol, if you are overweight, or if you smoke.

Estradiol should not be used to prevent heart disease, stroke, or dementia. This medicine may actually increase your risk of developing these conditions.

Tell your doctor if you have ever had:

- · heart disease;
- liver problems, or jaundice caused by pregnancy or taking hormones;
- · kidney disease;

- · gallbladder disease;
- · asthma;
- · epilepsy or other seizure disorder;
- · migraines;
- lupus;
- endometriosis or uterine fibroid tumors;
- hereditary angioedema;
- porphyria (a genetic enzyme disorder that causes symptoms affecting the skin or nervous system);
- a thyroid disorder; or
- high or low levels of calcium in your blood.

Using estradiol may increase your risk of cancer of the breast, uterus, or ovaries. Talk with your doctor about this risk.

Estradiol can slow breast milk production. Tell your doctor if you are breast-feeding.

How should I use Yuvafem (local)?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Estradiol may increase your risk of developing a condition that may lead to uterine cancer. Your doctor may prescribe a progestin to help lower this risk. **Call your doctor at once if you have any unusual vaginal bleeding.**

Wash your hands before and after inserting Yuvafem.

Use the applicator provided to measure the prescribed dose of estradiol **vaginal cream**. Take apart the **cream** applicator and wash it with mild soap and warm water after each use.

Each estradiol **vaginal tablet** is supplied in a single-use disposable applicator. Throw the **tablet** applicator away after one use.

You should not be able to feel the **vaginal ring** once it is in place. Leave the vaginal ring in place for 90 days, then remove it. Your doctor may want you to replace it with a new ring. The ring does not need to be removed during sexual intercourse. If the ring is bothersome, you may remove it, rinse it with warm water, and reinsert it after intercourse.

To remove the ring, loop a finger through the ring and gently pull it from the vagina. **Call your doctor if you have trouble removing a vaginal ring.**

Your doctor should check your progress on a regular basis to determine whether you should continue this treatment. Self-examine your breasts for lumps on a monthly basis and have a mammogram every year while using estradiol.

If you need major surgery or will be on long-term bed rest, you may need to stop using Yuvafem for a short time. Any doctor or surgeon who treats you should know that you are using estradiol.

Store at room temperature away from moisture and heat. Keep the vaginal ring in its protective pouch until you are ready to use it.

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

Remove the vaginal ring and insert a new one as soon as you remember. **Do not** use an extra vaginal ring to make up the missed wearing time.

If a vaginal ring falls out, rinse it with warm water and reinsert it. If it slides down into the lower part of the vagina, use your finger to push it in farther.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while using Yuvafem (local)?

Avoid smoking. It can greatly increase your risk of blood clots, stroke, or heart attack while using estradiol.

Avoid using other vaginal products without your doctor's advice.

Grapefruit may interact with estradiol and lead to unwanted side effects. Avoid the use of grapefruit products.

What other drugs will affect Yuvafem (local)?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Many drugs can affect estradiol. This includes prescription and over-the-counter medicines, vitamins, and herbal products. **Not all possible interactions are listed here.** Tell your doctor about all your current medicines and any medicine you start or stop using.

Yuvafem drug interactions (more detail)

Does Yuvafem interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Yuvafem	
+	
Enter a drug name	Add

More about Yuvafem (estradiol topical)

- Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (106)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: estrogens
- Breastfeeding
- En español

Patient resources

· Yuvafem patient information

Other brands

Vagifem, Imvexxy, Estrace Vaginal Cream, Estring

Professional resources

• Yuvafem prescribing information

Other brands

Estrace Vaginal Cream

Related treatment guides

- Atrophic Vaginitis
- Atrophic Urethritis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 10.02.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

প্ Risk data available

CSA Schedule*

N/A Not a controlled drug

User Reviews & Ratings

3.2 / 10

106 Reviews

Images

Yuvafem (vaginal insert) 10 mcg (AN 276)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and

natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

V