

Home

Dinoprostone

Dinoprostone topical

Generic name: dinoprostone topical [DYE-no-PROS-tone-TOP-ik-al]

Brand names: Cervidil, Prepidil, Prostin E2

Dosage forms: vaginal gel (0.5 mg/3 g), vaginal insert (10 mg)

Drug class: Uterotonic agents



Medically reviewed by Drugs.com on Aug 9, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is dinoprostone topical?

Dinoprostone is a prostaglandin, a hormone-like substance that is naturally produced by tissues in the body.

Dinoprostone topical is used in a pregnant woman to relax the muscles of the cervix (opening of the uterus) in preparation for inducing labor at the end of a pregnancy.

Dinoprostone topical may be used for purposes not listed in this medication guide.

Dinoprostone topical side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Tell your caregivers at once if your contractions slow down or become uneven, or if you have:

- · a light-headed feeling, like you might pass out;
- weak or shallow breathing;
- intense pain between contractions;
- · sudden vaginal bleeding;
- · unexpected stomach pain;
- · chest pain;
- · easy bruising, unusual bleeding;
- bleeding from a wound, surgical incision, or vein where an IV needle was placed; or
- any bleeding that will not stop.

Common side effects of dinoprostone topical may include:

- · contractions that are more frequent;
- slow heartbeats in the baby;
- nausea, stomach pain;
- · feeling of warmth in the vaginal area;
- · back pain; or
- fever.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i <u>Dinoprostone topical side effects</u> (more detail)

Related/similar drugs

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Korlym

Korlym (mifepristone) is used to control high blood sugar levels (hyperglycemia) in adults with ...

Reviews & ratings

Add a review

View more

Warnings

You should not be treated with dinoprostone topical if you are allergic to prostaglandins, or if you have active genital herpes with a vaginal lesion, placenta previa, or if your water has broken.

Be sure your doctor knows your entire pregnancy history, especially if you have ever had a C-section or major surgery on your uterus, if you have had a baby born in a breech position, or if you have had a difficult labor or delivery of a previous child.

Before taking this medicine

You should not be treated with this medicine if you are allergic to prostaglandins, or if you have:

- active genital herpes with a vaginal lesion;
- placenta previa (the placenta is below the fetus in your uterus); or
- if your water has broken.

Be sure your doctor knows your entire pregnancy history, especially:

- if you have ever had a C-section or major surgery on your uterus;
- if you have ever had a baby born in a breech position (not head-first); or
- if you have ever had a difficult labor or delivery of a previous child.

Tell your doctor if you have ever had:

- · liver or kidney disease;
- glaucoma;
- · asthma;
- if you are 30 years or older; or
- if your pregnancy is at full term (40 weeks).

Dinoprostone is not expected to be harmful to the unborn baby when used to induce labor.

You should not breastfeed while you are being treated with dinoprostone topical.

How is dinoprostone topical given?

Dinoprostone is a **gel** or **suppository** that is placed directly onto the cervix through the vagina using a special applicator. A healthcare provider will insert dinoprostone topical at the correct time best prepare your cervix.

Dinoprostone is usually given while you are lying on your back. Your doctor may use a vaginal speculum to view your cervix. This will help your doctor determine how much of this medication to use.

You will need to remain lying down for 15 to 30 minutes, and possibly for up to 2 hours, unless your doctor tells you otherwise.

Your contractions and your baby's heartbeat will be constantly monitored after you are treated with dinoprostone. Your doctor will also check your cervix frequently to determine how much it has dilated.

Dinoprostone topical can cause contractions to occur more often and they may also last longer.

If your uterus responds to dinoprostone, you may begin having regular uterine contractions within a few hours. You may also be given other medications to help stimulate your uterine contractions and make them more regular.

If your uterus does not respond to dinoprostone within 6 hours, your doctor may apply a second dose.

A dinoprostone topical suppository is usually removed once you begin labor, or 12 hours after the medicine was

Detailed Dinoprostone topical dosage information

What happens if I miss a dose?

Since dinoprostone is used when needed in a clinical setting, it does not have a daily dosing schedule.

What happens if I overdose?

Since dinoprostone topical is given by a healthcare professional in a medical setting, an overdose is unlikely to occur.

What should I avoid after receiving dinoprostone topical?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

What other drugs will affect dinoprostone topical?

Other drugs may affect dinoprostone, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

Dinoprostone topical drug interactions (more detail)

Does dinoprostone topical interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

dinoprostone topical
+
Enter a drug name
Add

More about dinoprostone topical

- · Check interactions
- · Compare alternatives
- Reviews (18)
- · Side effects
- Dosage information
- During pregnancy
- Drug class: uterotonic agents
- Breastfeeding

• En español

Patient resources

- · Dinoprostone advanced reading
- Dinoprostone Cervical Gel
- Dinoprostone Vaginal Suppositories

Other brands

Cervidil, Prostin E2, Prepidil

Professional resources

- Dinoprostone monograph
- Dinoprostone (FDA)

Other brands

Cervidil, Prostin E2, Prepidil

Related treatment guides

- Abortion
- Labor Induction
- Trophoblastic Disease

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 6.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

্প Risk data available

CSA Schedule*

N/A Not a controlled drug

V

User Reviews & Ratings

4.3 / 10

18 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





