

[Home](#)

2. Robaxin-750

Robaxin-750

Generic name: [methocarbamol](#) (oral/injection) [*meth-oh-KAR-ba-mal*]**Brand names:** [Robaxin](#), Robaxin-750**Drug class:** [Skeletal muscle relaxants](#)[Medically reviewed](#) by Drugs.com on Aug 4, 2023. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

The Robaxin-750 brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be [generic equivalents available](#).

See also: [Robaxin](#)

What is Robaxin-750?

[Robaxin](#)-750 is a muscle relaxer that is used together with rest and physical therapy to treat skeletal muscle conditions such as pain or injury.

Robaxin-750 **injection** is sometimes used in the treatment of [tetanus](#), (lockjaw) which causes painful tightening of the muscles.

Robaxin-750 may also be used for purposes not listed in this medication guide.

Robaxin-750 side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Robaxin-750 may cause serious side effects. Stop using Robaxin-750 and call your doctor at once if you have:

- a light-headed feeling, like you might pass out;
- slow heartbeats;
- a seizure;
- [jaundice](#) (yellowing of your skin or eyes); or
- pain, bruising, swelling, or skin changes where a Robaxin-750 injection was given.

Common side effects of Robaxin-750 may include:

- [headache](#), dizziness, [drowsiness](#);
- [fever](#);
- confusion, problems with memory;
- [nausea](#), [vomiting](#), [upset stomach](#);
- blurred vision, double vision;
- flushing (warmth, redness, or tingly feeling);
- sleep problems ([insomnia](#)); or
- lack of coordination.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Robaxin-750 side effects](#) (more detail)

Related/similar drugs

Flexeril

Flexeril (cyclobenzaprine) is a muscle relaxant used to treat skeletal muscle conditions such as ...

Reviews & ratings

6.1 / 10

[237 Reviews](#)

[View more](#)

Robaxin

Robaxin (methocarbamol) is used to treat skeletal muscle conditions such as pain or injury ...

Reviews & ratings

6.3 / 10

[80 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Valium

Valium is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn ...

Reviews & ratings

8.6 / 10

[341 Reviews](#)

[View more](#)

Zanaflex

Zanaflex (tizanidine) is a short-acting muscle relaxer used to treat spasticity by temporarily ...

Reviews & ratings

7.6 / 10

[116 Reviews](#)

[View more](#)

Soma

Soma (carisoprodol) is used treat injuries and other painful muscle conditions. Includes Soma side ...

Reviews & ratings

9.1 / 10

[200 Reviews](#)

[View more](#)

Valtoco

Valtoco nasal spray is used for the treatment of epilepsy cluster seizures. Includes Valtoco side ...

Reviews & ratings

[Add a review](#)

[View more](#)

Warnings

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use Robaxin-750 if you are allergic to it.

Tell your doctor if you have ever had:

- kidney disease;
- a seizure; or
- [myasthenia gravis](#).

Robaxin-750 may harm an unborn baby. Use effective [birth control](#) to prevent pregnancy, and tell your doctor if you become pregnant.

It may not be safe to breastfeed while using Robaxin-750. Ask your doctor about any risk.

Robaxin-750 is not approved for use by anyone younger than 16 years old unless to treat tetanus.

How should I use Robaxin-750?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed. Robaxin-750 is only part of a complete treatment program that may also include rest, physical therapy, or other pain relief measures.

Robaxin-750 **oral** is taken by mouth. You may need to reduce your dose after the first 2 or 3 days of treatment.

Carefully follow your doctor's dosing instructions.

Robaxin-750 **injection** is injected into a muscle or given as an infusion into a vein. A healthcare provider will give you this injection.

Robaxin-750 **injection** is usually given as a single dose before you start taking the oral form.

Tell your caregivers if you feel any burning, pain, or swelling around the IV needle when Robaxin-750 is injected.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using Robaxin-750.

Store at room temperature away from moisture and heat. Keep the bottle tightly closed when not in use.

 [Detailed Robaxin-750 dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include nausea, extreme drowsiness, [fainting](#), seizure, or coma.

What should I avoid while taking Robaxin-750?

Avoid driving or hazardous activity until you know how Robaxin-750 will affect you. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

Drinking alcohol with this medicine can cause side effects.

What other drugs will affect Robaxin-750?

Using Robaxin-750 with other drugs that make you sleepy or slow your breathing can cause dangerous side effects or death. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for [anxiety](#) or [seizures](#).

Other drugs may affect Robaxin-750, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Tell your doctor about all your current medicines and any medicine you start or stop using.

 [Robaxin-750 drug interactions](#) (more detail)

Does Robaxin-750 interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Robaxin-750

+

Enter a drug name

Add

Popular FAQ

How long does it take for methocarbamol to work?



Is methocarbamol addictive?



Is methocarbamol a controlled substance?



Can you take methocarbamol with diclofenac?



Does methocarbamol cause weight gain?



Does methocarbamol make you sleepy?



What is Ardosons called in the U.S?



More FAQ

- [What is Miderizone used for?](#)
- [What is the maximum dose of methocarbamol?](#)

[View more FAQ...](#)

More about Robaxin-750 (methocarbamol)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(41\)](#)
- [Drug images](#)
- [Latest FDA alerts \(7\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: skeletal muscle relaxants](#)

Patient resources

- [Robaxin-750 advanced reading](#)

Professional resources

- [Methocarbamol monograph](#)
- [Methocarbamol Injection \(FDA\)](#)

Other brands

[Tanlor](#)

Other formulations

- [Robaxin](#)

Related treatment guides

- [Muscle Spasm](#)
- [Tetanus](#)

Further information


Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.


Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)


Copyright 1996-2025 Cerner Multum, Inc. Version: 10.01.

DRUG STATUS

Availability
 Discontinued

Pregnancy & Lactation
 Risk data available

CSA Schedule*
N/A Not a controlled drug

Approval History
 FDA approved 1957



User Reviews & Ratings

6.2 / 10

[41 Reviews](#)

Images

[Robaxin-750 750 MG \(ROBAXIN 750\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.