

Home

2. Naratriptan

Naratriptan

Generic name: naratriptan [NAR-a-TRIP-tan]

Brand name: Amerge

Dosage form: oral tablet (1 mg; 2.5 mg) **Drug class:** Antimigraine agents



<u>Medically reviewed</u> by Drugs.com on Aug 11, 2023. Written by <u>Cerner Multum</u>.

Uses Side effects Warnings Before taking Dosage Interactions

What is naratriptan?

Naratriptan is a headache medicine that narrows blood vessels around the brain. Naratriptan also reduces substances in the body that can trigger headache pain, sensitivity to light and sound, and other migraine symptoms.

Naratriptan is used to treat migraine headaches. Naratriptan will only treat a headache that has already begun. It will not prevent headaches or reduce the number of attacks.

Naratriptan should not be used to treat a common tension headache, a headache that causes loss of movement on one side of your body, or any headache that seems to be different from your usual migraine headaches. **Use naratriptan only if your condition has been confirmed by a doctor as migraine headaches.**

Naratriptan may also be used for purposes not listed in this medication guide.

Naratriptan side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Naratriptan may cause serious side effects. Stop using naratriptan and call your doctor at once if you have:

- · fast or pounding heartbeats;
- numbness or tingling and a pale or blue-colored appearance in your fingers or toes;
- pain or heavy feeling in your legs, hip pain, burning pain in your feet;
- sudden and severe stomach pain, bloody diarrhea, constipation, fever, weight loss;
- dangerously high blood pressure--severe headache, blurred vision, pounding in your neck or ears, nosebleed, anxiety, confusion, severe chest pain, shortness of breath, irregular heartbeats, seizure;
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;

- high levels of serotonin in the body--agitation, hallucinations, fever, fast heart rate, overactive reflexes, nausea, vomiting, diarrhea, loss of coordination, fainting; or
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance.

Common side effects of naratriptan may include:

- · dizziness, drowsiness;
- · feeling weak or tired;
- · numbness or tingling;
- flushing (warmth, redness, or tingly feeling);
- · nausea; or
- pain or tightness in your jaw, neck, or throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Naratriptan side effects (more detail)

Related/similar drugs

Celebrex

Celebrex is a nonsteroidal antiinflammatory drug used to treat pain or inflammation. Learn about ...

Reviews & ratings

7.0 / 10

226 Reviews

View more

Botox

Botox is used to treat chronic migraines, excessive sweating, bladder conditions, eye muscle ...

Reviews & ratings

5.7 / 10

473 Reviews

View more

FEATURED

Nurtec ODT

Murtae ODT (rimenanant) is used to

treat acute migraines and prevent episodic migraines, by ...

Reviews & ratings

6.3 / 10

320 Reviews

View more

Qulipta

Qulipta is used to help prevent episodic or chronic migraine headaches in adults. Qulipta is an ...

Reviews & ratings

8.1 / 10

174 Reviews

View more

Metoclopramide

Metoclopramide is used short-term to treat heartburn caused by gastroesophageal reflux. Learn about

Reviews & ratings

3.9 / 10

517 Reviews

View more

Celecoxib

Celecoxib is a nonsteroidal antiinflammatory drug (NSAID) used to treat pain or inflammation ...

Reviews & ratings

7.0 / 10

280 Reviews

View more

Diclofenac

Diclofenac is a nonsteroidal antiinflammatory drug used to treat pain and inflammation associated ...

Reviews & ratings

7.3 / 10

695 Reviews

View more

Propranolol

Propranolol is a beta-blocker that is used to treat tremors, chest pain, high blood pressure, heart ...

Reviews & ratings

8.2 / 10

1,376 Reviews

View more

Warnings

You should not use naratriptan if you have uncontrolled high blood pressure, heart problems, certain heart rhythm disorders, severe liver or kidney disease, a history of heart attack or stroke, or circulation problems that cause a lack of blood supply within the body.

Do not take naratriptan within 24 hours before or after using another migraine headache medicine.

Before taking this medicine

You should not use this medicine if you are allergic to naratriptan, or if you have:

- severe or uncontrolled high blood pressure;
- past or present heart problems;
- history of coronary artery disease, heart attack, or stroke, including "mini-stroke";
- Wolff-Parkinson-White syndrome or other heart rhythm disorder;
- a blood vessel disorder or circulation problems that cause a lack of blood supply within the body;
- · severe liver or kidney disease; or
- a headache that seems different from your usual migraine headaches.

To make sure naratriptan is safe for you, tell your doctor if you have:

- liver or kidney disease;
- high blood pressure, a heart rhythm disorder; or
- coronary heart disease (or risk factors such as diabetes, menopause, smoking, being overweight, having high

cholesterol, having a family history of coronary artery disease, being older than 40 and a man, or being a woman who has had a hysterectomy).

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether naratriptan passes into breast milk or if it could harm a nursing baby. You should not breast-feed while using this medicine.

Do not give this medicine to anyone younger than 18.

How should I use naratriptan?

Your doctor may want to give your first dose of naratriptan in a hospital or clinic setting to quickly treat any serious side effects that occur.

Follow all directions on your prescription label. **Never use more than your recommended dose.** Overuse of migraine headache medicine can make headaches worse. Tell your doctor if the medicine seems to stop working as well in treating your migraine attacks.

Take naratriptan as soon as you notice migraine symptoms.

Take 1 tablet whole with a full glass of water.

After taking a tablet: If your headache does not go away completely, or if it goes away and comes back, call your doctor before taking a second tablet.

Do not take the second tablet until at least 4 hours have passed since you took the first tablet. **Do not take more than 5** milligrams (mg) of naratriptan in 24 hours.

If your symptoms have not improved after taking 2 tablets in 24 hours, contact your doctor before taking any more tablets.

Tell your doctor if you have more than four headaches in one month (30 days).

Naratriptan can raise blood pressure to dangerous levels. Your blood pressure may need to be checked often while you are using this medicine. If you use naratriptan long-term, your heart function may need to be checked using an electrocardiograph or ECG (sometimes called an EKG).

Store at room temperature away from moisture, heat, and light.

Naratriptan dosing information

Usual Adult Dose for Migraine:

Use only after a clear diagnosis of migraine has been established

Initial dose: 1 mg or 2.5 mg orally, once

-Provided there has been some response to first dose, a second dose may be administered at least 4 hours later if migraine returns or symptoms recur.

Maximum dose: 5 mg in a 24-hour period

Comments:

- -This drug should not be used to treat basilar or hemiplegic migraines because these patients are at a greater risk of stroke.
- -The safety of treating an average of 4 or more migraine attacks in a 30-day period has not been established.

Use: For the acute treatment of migraine with or without aura.

Detailed Naratriptan dosage information

What happens if I miss a dose?

Since naratriptan is used as needed, it does not have a daily dosing schedule. Call your doctor if your symptoms do not improve after using naratriptan.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include tiredness, neck stiffness, loss of coordination, or feeling light-headed.

What should I avoid while using naratriptan?

Do not take naratriptan within 24 hours before or after using another migraine headache medicine, including:

- medicines like naratriptan--almotriptan, eletriptan, frovatriptan, rizatriptan, sumatriptan, zolmitriptan, and others; or
- ergot medicine--dihydroergotamine, ergotamine, ergonovine, methylergonovine.

naratriptan may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

What other drugs will affect naratriptan?

Taking naratriptan while you are using certain other medicines can cause **high levels of serotonin to build up in your body, a condition called "serotonin syndrome,"** which can be fatal. Tell your doctor if you also use:

- medicine to treat depression;
- medicine to treat a psychiatric disorder;
- a narcotic (opioid) medication; or
- medicine to prevent nausea and vomiting.

This list is not complete. Other drugs may interact with naratriptan, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

Naratriptan drug interactions (more detail)

Does naratriptan interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

naratriptan	
+	
Enter a drug name	Add

More about naratriptan

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (51)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: antimigraine agents
- Breastfeeding
- En español

Patient resources

Naratriptan advanced reading

Other brands

Amerge

Professional resources

- Naratriptan monograph
- Naratriptan (FDA)

Other brands

Amerge

Related treatment guides

• Migraine

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 12.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

്യ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

8.5 / 10

51 Reviews

Related News

Combination Therapy Recommended For Migraines

Images

Naratriptan Hydrochloride 1 mg (I53)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×