

Home

2. Luxiq

Luxiq

Generic name: betamethasone topical [BAY-ta-METH-a-sone-TOP-ik-al]

Brand names: Diprolene, Luxiq, Sernivo

Drug class: Topical steroids



Medically reviewed by Drugs.com on Dec 17, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is Luxiq?

Betamethasone is a highly potent steroid that prevents the release of substances in the body that cause inflammation.

Luxiq (for the skin) is used to treat the inflammation and itching caused by a number of skin conditions such as eczema or psoriasis.

Luxig may also be used for purposes not listed in this medication guide.

Luxiq side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Betamethasone may cause serious side effects. Stop using betamethasone and call your doctor at once if you have:

- severe skin irritation where the medicine was applied; or
- signs of skin infection (swelling, redness, warmth, oozing).

Your skin can absorb topical steroid medicine, which may cause steroid side effects throughout the body. Stop using betamethasone and call your doctor if you have:

- blurred vision, tunnel vision, eye pain, or seeing halos around lights;
- slow wound healing, thinning skin, increased body hair;
- increased thirst or urination, dry mouth, fruity breath odor;
- · weight gain, puffiness in your face; or
- muscle weakness, tired feeling, depression, anxiety, feeling irritable.

Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Luxiq.

Common side effects of Luxiq may include:

- · itching, redness, burning, stinging, or blistering of treated skin;
- skin bruising or shiny appearance; or
- folliculitis (redness or crusting around your hair follicles).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



Related/similar drugs

Skyrizi

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis, ulcerative colitis ...

Reviews & ratings

7.9 / 10

56 Reviews

View more

Dupixent

Dupixent may be used to treat eczema, eosinophilic or oral-corticosteroid-dependent asthma, chronic ...

Reviews & ratings

7.2 / 10

499 Reviews

View more

FEATURED

Otezla

Otezla (apremilast) is used to treat plaque psoriasis, psoriatic arthritis, and oral ulcers ...

Reviews & ratings

5.2 / 10

439 Reviews

View more

Stelara

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

Reviews & ratings

7.6 / 10

252 Reviews

View more

Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

Reviews & ratings

7.0 / 10

223 Reviews

View more

Humira

Humira is a tumor necrosis factor blocker that is used to treat many inflammatory conditions such ...

Reviews & ratings

6.4 / 10

698 Reviews

View more

Enbrel

Enbrel is used to treat rheumatoid arthritis, ankylosing spondylitis, and psoriasis. Learn about ...

Reviews & ratings

7.7 / 10

306 Reviews

View more

Adalimumab

Adalimumah is usad to treat planua

psoriasis, Crohn's disease, ulcerative colitis, rheumatoid ...

Reviews & ratings

6.4 / 10

737 Reviews

View more

Clobetasol topical

Clobetasol is available as a cream, gel, ointment, lotion, solution, foam or spray and is used to ...

Reviews & ratings

8.2 / 10

164 Reviews

View more

Warnings

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use Luxiq if you are allergic to it.

Tell your doctor if you have ever had:

- skin irritation after using steroid medicine;
- thinning of the skin where you will apply Luxiq;
- cataracts or glaucoma;
- · diabetes;
- liver disease;
- · adrenal gland problems; or
- any type of skin infection.

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant.

It may not be safe to breast-feed a baby while you are using this medicine. Ask your doctor about any risks. If you apply betamethasone to your chest, avoid areas that may come into contact with the baby's mouth.

Do not use Luxiq on a child without a doctor's advice. Children can absorb larger amounts of this medicine through

the skin and may be more likely to have side effects.

Diprolene is not approved for use by anyone younger than 13 years old. **Sernivo** and **Luxiq** are not approved for anyone younger than 18 years old.

How should I use Luxiq?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Do not take by mouth. Topical medicine is for use only on the skin. Do not use on open wounds or on sunburned, windburned, dry, or irritated skin. Rinse with water if this medicine gets in your eyes or mouth.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions.

Wash your hands before and after using this medicine, unless you are using the medicine to treat the skin on your hands.

Apply a small amount to the affected area and rub it gently into the skin. Do not apply Luxiq over a large area of skin.

Do not cover the treated skin area unless your doctor tells you to. Covering treated areas can increase the amount of medicine absorbed through your skin and may cause harmful effects.

Use Luxiq regularly to get the most benefit. Stop using the medicine once your symptoms clear up.

Call your doctor if your skin condition does not improve after 2 weeks, or if it gets worse.

Do not use **Diprolene** for longer than 2 weeks in a row. Do not use **Sernivo** or **Luxiq** for longer than 4 weeks.

Store at room temperature away from moisture and heat.

The **foam** is flammable. Do not use near high heat or open flame. Do not smoke until the foam has completely dried on your skin.

Detailed Luxiq dosage information

What happens if I miss a dose?

Apply the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** apply two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 if anyone has accidentally swallowed the medication.

Long term use of high doses can lead to thinning skin, easy bruising, changes in body fat (especially in your face, neck, back, and waist), increased acne or facial hair, menstrual problems, impotence, or loss of interest in sex.

What should I avoid while using Luxiq?

Do not get Luxiq in your eyes. If contact does occur, rinse with water.

Avoid applying Luxiq to the skin of your face, underarms, or groin area without your doctor's instruction.

Do not use Luxiq to treat any condition that has not been checked by your doctor.

What other drugs will affect Luxiq?

Medicine used on the skin is not likely to be affected by other drugs you use. But many drugs can interact with each other. Tell each of your health care providers about all medicines you use, including prescription and over-the-counter medicines, vitamins, and herbal products.

1 Luxiq drug interactions (more detail)

Does Luxiq interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

| Does Luxiq Foam help with hair loss? | ~ |
|--|---|
| Can you buy Luxiq foam over-the-counter (OTC)? | ~ |
| What is Luxiq Foam used to treat? | ~ |
| How do you use Luxiq Foam? | ~ |

More about Luxiq (betamethasone topical)

- Check interactions
- · Compare alternatives
- Reviews (4)
- Side effects
- · Dosage information
- During pregnancy

- · Drug class: topical steroids
- Breastfeeding
- En español

Patient resources

Luxiq advanced reading

Other brands

Diprolene, Valisone, Diprosone, Sernivo, ... +4 more

Professional resources

- Betamethasone, Betamethasone Benzoate, Betamethasone Dipropionate, Betamethasone Valerate (Topical) monograph
- Betamethasone Valerate Lotion (FDA)

Other brands

Diprolene, Sernivo, Diprolene AF

Related treatment guides

- Lichen Sclerosus
- Atopic Dermatitis
- Dermatitis
- Dermatological Disorders

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 10.02.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ড Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

WADA Class

Anti-Doping Classification

User Reviews & Ratings

4 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







