

[Home](#)2. [Dexedrine](#)

Dexedrine Spansule

Generic name: [dextroamphetamine](#) [*DEX-tro-am-FET-a-meen*]**Brand names:** Dexedrine Spansule, [ProCentra](#), [Xelstrym](#), [Zenzedi](#)**Drug class:** [CNS stimulants](#)[Medically reviewed](#) by Drugs.com on Jun 22, 2023. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

What is Dexedrine Spansule?

Dexedrine Spansule **tablets** and **oral solution (liquid)** are used to treat attention deficit hyperactivity disorder (ADHD) in children who are 3 to 16 years old.

Dexedrine Spansule **extended-release capsules** are used to treat attention deficit hyperactivity disorder (ADHD) in children who are 6 to 16 years old.

Dexedrine Spansule **skin patch** is used to treat attention deficit hyperactivity disorder (ADHD) in adults and children at least 6 years old.

Dexedrine Spansule **tablets**, **oral solution (liquid)** and **extended-release capsules** are also used in the treatment of a sleep disorder called narcolepsy.

Dexedrine Spansule may also be used for purposes not listed in this medication guide.

Dexedrine Spansule side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) or **a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Stimulants have caused stroke, heart attack, and sudden death in certain people. Get emergency medical help if you have **signs of a heart attack or stroke**: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, or feeling short of breath.

Dexedrine Spansule may cause serious side effects. Call your doctor at once if you have:

- **signs of heart or blood circulation problems** --chest pain, trouble breathing, numbness, cold feeling, or pale appearance of your fingers or toes;
- **signs of mania or psychosis**-- seeing or hearing things that are not real, aggression, paranoia, decreased need for sleep, risk-taking behavior, being agitated or talkative;

- a seizure, muscle twitches (tics), changes in your vision, blurry vision; or
- mild skin irritation where the patch is worn.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, [fast heart rate](#), muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Dexedrine Spansule can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Dexedrine Spansule.

Common side effects of Dexedrine Spansule may include:

- dry mouth, upset stomach, loss of appetite, weight loss;
- headache, dizziness, sleep problems (insomnia), irritability, [muscle twitching](#) (tics); or
- nausea, vomiting, increased blood pressure, tremors, fast heartbeats.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Dexedrine side effects](#) (more detail)

Related/similar drugs

Adderall

Adderall (amphetamine and dextroamphetamine) is used to treat narcolepsy and ADHD. Includes side ...

Reviews & ratings

7.2 / 10

[505 Reviews](#)

[View more](#)

Vyvanse

Vyvanse is a prescription stimulant used to treat ADHD in children and adults, and binge eating ...

Reviews & ratings

7.4 / 10

[973 Reviews](#)

[View more](#)

FEATURED

Xywav

Xywav is used for the treatment of cataplexy and excessive daytime sleepiness (EDS) in patients ...

Reviews & ratings

8.1 / 10

[16 Reviews](#)

[View more](#)

Strattera

Strattera (atomoxetine) is used to treat attention deficit hyperactivity disorder.

Learn about side ...

Reviews & ratings

5.4 / 10

[541 Reviews](#)

[View more](#)

Concerta

Concerta is used to treat attention deficit disorder (ADD) and attention deficit hyperactivity ...

Reviews & ratings

6.9 / 10

[424 Reviews](#)

[View more](#)

Ritalin

Ritalin (methylphenidate) is used to treat attention deficit disorder (ADD) and narcolepsy ...

Reviews & ratings

7.0 / 10

[221 Reviews](#)

[View more](#)

Focalin

Focalin (dexamethylphenidate) is used to treat attention deficit hyperactivity disorder (ADHD) ...

Reviews & ratings

5.6 / 10

[139 Reviews](#)

[View more](#)

Provigil

Provigil promotes wakefulness and is used to treat excessive daytime sleepiness caused by ...

Reviews & ratings

7.4 / 10

[169 Reviews](#)

[View more](#)

Armodafinil

Armodafinil may be used to improve wakefulness in adults who are very sleepy due to narcolepsy ...

Reviews & ratings

7.3 / 10

[435 Reviews](#)

[View more](#)

Warnings

You should not use this medicine if you have glaucoma, overactive thyroid, severe agitation, moderate to severe [high blood pressure](#), heart disease or [coronary artery disease](#), or a history of drug or alcohol addiction.

Dexedrine Spansule may be habit-forming. **Misuse can cause addiction, serious heart problems, or death.**

Call your doctor right away if you have: chest pain, seeing or hearing things that are not real, decreased need for sleep, being agitated, unexplained wounds on your fingers or toes.

Before taking this medicine

You should not use Dexedrine Spansule if you are allergic to any stimulant medicine, or if you have:

- moderate to severe high blood pressure, heart problems or coronary artery disease (clogged arteries);
- overactive thyroid, glaucoma;
- severe anxiety, tension, or agitation (stimulant medicine can make these symptoms worse); or

- a history of drug or alcohol addiction.

Do not use Dexedrine Spansule if you have used an MAO inhibitor in the past 14 days. A dangerous [drug interaction](#) could occur. MAO inhibitors include [isocarboxazid](#), linezolid, [methylene blue](#) injection, phenelzine, [tranylcypromine](#), and others.

Tell your doctor if you also use opioid medicine, [herbal products](#), or medicine for depression, mental illness, [Parkinson's disease](#), migraine headaches, serious infections, or prevention of nausea and vomiting. **An interaction with Dexedrine Spansule could cause a serious condition called serotonin syndrome.**

Tell your doctor if you or a family member has ever had:

- a heart defect, high blood pressure, a family history of heart disease or sudden death;
- depression, mental illness, [bipolar disorder](#), psychosis, or suicidal thoughts or actions;
- motor tics (muscle twitches) or [Tourette's syndrome](#);
- seizures or epilepsy, an abnormal brain wave test (EEG);
- [contact dermatitis](#);
- blood circulation problems or [peripheral vascular disease](#) such as [Raynaud's syndrome](#); or
- kidney disease.

Tell your doctor if you are pregnant or plan to become pregnant. Using Dexedrine Spansule during pregnancy may increase the risk of premature birth, withdrawal symptoms or low birth weight.

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of [dextroamphetamine skin patch](#) on the baby.

You should not breastfeed while using Dexedrine Spansule.

How should I use Dexedrine Spansule?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Dexedrine Spansule may be habit-forming. **Misuse can cause addiction, overdose, serious heart problems or death.** Keep the medication where others cannot get to it. Selling or giving away this medicine is against the law.

Dexedrine Spansule is sometimes taken 2 or 3 times per day. **Extended-release** this medicine is taken only once per day.

Apply the patch to clean, dry, hairless skin. Press firmly into place for about 10 seconds. Leave the patch on while bathing, showering, or swimming. **Apply the patch 2 hours before the effect is needed and remove the patch within 9 hours after application.**

Choose a different place on your body each time you apply a new patch. Do not apply to the same skin twice in 7 days. **Do not wear more than one patch at a time. Use only 1 patch per day.** Never cut a skin patch.

Swallow the Dexedrine Spansule **extended release capsule** whole and do not crush, chew, break, or open it.

Measure **liquid medicine** with the supplied measuring device (not a kitchen spoon).

Your doctor will need to check your progress on a regular basis. You may need medical tests. **Tell any doctor who treats you that you are using this medicine.**

Your dose needs may change if you switch to a different brand, strength, or form of this medicine. **Avoid medication errors by using only the medicine your doctor prescribes.**

Store at room temperature away from moisture, heat, and light.

Read and carefully follow the instructions provided with this medicine about how to safely dispose of any unused portion. **Do not** flush Dexedrine Spansule **skin patch** down the toilet.

 [Detailed Dexedrine dosage information](#)

What happens if I miss a dose?

Skip the missed dose and use your next dose at the regular time. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of Dexedrine Spansule can be fatal.**

Overdose symptoms may include restlessness, rapid breathing, hallucinations, aggressiveness, vomiting, diarrhea, uneven heartbeats, fainting, seizures, or coma.

What should I avoid while using dextroamphetamine?

Dexedrine Spansule may cause blurred vision and may impair your reactions. Avoid driving or hazardous activity until you know how this medicine will affect you.

Avoid external heat sources like hair dryers, heat lamps or saunas while wearing Dexedrine Spansule **skin patch**.

Avoid drinking fruit juices or taking vitamin C while you take Dexedrine Spansule. These can make your body absorb less of the medicine.

What other drugs will affect Dexedrine Spansule?

Sometimes it is not safe to use certain medicines at the same time. Some drugs can affect your blood levels of other drugs you use, which may increase side effects or make the medicines less effective.

Tell your doctor about all your current medicines. Many drugs can affect Dexedrine Spansule, especially:

- an antacid, [migraine headache](#) medicine, seizure medicine;
- fentanyl, tramadol, lithium, buspirone;
- [acetazolamide](#), tryptophan, methenamine, vitamin c products;
- an antidepressant, a diuretic or "water pill"; or
- blood pressure medication.

This list is not complete and many other drugs may affect Dexedrine Spansule. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible [drug interactions](#) are listed here.

 [Dexedrine drug interactions](#) (more detail)

Does Dexedrine interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Dexedrine

+

Enter a drug name

Add

Popular FAQ

'Black Beauties' capsules in the 1970's - What drug was that?



More FAQ

- [Concerta vs Adderall - What's the difference between them?](#)
- [What pill has a half moon and 26 On same side. orange, and round?](#)

[View more FAQ...](#)

More about Dexedrine (dextroamphetamine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(123\)](#)
- [Drug images](#)
- [Latest FDA alerts \(4\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [Drug class: CNS stimulants](#)
- [Breastfeeding](#)

- [En español](#)

Patient resources

- [Dexedrine advanced reading](#)
- [Dexedrine Spansules \(Advanced Reading\)](#)
- [Dexedrine](#)

Other brands

[ProCentra](#), [Xelstrym](#), [Zenzedi](#), [Dextrostat](#), [Liquadd](#)

Professional resources

- [Dexedrine prescribing information](#)
- [Dextroamphetamine \(AHFS Monograph\)](#)

Other brands

[ProCentra](#), [Xelstrym](#), [Zenzedi](#), [Dextrostat](#)

Related treatment guides

- [ADHD](#)
- [Narcolepsy](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 9.02.

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
2	High potential for abuse
Approval History	
	Drug history at FDA

User Reviews & Ratings

8.6 / 10

[123 Reviews](#)

Images

[Dexedrine 5 mg \(SKF E19\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices

ADVERTISEMENT



