

Home

2. Prolensa

Prolensa (ophthalmic)

Generic name: bromfenac (ophthalmic) [BROM-fen-ak-off-THAL-mik]

Brand names: BromSite, Prolensa

Drug class: Ophthalmic anti-inflammatory agents



Medically reviewed by Drugs.com on Jan 6, 2025. Written by Cerner Multum.

Uses Side effects Before taking Dosage What to avoid Interactions

What is Prolensa ophthalmic?

Prolensa is a non-steroidal anti-inflammatory drug (NSAID).

Prolensa (for the eyes) is used to treat swelling and pain caused by cataract surgery.

Prolensa may also be used for purposes not listed in this medication guide.

Prolensa ophthalmic side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Prolensa may cause serious side effects. Stop using Prolensa and call your doctor at once if you have:

- any change in your vision;
- slow healing after your eye surgery;
- eye redness or bleeding;
- eye pain or swelling; or
- severe eye discomfort, increased tears, crusting or drainage.

Common side effects of Prolensa may include:

- increased sensitivity of your eyes to light;
- mild eye irritation, or feeling like something is in your eye;
- · headache; or
- seeing flashes of light or "floaters" in your vision.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.

You may report side effects to FDA at 1-800-FDA-1088.

• Prolensa side effects (more detail)

Related/similar drugs

Lotemax

Lotemax and Lotemax SM (loteprednol etabonate) are corticosteroid medications used to help treat ...

Reviews & ratings

6.4 / 10

16 Reviews

View more

Ozurdex

Ozurdex is an implant injected into the eye to treat swelling that may occur when there is a ...

Reviews & ratings

6.1 / 10

8 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Durezol

Durezol (difluprednate) is used to treat pain and swelling following eye surgery. Includes Durezol ...

Reviews & ratings

6.0 / 10

22 Reviews

View more

ophthalmic

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use Prolensa if you are allergic to it.

Tell your doctor if you have ever had:

- · asthma or sulfite allergy;
- diabetes;
- rheumatoid arthritis;
- dry eye syndrome;
- an allergy to aspirin or other NSAIDs;
- · a bleeding or blood-clotting disorder; or
- more than one eye surgery in a short period of time.

If you are pregnant, you should not use Prolensa unless your doctor tells you to. Using an NSAID during the last 20 weeks of pregnancy can cause serious heart or kidney problems in the unborn baby and possible complications with your pregnancy.

Tell your doctor if you are breastfeeding.

How should I use Prolensa ophthalmic?

Follow all directions on your prescription label. **Do not use this medication for more than the number of days your doctor has recommended.** Using an NSAID eye drop too long can increase your risk of serious side effects that may damage your vision.

Do not use Prolensa while wearing contact lenses. This medicine may contain a preservative that can discolor soft contact lenses. Wait at least 10 minutes after using Prolensa before putting in your contact lenses.

The **Prolensa** brand of bromfenac ophthalmic is usually given as 1 drop in each affected eye **once daily** for 14 days.

The Bromsite brand of bromfenac ophthalmic is usually given as 1 drop in each affected eye twice daily for 14 days.

Use only the number of drops your doctor has prescribed.

Wash your hands before using eye medication.

To apply the eye drops: Tilt your head back slightly and pull down your lower eyelid to create a small pocket. Hold the dropper above the eye and squeeze a drop into this pocket. Close your eyes for 1 or 2 minutes.

Do not touch the tip of the eye dropper or place it directly on your eye. A contaminated dropper can infect your eye, which could lead to serious vision problems.

Do not use the eye drops if the liquid has changed colors or has particles in it. Call your pharmacist for new medicine.

Store at room temperature away from moisture and heat. Do not freeze. Keep the bottle tightly closed when not in use.

Detailed Prolensa dosage information

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

Call your doctor for instructions if you miss your first dose, which should be given 1 day before surgery.

What happens if I overdose?

An overdose of Prolensa ophthalmic is not expected to be dangerous. Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 if anyone has accidentally swallowed the medication.

What should I avoid while using bromfenac ophthalmic?

Prolensa may cause blurred vision and may impair your reactions. Avoid driving or hazardous activity until you know how this medicine will affect you.

Do not use other eye medications unless your doctor tells you to.

What other drugs will affect Prolensa ophthalmic?

Medicine used in the eyes is not likely to be affected by other drugs you use. But many drugs can interact with each other. Tell each of your healthcare providers about all medicines you use, including prescription and over-the-counter medicines, vitamins, and herbal products.

1 Prolensa drug interactions (more detail)

Does Prolensa interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Prolensa	
+	
Enter a drug name	Add

More about Prolensa (bromfenac ophthalmic)

· Check interactions

- · Compare alternatives
- Pricing & coupons
- Reviews (8)
- Side effects
- Dosage information
- During pregnancy
- · Generic availability
- FDA approval history
- Drug class: ophthalmic anti-inflammatory agents
- Breastfeeding
- En español

Patient resources

- · Prolensa advanced reading
- Prolensa

Other brands

Bromday, BromSite, Xibrom

Professional resources

- Prolensa prescribing information
- Bromfenac (AHFS Monograph)

Other brands

BromSite

Related treatment guides

• Postoperative Ocular Inflammation

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 2.04.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ড Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

The Drug history at FDA

User Reviews & Ratings

8.6 / 10

8 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









