



[Home](#)

2. Tylenol with Codeine #3

## Tylenol with Codeine #3

**Generic name:** [acetaminophen and codeine](#) [ a-SEET-a-MIN-o-fen-and-KOE-deen ]

**Brand names:** Tylenol with Codeine #3, [Tylenol with Codeine #4](#)

**Drug class:** [Narcotic analgesic combinations](#)



[Medically reviewed](#) by Drugs.com on Aug 3, 2023. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)



The Tylenol with Codeine #3 brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be [generic equivalents available](#).

## What is Tylenol with Codeine #3?

[Tylenol with Codeine](#) #3 is a combination medicine used to relieve moderate to severe pain.

Tylenol with Codeine #3 may also be used for purposes not listed in this medication guide.

## Tylenol with Codeine #3 side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

**Opioid medicine can slow or stop your breathing, and death may occur.** A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up.

**In rare cases, acetaminophen may cause a severe skin reaction that can be fatal.** This could occur even if you have taken [acetaminophen](#) in the past and had no reaction. **Stop taking Tylenol with Codeine #3 and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling.**

**Tylenol with Codeine #3 may cause serious side effects. Call your doctor at once if you have:**

- noisy breathing, sighing, shallow breathing, breathing that stops;
- a light-headed feeling, like you might pass out;
- confusion, severe drowsiness;
- **liver problems**--nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes); or

- **high levels of serotonin in the body**--agitation, hallucinations, fever, sweating, shivering, [fast heart rate](#), muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea.

**Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders.**

Common side effects include:

- drowsiness, dizziness, feeling tired;
- nausea, vomiting, stomach pain;
- [constipation](#); or
- headache.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Tylenol with Codeine #3 side effects](#) (more detail)

## Related/similar drugs

### Naproxen

Naproxen is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation caused by ...

#### Reviews & ratings

6.8 / 10

[683 Reviews](#)

[View more](#)

### Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

#### Reviews & ratings

5.9 / 10

[1,404 Reviews](#)

[View more](#)

#### FEATURED

### OmvoH

OmvoH is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

## Reviews & ratings

[1 Reviews](#)

[View more](#)

## Tramadol

Tramadol is an opioid medication that may be used to treat moderate to moderately severe chronic ...

## Reviews & ratings

7.0 / 10

[1,876 Reviews](#)

[View more](#)

## Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

## Reviews & ratings

6.0 / 10

[618 Reviews](#)

[View more](#)

## Duloxetine

Duloxetine is a selective serotonin and norepinephrine reuptake inhibitor antidepressant used to ...

## Reviews & ratings

6.2 / 10

[2,456 Reviews](#)

[View more](#)

## Meloxicam

Meloxicam is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation caused by ...

## Reviews & ratings

6.3 / 10

[600 Reviews](#)

[View more](#)

## Aspirin

Aspirin is used to treat mild to moderate pain and to reduce fever or inflammation. Learn about ...

### Reviews & ratings

7.8 / 10

[51 Reviews](#)

[View more](#)

## Ibuprofen

Ibuprofen (Advil, Motrin) treats minor aches and pains caused by the common cold, headaches ...

### Reviews & ratings

7.3 / 10

[240 Reviews](#)

[View more](#)

## Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-the-counter painkiller and fever ...

### Reviews & ratings

6.1 / 10

[13 Reviews](#)

[View more](#)

## Warnings

**MISUSE OF OPIOID MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH.** Keep the medication in a place where others cannot get to it.

**Do not** give this medicine to anyone younger than 12 years old, or anyone under 18 who recently had surgery to remove the tonsils or adenoids.

**Taking opioid medicine during pregnancy may cause life-threatening withdrawal symptoms in the newborn.**

**Fatal side effects can occur if you use opioid medicine with alcohol, or with other drugs that cause drowsiness**

or slow your breathing.

**Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling.**

## Before taking this medicine

You should not use Tylenol with Codeine #3 if you are allergic to acetaminophen or codeine, or if you have:

- severe asthma or breathing problems; or
- a blockage in your stomach or intestines.

In some people, codeine breaks down rapidly in the liver and reaches higher than normal levels in the body. This can cause dangerously slow breathing and may cause death.

Codeine is **not approved** for use by anyone younger than 12 years old.

**Do not** give this medicine to anyone younger than 18 years old who recently had surgery to remove the tonsils or adenoids.

Tell your doctor if you have ever had:

- breathing problems, [sleep apnea](#);
- [liver disease](#);
- [alcoholism](#) or [drug addiction](#);
- urination problems;
- kidney disease; or
- problems with your pancreas, thyroid, or gallbladder.

**If you use opioid medicine while you are pregnant, your baby could become dependent on the drug.** This can cause life-threatening withdrawal symptoms in the baby after it is born. Babies born dependent on opioids may need medical treatment for several weeks.

**Do not breastfeed.** Codeine can pass into breast milk and cause drowsiness, breathing problems, or death in a nursing baby.

## How should I take Tylenol with Codeine #3?

Follow the directions on your prescription label and read all medication guides. **Never use this medicine in larger amounts, or for longer than prescribed. An overdose can damage your liver or cause death.** Tell your doctor if you feel an increased urge to use more of this medicine.

**Never share opioid medicine with another person, especially someone with a history of drug abuse or addiction. MISUSE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH.** Keep the medication in a place where others cannot get to it. Selling or giving away opioid medicine is against the law.

If you need surgery or medical tests, tell the surgeon or doctor ahead of time that you are using this medicine.

**Do not stop using Tylenol with Codeine #3 suddenly after long-term use**, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using this medicine.

Store at room temperature away from moisture and heat. Keep track of your medicine. You should be aware if anyone is using it improperly or without a prescription.

Do not keep leftover opioid medication. **Just one dose can cause death in someone using this medicine accidentally or improperly.** Ask your pharmacist where to locate a drug take-back disposal program. If there is no take-back program, flush the unused medicine down the toilet.

 [Detailed Tylenol with Codeine #3 dosage information](#)

## What happens if I miss a dose?

Since Tylenol with Codeine #3 is used for pain, you are not likely to miss a dose. Skip any missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An opioid overdose can be fatal, especially in a child or other person using the medicine without a prescription.** Overdose symptoms may include severe drowsiness, pinpoint pupils, slow breathing, or no breathing.

Your doctor may recommend you get naloxone (a medicine to reverse an [opioid overdose](#)) and keep it with you at all times. A person caring for you can give the naloxone if you stop breathing or don't wake up. Your caregiver must still get emergency medical help and may need to perform CPR (cardiopulmonary resuscitation) on you while waiting for help to arrive.

Anyone can buy naloxone from a pharmacy or local health department. Make sure any person caring for you knows where you keep naloxone and how to use it.

## What should I avoid while taking Tylenol with Codeine #3?

Avoid driving or operating machinery until you know how Tylenol with Codeine #3 will affect you. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

**Do not drink alcohol.** Dangerous side effects or death could occur.

Ask a doctor or pharmacist before using any other medicine that may contain acetaminophen (sometimes abbreviated as APAP). **Taking certain medications together can lead to a fatal overdose.**

## What other drugs will affect Tylenol with Codeine #3?

You may have breathing problems or withdrawal symptoms if you start or stop taking certain other medicines. Tell your doctor if you also use an [antibiotic](#), antifungal medication, heart or blood pressure medication, seizure medication, or medicine to treat HIV or [hepatitis C](#).

**Opioid medication can interact with many other drugs and cause dangerous side effects or death.** Be sure your doctor knows if you also use:

- cold or allergy medicines, bronchodilator asthma/COPD medication, or a diuretic ("water pill");
- medicines for [motion sickness](#), [irritable bowel syndrome](#), or [overactive bladder](#);
- **other opioids**--opioid pain medicine or prescription cough medicine;
- **a sedative like Valium**--diazepam, [alprazolam](#), lorazepam, Xanax, Klonopin, Versed, and others;
- **drugs that make you sleepy or slow your breathing**--a sleeping pill, muscle relaxer, medicine to treat mood disorders or mental illness;
- **drugs that affect serotonin levels in your body**--a stimulant, or medicine for [depression](#), [Parkinson's disease](#), migraine headaches, serious infections, or nausea and vomiting.

This list is not complete. Other drugs may affect Tylenol with Codeine #3, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). **Not all possible interactions are listed here.**

**i** [Tylenol with Codeine #3 drug interactions](#) (more detail)

## Does Tylenol with Codeine #3 interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Tylenol with Codeine #3

+

Add

## More about Tylenol with Codeine #3 (acetaminophen / codeine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(159\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: narcotic analgesic combinations](#)

## Patient resources

### Other brands

[Tylenol with Codeine](#), [Capital w/Codeine](#), [Tylenol with Codeine #4](#)

## Professional resources

- [Acetaminophen and Codeine Oral Solution prescribing information](#)

Other brands

[Tylenol with Codeine](#)

Other formulations

- [Tylenol](#)

Related treatment guides

- [Osteoarthritis](#)
- [Cough](#)
- [Pain](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.


[Medical Disclaimer](#)


Copyright 1996-2025 Cerner Multum, Inc. Version: 20.01.

Tylenol Pediatric Dosage Guide

Use these dosage charts for infants and children under 12 years old:  
[Tylenol Dosage Charts \(Infants & Children\)](#)

DRUG STATUS

**Availability**  
 Discontinued

**Pregnancy & Lactation**  
 Risk data available

**CSA Schedule\***  
**M** Multiple: 3,5



User Reviews & Ratings

4.6 / 10

[159 Reviews](#)

Images



# Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



## About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

## Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

## Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT

Copyright © 2000-2025 Drugs.com. All rights reserved.

