

Home

2. Albuterol/ipratropium

Albuterol and ipratropium (inhalation)

Generic name: albuterol and ipratropium (inhalation) [al-BUE-ter-ol-and-IP-ra-TRO-pee-um]

Brand names: Combivent Respirat, Combivent, DuoNeb

Dosage forms: inhalation aerosol (100 mcg-20 mcg/inh), inhalation solution (2.5 mg-0.5 mg/3 mL)

Drug class: Bronchodilator combinations



Medically reviewed by Drugs.com on Sep 25, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is albuterol and ipratropium inhalation?

Albuterol and ipratropium are bronchodilators that relax muscles in the airways and increase air flow to the lungs.

Albuterol and ipratropium inhalation is used to prevent bronchospasm in people with chronic obstructive pulmonary disease (COPD) who are also using other medicines to control their condition.

Albuterol and ipratropium inhalation may also be used for purposes not listed in this medication guide.

Albuterol and ipratropium inhalation side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Albuterol and ipratropium may cause serious side effects. Call your doctor at once if you have:

- wheezing, choking, or other breathing problems after using albuterol and ipratropium;
- · chest pain;
- · fast or pounding heartbeats, fluttering in your chest;
- tremors, nervousness;
- · swelling of your ankles or feet;
- blurred vision, tunnel vision, eye pain, or seeing halos around lights;
- painful or difficult urination; or
- **low potassium**--leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling.

Common side effects of albuterol and ipratropium may include:

- · headache;
- trouble breathing; or
- cold symptoms such as stuffy nose, sneezing, cough, or sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Albuterol / ipratropium side effects (more detail)

Related/similar drugs

Symbicort

Symbicort (budesonide and formoterol) is used to prevent bronchospasm in people with asthma or ...

Reviews & ratings

6.6 / 10

226 Reviews

View more

Ventolin HFA

Ventolin HFA (albuterol) is used to treat or prevent breathing problems in patients who have asthma ...

Reviews & ratings

7.5 / 10

18 Reviews

View more

FEATURED

Breztri Aerosphere

Breztr

(budesonide/glycopyrrolate/formoterol fumarate) is a combination inhaler that may be used ...

Reviews & ratings

6.9 / 10

119 Reviews

View more

Dupixent

Dupixent may be used to treat eczema, eosinophilic or oral-corticosteroid-dependent asthma, chronic ...

Reviews & ratings

7.2 / 10

499 Reviews

View more

Breo Ellipta

Breo Ellipta (fluticasone and vilanterol) is used to prevent airflow obstruction or bronchospasm in ...

Reviews & ratings

5.8 / 10

228 Reviews

View more

Spiriva

Spiriva (tiotropium) is used to prevent bronchospasm in people with bronchitis, emphysema, or COPD ...

Reviews & ratings

5.9 / 10

122 Reviews

View more

Anoro Ellipta

Anoro (umeclidinium and vilanterol inhalation powder) is used to treat chronic obstructive ...

Reviews & ratings

7.1 / 10

69 Reviews

View more

Xopenex

Yonanav is a short-action

bronchodilator used to treat or prevent bronchospasm in asthma and other ...

Reviews & ratings

9.0 / 10

15 Reviews

View more

Combivent

Combivent Respimat is used to prevent bronchospasm in people with chronic obstructive pulmonary ...

Reviews & ratings

8.1 / 10

23 Reviews

View more

Warnings

Seek medical attention if your breathing problems get worse quickly, or if you think your medications are not working as well. Overuse of albuterol and ipratropium may increase the risk of death. It is critical that you use only the prescribed dose of this medicine.

Before taking this medicine

You should not use albuterol and ipratropium if you are allergic to albuterol, ipratropium, or atropine.

Tell your doctor if you have ever had:

- · heart disease, high blood pressure, coronary artery disease, or heart rhythm disorder;
- a seizure disorder such as epilepsy;
- · diabetes;
- · overactive thyroid;
- glaucoma;
- liver or kidney disease; or
- enlarged prostate, problems with urination.

Tell your doctor if you are pregnant or breastfeeding.

This medicine is not approved for use by anyone younger than 18 years old.

How should I use albuterol and ipratropium inhalation?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Albuterol and ipratropium inhalation is usually used 4 times per day. Follow your doctor's dosing instructions very carefully. **Do not use more than 6 inhalations in a 24-hour period.**

Overuse of this medicine may increase the risk of death. It is critical that you use only the prescribed dose of this medicine.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions.

Always use the new inhaler device provided with your refill.

To use the inhaler:

- · You do not need to shake the inhaler before use.
- Uncap the mouthpiece of the inhaler. Breathe out fully. Put the mouthpiece into your mouth and close your lips. Keep
 your eyes closed to prevent spraying any medicine into your eyes. Breathe in slowly while pressing the dose-release
 button on the inhaler. Hold your breath for 10 seconds, then breathe out slowly.
- Close the cap until you use your inhaler again.
- · Carefully follow all directions for cleaning your specific inhaler device once per week.

Keep track of the number of sprays you have used. Throw away the **Combivent Respirat** inhaler canister after 3 months or 120 sprays, whichever comes first.

To use the solution with a nebulizer:

- Open the foil pouch and remove one vial. Empty the medicine into the chamber of the nebulizer. Attach the mouthpiece or face mask, then attach the drug chamber to the compressor.
- Sit upright in a comfortable position. Place the mouthpiece into your mouth or put on the face mask, covering your nose and mouth. Turn on the compressor.
- Breathe in slowly and evenly until no more mist is formed by the nebulizer and the drug chamber is empty.
- Clean the nebulizer after each use. Follow the cleaning directions that came with your nebulizer.

Seek medical attention if your breathing problems get worse quickly, or if you think your medications are not working as well.

To make sure this medicine is not causing harmful effects on your lungs, you may need to have chest X rays or other frequent lung function tests.

Store at room temperature away from moisture, heat, and light. Do not freeze. Keep each vial in its foil pouch until you are ready to use it.

Keep the cover on your inhaler when not in use. Keep away from open flame or high heat. The canister may explode if it gets too hot. Do not puncture or burn an empty inhaler canister.

Albuterol and ipratropium dosing information

Usual Adult Dose for Chronic Obstructive Pulmonary Disease -- Maintenance:

Inhalation Aerosol:

1 inhalation (albuterol-ipratropium bromide 100 mcg-20 mcg) orally four times a day. Additional inhalations can be taken as required.

Maximum dose: 6 inhalations in 24 hours

Inhalation solution:

One 3 mL vial (albuterol-ipratropium bromide 2.5 mg-0.5 mg) four times a day via nebulization

Maximum dose: Up to 2 additional 3 mL doses per day, if needed

Comments:

- -The inhalation solution can be continued as medically indicated to control recurring bouts of bronchospasm.
- -If a previously effective regimen fails to provide the usual relief, medical advice should be sought immediately, as this is often a sign of worsening COPD and may require reassessment of therapy.

Use: In patients with chronic obstructive pulmonary disease (COPD) on a regular aerosol bronchodilator who continue to have evidence of bronchospasm and who require a second bronchodilator

Detailed Albuterol / ipratropium dosage information

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

Get your prescription refilled before you run out of medicine completely.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of albuterol and ipratropium can be fatal.** Overdose symptoms may include chest pain, fast or pounding heartbeats, tremors, dry mouth, extreme thirst, muscle weakness or limp feeling, severe headache, pounding in your neck or ears, or feeling like you might pass out.

What should I avoid while using albuterol and ipratropium inhalation?

If this medication gets in your eyes, rinse with water and seek medical attention.

Avoid driving or hazardous activity until you know how albuterol and ipratropium will affect you. Your vision or reactions could be impaired.

What other drugs will affect albuterol and ipratropium inhalation?

Tell your doctor about all your other medicines, especially:

- a diuretic or "water pill";
- · heart or blood pressure medicine;
- other beta-blockers; or
- an antidepressant.

This list is not complete. Other drugs may affect albuterol and ipratropium, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Albuterol / ipratropium drug interactions (more detail)

Does albuterol/ipratropium interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Enter a drug name Add

More about albuterol / ipratropium

- Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (47)
- Side effects
- Dosage information
- During pregnancy
- Drug class: bronchodilator combinations
- En español

Patient resources

- · Ipratropium and albuterol advanced reading
- Ipratropium and Albuterol Inhalation Solution
- Ipratropium and Albuterol Inhalation Spray

Other brands

DuoNeb, Combivent, Combivent Respirat

Professional resources

• Ipratropium and Albuterol prescribing information

Other brands

DuoNeb, Combivent, Combivent Respimat

Related treatment guides

- COPD, Maintenance
- COPD

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ം Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

6.6 / 10

47 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.