

[Home](#)

2. Vagistat-1

Vagistat-1

Generic name: [tioconazole vaginal](#) [*tee-oh-CON-a-zole*]**Drug class:** [Vaginal anti-infectives](#)[Medically reviewed](#) by Drugs.com on Jun 12, 2024. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

What is Vagistat-1?

[Tioconazole](#) is an antifungal medication that fights infections caused by fungus.

Vagistat-1 (for use in the vagina) is used to treat vaginal Candida (yeast) infections.

Vagistat-1 may also be used for other purposes not listed in this medication guide.

Vagistat-1 side effects

Get emergency medical help if you have any of these **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have new or worsening vaginal symptoms.

Common side effects of Vagistat-1 may include:

- vaginal redness or swelling;
- mild stomach pain;
- pain or burning when you urinate;
- [headache](#), runny or stuffy nose; or
- mild vaginal burning, [itching](#) or irritation after use.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

[Vagistat-1 side effects](#) (more detail)

Related/similar drugs

Diflucan

Diflucan (fluconazole) is used to treat and prevent fungal infections. Includes Diflucan side ...

Reviews & ratings

5.9 / 10

[144 Reviews](#)

[View more](#)

Acidophilus

Acidophilus helps maintain an acidic environment, which can prevent the growth of harmful bacteria ...

Reviews & ratings

9.5 / 10

[29 Reviews](#)

[View more](#)

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

[685 Reviews](#)

[View more](#)

Warnings

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use Vagistat-1 if you are allergic to it.

Do not use Vagistat-1 if you have never had a [vaginal yeast infection](#) that has been confirmed by a doctor.

Ask a doctor or pharmacist if it is safe for you to use this medicine if you have other medical conditions, especially:

- fever, chills, vomiting, pain in your back or shoulder;
- pelvic pain, vaginal discharge with a bad odor;
- if you are having vaginal itching or discomfort for the first time; or
- if you think you may have been exposed to HIV (human immunodeficiency virus).

FDA pregnancy category C. It is not known whether Vagistat-1 will harm an unborn baby. **Do not use this medicine without your doctor's advice if you are pregnant.**

It is not known whether tioconazole vaginal passes into breast milk or if it could harm a nursing baby. **Do not use this medicine without your doctor's advice if you are breast-feeding.**

Do not give this medication to anyone under 12 years old without medical advice.

How should I use Vagistat-1?

Use exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended.

Do not take Vagistat-1 by mouth. It is for use only in your vagina.

This medication comes with patient instructions for safe and effective use. Follow these directions carefully. Ask your doctor or pharmacist if you have any questions.

This medicine is usually given as a single dose. You should notice some improvement in your symptoms after the 1-day treatment. It may take up to 7 days after using the 1-day dose before your symptoms go away completely.

Call your doctor if your symptoms do not improve after 3 days, or if you still have symptoms for longer than 7 days. Frequent vaginal yeast infections that do not clear up with treatment may be a sign of a more serious condition.

The single-use applicator is for one use only. Throw the applicator away after one use, even if there is still some medicine left in it after inserting your dose.

Vagistat-1 contains mineral oil or petrolatum, which can weaken the latex rubber in a condom or diaphragm.

These forms of birth control may be less effective if you use them during your treatment with this medicine. Use another form of birth control while you are using tioconazole and for at least 3 full days after your treatment ends.

Store at room temperature away from moisture and heat. Do not warm the medicine applicator before using Vagistat-1.

 [Detailed Vagistat-1 dosage information](#)

What happens if I miss a dose?

Since Vagistat-1 is used as a single dose, it does not have a daily dosing schedule.

What happens if I overdose?

Since Vagistat-1 is used as a single dose from a prefilled applicator, an overdose is unlikely to occur.

What should I avoid while using Vagistat-1?

Avoid wearing tight-fitting, synthetic clothing such as nylon underwear or panty hose that does not allow air circulation. Wear loose-fitting clothing made of cotton and other natural fibers until your [infection](#) is healed.

Do not use a tampon, vaginal douche, spermicide, or other vaginal products while you are using Vagistat-1. Do not have vaginal intercourse during treatment.

What other drugs will affect Vagistat-1?

It is not likely that other drugs you take orally or inject will have an effect on tioconazole used in the vagina. But many drugs can interact with each other. Tell each of your healthcare providers about all medicines you use, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#).

Does Vagistat-1 interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Vagistat-1

+

Enter a drug name

Add

Frequently asked questions

- [Can antibiotics cause yeast infections?](#)

More about Vagistat-1 (tioconazole topical)

- [Compare alternatives](#)
- [Reviews \(26\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: vaginal anti-infectives](#)
- [Breastfeeding](#)
- [En español](#)

Professional resources

- [Tioconazole monograph](#)

Related treatment guides

- [Vaginal Yeast Infection](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 6.04.

DRUG STATUS

Availability

OTC Over the counter

[Pregnancy & Lactation](#)


 Risk data available

CSA Schedule*

N/A Not a controlled drug



Approval History

 Drug history at FDA



User Reviews & Ratings

6.6 / 10

[26 Reviews](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.