

[Home](#)

2. Prevalite

# Prevalite

**Generic name:** [cholestyramine](#) [ *koe-le-STYE-ra-meen* ]**Brand names:** Prevalite, Prevalite Packets, [Questran](#), [Questran Light](#), Questran Packets, [Cholestyramine Light](#)**Drug class:** [Bile acid sequestrants](#)[Medically reviewed](#) by Drugs.com on Aug 3, 2023. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

## What is Prevalite?

Prevalite is used to lower high levels of cholesterol in the blood, especially low-density lipoprotein (LDL) ("bad" cholesterol).

Prevalite is also used to treat itching caused by a blockage in the bile ducts of the gallbladder.

Prevalite may also be used for purposes not listed in this medication guide.

## Prevalite side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

**Prevalite may cause serious side effects. Call your doctor at once if you have:**

- ongoing or worsening [constipation](#);
- severe stomach pain;
- blood in your urine;
- black, bloody, or tarry stools; or
- easy bruising, unusual bleeding.

Side effects such as constipation may be more likely in older adults.

**Common side effects of Prevalite may include:**

- mild constipation, [diarrhea](#);
- stomach pain, nausea, loss of appetite;
- bloating or gas;

- irritation of your tongue; or
- itching or irritation around your rectal area.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Prevalite side effects](#) (more detail)

## Related/similar drugs

### Lipitor

Lipitor is used to treat high cholesterol.

Learn about side effects, interactions and indications.

#### Reviews & ratings

4.3 / 10

[102 Reviews](#)

[View more](#)

### Crestor

Crestor (rosuvastatin) is used to treat high cholesterol and high triglycerides in the blood ...

#### Reviews & ratings

5.6 / 10

[138 Reviews](#)

[View more](#)

#### FEATURED

### Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

#### Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

### Zocor

Zocor (simvastatin) reduces low-density lipoprotein cholesterol and total

cholesterol in the blood ...

### Reviews & ratings

5.0 / 10

[22 Reviews](#)

[View more](#)

## Gemfibrozil

Gemfibrozil is used to treat high cholesterol and triglyceride levels in people with pancreatitis ...

### Reviews & ratings

6.1 / 10

[27 Reviews](#)

[View more](#)

## Pravastatin

Pravastatin (Pravachol) is a statin medicine used to lower cholesterol and triglycerides in the ...

### Reviews & ratings

4.5 / 10

[78 Reviews](#)

[View more](#)

## Fenofibrate

Fenofibrate is used to treat high cholesterol and high triglyceride levels. Learn about side ...

### Reviews & ratings

5.7 / 10

[69 Reviews](#)

[View more](#)

## Warnings

You should not use Prevalite if you have a blockage in your stomach or intestines.

## Before taking this medicine

You should not use Prevalite if you are allergic to it, or if you have:

- a blockage in your biliary tract.

Tell your doctor if you have ever had:

- chronic constipation;
- a thyroid disorder;
- diabetes;
- kidney disease;
- [liver disease](#); or
- [coronary artery disease](#) (clogged arteries).

This medicine may contain phenylalanine. Talk to your doctor before using Prevalite if you have [phenylketonuria](#) (PKU).

Taking Prevalite can make it harder for your body to absorb certain [vitamins](#). Your doctor may recommend you take a vitamin supplement.

Tell your doctor if you are pregnant or breastfeeding.

**You may have an even greater need for vitamin supplements during pregnancy or while you are breast-feeding a baby.** Follow your doctor's instructions about taking vitamin supplements during treatment with Prevalite.

## How should I take Prevalite?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Although Prevalite is usually taken once or twice per day, Prevalite may be taken up to 6 times per day. **Carefully follow your doctor's dosing instructions.**

Mix the Prevalite powder with at least 2 to 6 ounces of water or other non-carbonated beverage. You may also mix the powder with a brothy soup, crushed pineapple, or applesauce. Measure the powder using the scoop provided with your medication. Do not use any other scoop or measuring cup to measure your this medicine dose.

Use Prevalite regularly to get the most benefit. Get your prescription refilled before you run out of medicine completely.

**Prevalite can affect the surfaces of your teeth.** Sipping the this medicine/liquid mixture slowly or holding the liquid in your mouth for too long may result in tooth discoloration, enamel erosion, or tooth decay. Be sure to brush your teeth regularly while you are using this medicine.

Drink extra fluids to prevent constipation while you are taking Prevalite.

While using Prevalite, you may need frequent blood tests.

Prevalite is only part of a complete program of treatment that may also include diet, exercise, and weight control. Follow your diet, medication, and exercise routines very closely.

Store at room temperature away from moisture and heat.

 [Detailed Prevalite dosage information](#)

## What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe stomach pain or constipation.

## What should I avoid while taking Prevalite?

Avoid taking other medications at the same time you take Prevalite. Wait at least 1 hour before or 4 to 6 hours after taking this medicine before you take any other medications.

## What other drugs will affect Prevalite?

**Tell your doctor about all your current medicines.** Many drugs can affect Prevalite, especially:

- [birth control pills](#) or hormone replacement therapy;
- digoxin (digitalis);
- a diuretic or "water pill";
- penicillin G;
- [phenobarbital](#);
- [phenylbutazone](#);
- [propranolol](#);
- [spironolactone](#);
- [tetracycline](#);
- thyroid medication such as [levothyroxine](#); or
- [warfarin](#) ([Coumadin](#), [Jantoven](#)).

This list is not complete. Other drugs may interact with [cholestyramine](#), including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible interactions are listed in this medication guide.

 [Prevalite drug interactions](#) (more detail)

## Does Prevalite interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Prevalite

+

Enter a drug name

Add

## More about Prevalite (cholestyramine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(4\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: bile acid sequestrants](#)
- [Breastfeeding](#)

## Patient resources

- [Prevalite Packets drug information](#)
- [Prevalite \(Advanced Reading\)](#)

## Other brands

[Questran](#), [Cholestyramine Light](#), [Questran Light](#)

## Professional resources

- [Prevalite prescribing information](#)
- [Cholestyramine \(AHFS Monograph\)](#)

## Other brands

[Questran](#), [Cholestyramine Light](#)

## Related treatment guides

- [Hyperlipoproteinemia Type IIa, Elevated LDL](#)
- [Hyperlipoproteinemia Type IIb, Elevated LDL VLDL](#)
- [Pruritus of Partial Biliary Obstruction](#)
- [Hyperlipoproteinemia](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 9.01.

### DRUG STATUS

#### Availability

**Rx** Prescription only

#### [Pregnancy & Lactation](#)


 Risk data available

#### CSA Schedule\*

**N/A** Not a controlled drug



#### Approval History

 Drug history at FDA



## User Reviews & Ratings

[4 Reviews](#)

## Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



### About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

### Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

### Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.