

#### Home

Coagulation factor ix

# Coagulation factor IX

Generic name: coagulation factor IX [ koe-AG-yoo-LAY-shun-FAK-tor-NINE ]

Brand names: AlphaNine SD, Alprolix, BeneFIX, Idelvion, Ixinity, ... show all 10 brands

Dosage forms: intravenous kit (recombinant; recombinant albumin fusion protein), intravenous powder for injection (human; recombinant;

recombinant fc fusion protein; recombinant glycoPEGylated)

Drug class: Miscellaneous coagulation modifiers



Medically reviewed by Drugs.com on Aug 16, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

## What is coagulation factor IX?

Coagulation factor IX is used to treat or prevent bleeding in people with hemophilia B. This medication is not for treating people with hemophilia A factor VIII deficiency.

Coagulation factor IX may also be used for purposes not listed in this medication guide.

## Coagulation factor IX side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives, itching; wheezing, tightness in your chest, difficult breathing, fast heartbeats, blue lips, feeling like you might pass out; swelling of your face, lips, tongue, or throat.

#### Coagulation factor IX may cause serious side effects. Call your doctor at once if you have:

- weight gain, swelling in your waist, hands, or lower legs;
- · loss of appetite;
- fever or chills;
- continued bleeding after treatment;
- · new or worsened bleeding; or
- signs of excessive blood clotting--sudden numbness or weakness (especially on one side of the body), slurred speech, problems with vision or balance, chest pain, coughing up blood, or pain, swelling, warmth and redness in one or both legs.

#### Common side effects of coagulation factor IX may include:

- nausea;
- headache;

- · dizziness;
- · altered sense of taste;
- mild skin rash; or
- pain, redness, itching, stinging, or other irritation where the medicine was injected.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Coagulation factor ix side effects (more detail)

## Related/similar drugs

### Hemgenix

Hemgenix (etranacogene dezaparvovec-drlb) is a gene therapy indicated for the treatment of adults ...

### **Reviews & ratings**

Add a review

View more

## Lysteda

Lysteda (tranexamic acid) is used to treat heavy menstrual bleeding. Includes Lysteda side effects ...

### **Reviews & ratings**

7.9 / 10

70 Reviews

View more

FEATURED

### **Ozempic**

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

#### **Reviews & ratings**

6.6 / 10

1,504 Reviews

View more

#### **BeneFix**

BeneFix (coagulation factor IX) is a once-weekly, injectable, treatment for hemophilia B that may ...

### **Reviews & ratings**

1 Reviews

#### View more

## **Hympavzi**

Hympavzi (marstacimab-hncq) may be used to prevent or reduce the frequency of bleeding episodes ...

## **Reviews & ratings**

Add a review

#### View more

## **Alprolix**

Alprolix is used in hemophilia B to help control and prevent bleeding, it is a coagulation Factor ...

### **Reviews & ratings**

Add a review

### View more

### **Idelvion**

Idelvion (coagulation factor ix recombinant, albumin fusion protein) is a long-acting, injectable ...

### **Reviews & ratings**

1 Reviews

### View more

### **Beqvez**

Beqvez (fidanacogene elaparvovecdzkt) is an one-time, injectable, viral vector-based gene therapy ...

## **Reviews & ratings**

Add a review

### View more

## **Warnings**

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

## Before taking this medicine

You should not use coagulation factor IX if:

- you have had an allergic reaction to clotting factor medicine;
- · you have signs of excessive blood clotting; or
- you are allergic to hamster proteins.

Tell your doctor if you have ever had:

- · kidney disease;
- liver disease;
- coronary artery disease (hardening of the arteries); or
- a heart attack or stroke.

Tell your doctor if you are pregnant or breastfeeding.

Coagulation factor IX is made from donated human plasma and may contain viruses or other infectious agents. Donated plasma is tested and treated to reduce the risk of contamination, but there is still a small possibility it could transmit disease. Ask your doctor about any possible risk.

## How should I use coagulation factor IX?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

You may receive your first dose in a hospital or clinic setting to quickly treat any serious side effects.

Coagulation factor IX is given as an infusion into a vein. A healthcare provider will give your first dose and may teach you how to properly use the medication by yourself.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you don't understand all instructions.

Coagulation factor IX must be mixed with a liquid (diluent) before using it. When using injections by yourself, be sure you understand how to properly mix and store the medicine.

The mixture should appear clear, yellow, or colorless. Prepare an injection only when you are ready to give it. **Do not** use if the medicine looks cloudy, has changed colors, or has particles in it. Call your pharmacist for new medicine.

Gently swirl but do not shake the medicine while mixing or you may ruin the medicine. Use the injection within 3 hours after mixing your dose.

You may need frequent blood tests.

In case of emergency, wear or carry medical identification to let others know you have a bleeding disorder.

Carefully follow directions on your medicine label about how to store the powder medicine and the diluent. Some forms of coagulation factor IX can be stored at room temperature, and others should be kept in a refrigerator. Do not freeze. Avoid exposing the medication to light.

Each vial (bottle) is for one use only. Throw it away after one use, even if there is still medicine left inside.

Throw away any leftover coagulation factor IX once the expiration date on the label has passed.

Use a needle and syringe only once and then place them in a puncture-proof "sharps" container. Follow state or local laws about how to dispose of this container. Keep it out of the reach of children and pets.

Detailed Coagulation factor ix dosage information

## What happens if I miss a dose?

Call your doctor for instructions if you miss a dose.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

## What should I avoid while using coagulation factor IX?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

## What other drugs will affect coagulation factor IX?

Other drugs may affect coagulation factor IX, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

Coagulation factor ix drug interactions (more detail)

## Does coagulation factor ix interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

coagulation factor ix	
+	
Enter a drug name	Add

## More about coagulation factor ix

- · Check interactions
- Compare alternatives
- Reviews (5)
- · Side effects
- Dosage information
- During pregnancy
- Drug class: miscellaneous coagulation modifiers
- Breastfeeding
- En español

## **Patient resources**

- Coagulation factor ix recombinant advanced reading
- Coagulation factor ix recombinant, glycopegylated (Advanced Reading)
- Factor ix albumin fusion protein recombinant (Advanced Reading)
- Factor ix fc fusion protein recombinant (Advanced Reading)
- Factor IX (Recombinant)

### Other brands

BeneFix, Alprolix, Idelvion, Alphanine SD, ... +4 more

### **Professional resources**

- Factor IX (Human), Factor IX Complex (Human) monograph
- Factor IX (Recombinant) (AHFS Monograph)
- Factor IX (Recombinant), Fc fusion protein (AHFS Monograph)
- Factor IX (Recombinant), albumin fusion protein (AHFS Monograph)

### Other brands

BeneFix, Alprolix, Idelvion, Alphanine SD, ... +4 more

## Related treatment guides

- Factor IX Deficiency
- Hemophilia B

## **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 6.01.

### **DRUG STATUS**

### **Availability**

Rx Prescription only

**Pregnancy & Lactation** 

ஆ Risk data available

**CSA Schedule\*** 

N/A Not a controlled drug

## **User Reviews & Ratings**

5 Reviews

## **Images**

Alprolix (coagulation factor ix) multiple strengths

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





### **About**

About Drugs.com
Advertising policy

Attribution & citations

### Terms & privacy

Terms of use

**Editorial policy** 

Privacy policy

### Support

Help center

Sitemap

Contact us









Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.