

[Home](#)

2. Risperdal

Risperdal

Generic name: [risperidone](#) (oral) [*ris-PER-i-done*]**Drug class:** [Atypical antipsychotics](#)Medically reviewed by [Melisa Puckey, BPharm](#). Last updated on Mar 1, 2024.[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#) [FAQ](#)

What is Risperdal?

Risperdal is an [antipsychotic medicine](#) that works by changing the effects of chemicals in the brain.

Risperdal is used to treat [schizophrenia](#) in adults and children who are at least 13 years old.

Risperdal is also used to treat symptoms of [bipolar disorder](#) (manic depression) in adults and children who are at least 10 years old.

Risperdal is also used to treat symptoms of irritability in autistic children who are 5 to 16 years old.

Warnings

Risperdal is not approved for use in older adults with [dementia-related](#) psychosis.

Do not give Risperdal to a child without a doctor's advice.

While you are taking this medicine, you may be more sensitive to temperature extremes such as very hot or cold conditions. Avoid getting too cold, or becoming overheated or dehydrated. Drink plenty of fluids, especially in hot weather and during exercise. It is easier to become dangerously overheated and dehydrated while you are taking this medication.

Risperidone may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert. Drinking alcohol can increase certain side effects of risperidone.

Stop using Risperdal and call your doctor at once if you have fever, stiff muscles, confusion, sweating, fast or uneven heartbeats, restless muscle movements in your face or neck, tremor (uncontrolled shaking), trouble swallowing, feeling light-headed, or fainting.

Before taking this medicine

You should not use Risperdal if you are allergic to risperidone.

Risperidone may increase the risk of death in older adults with dementia-related psychosis and is not approved for this

use.

To make sure this medicine is safe for you, tell your doctor if you have ever had:

- heart disease, [high blood pressure](#), heart rhythm problems, stroke or [heart attack](#);
- diabetes (or risk factors such as [obesity](#) or family history of diabetes);
- low white blood cell (WBC) counts;
- liver or kidney disease;
- [seizures](#);
- [breast cancer](#);
- low bone mineral density;
- trouble swallowing;
- [Parkinson's disease](#); or
- if you are dehydrated.

The Risperdal M-Tab orally disintegrating tablet may contain phenylalanine. Tell your doctor if you have [phenylketonuria](#) (PKU).

Taking antipsychotic medication during the last 3 months of pregnancy may cause problems in the newborn, such as withdrawal symptoms, breathing problems, feeding problems, fussiness, tremors, and limp or stiff muscles. However, you may have withdrawal symptoms or other problems if you stop taking your medicine during pregnancy. **If you become pregnant while taking , do not stop taking it without your doctor's advice.**

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of risperidone on the baby.

Risperidone may temporarily affect fertility (ability to have children) in women.

Risperidone can pass into breast milk and may cause side effects in the baby. **If you breast-feed while using Risperdal, tell your doctor if the baby has symptoms such as drowsiness, tremors, or involuntary muscle movements.**

Do not give this medicine to a child without a doctor's advice.

 [Risperdal pregnancy and breastfeeding warnings](#) (more detail)

How should I take Risperdal?

Take Risperdal exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

Risperdal can be taken with or without food.

Remove the Risperdal M-Tab orally disintegrating tablet from the package only when you are ready to take the medicine. Place the tablet in your mouth and allow it to dissolve, without chewing. Swallow several times as the tablet dissolves.

Measure liquid medicine carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

Do not mix the Risperdal **liquid** with cola or tea.

It may take up to several weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

Store at room temperature away from moisture, heat, and light. Do not liquid medicine to freeze.

 [Detailed Risperdal dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

Get your prescription refilled before you run out of medicine completely.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe drowsiness, [fast heart rate](#), feeling light-headed, fainting, and restless muscle movements in your eyes, tongue, jaw, or neck.

What to avoid

Avoid driving or hazardous activity until you know how risperidone will affect you. Your reactions could be impaired.

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. [Dizziness](#) or severe drowsiness can cause falls, fractures, or other injuries.

Avoid drinking alcohol. Dangerous side effects could occur.

While you are taking risperidone, you may be more sensitive to temperature extremes such as very hot or cold conditions. Avoid getting too cold, or becoming overheated or dehydrated. Drink plenty of fluids, especially in hot weather and during exercise.

Risperdal side effects

Get emergency medical help if you have **signs of an allergic reaction to Risperdal:** [hives](#); difficulty breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement);
- breast swelling or tenderness (in men or women), nipple discharge, [impotence](#), lack of interest in sex, missed menstrual periods;
- **severe nervous system reaction** - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out;

- **low white blood cells** - sudden weakness or ill feeling, fever, chills, [sore throat](#), mouth sores, red or swollen gums, trouble swallowing, skin sores, cold or flu symptoms, cough, trouble breathing;
- **low levels of platelets in your blood** - easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin;
- **high blood sugar** - increased thirst, increased urination, dry mouth, fruity breath odor; or
- penis erection that is painful or lasts 4 hours or longer.

Common Risperdal side effects may include:

- [headache](#);
- dizziness, drowsiness, feeling tired;
- tremors, twitching or uncontrollable muscle movements;
- [agitation](#), [anxiety](#), restless feeling;
- depressed mood;
- dry mouth, [upset stomach](#), [diarrhea](#), [constipation](#);
- weight gain; or
- [cold symptoms](#) such as stuffy nose, sneezing, sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Risperdal side effects](#) (more detail)

Related/similar drugs

Lybalvi

Lybalvi is used to treat adults with manic or mixed episodes associated with bipolar I disorder ...

Reviews & ratings

8.1 / 10

[16 Reviews](#)

[View more](#)

Fanapt

Fanapt (iloperidone) is an antipsychotic medication used to treat schizophrenia. Includes Fanapt ...

Reviews & ratings

6.4 / 10

[30 Reviews](#)

[View more](#)

FEATURED

Caplyta

Caplyta (lumateperone) may be used to treat schizophrenia or depression associated with bipolar ...

Reviews & ratings

6.6 / 10

[184 Reviews](#)

[View more](#)

Venlafaxine

Venlafaxine is used to treat major depressive disorder, anxiety, and panic disorder. Learn about ...

Reviews & ratings

6.5 / 10

[2,786 Reviews](#)

[View more](#)

Aripiprazole

Easy-to-read patient leaflet for aripiprazole. Includes indications, proper use, special ...

Reviews & ratings

6.1 / 10

[1,540 Reviews](#)

[View more](#)

Quetiapine

Quetiapine is an atypical antipsychotic used to treat schizophrenia, bipolar disorder, and major ...

Reviews & ratings

7.2 / 10

[2,249 Reviews](#)

[View more](#)

Fluoxetine

Fluoxetine is used to treat depression or obsessive-compulsive disorder in adults. Learn about side ...

Reviews & ratings

7.3 / 10

[2,232 Reviews](#)

[View more](#)

Abilify

Abilify is an antipsychotic medicine used to treat the symptoms of schizophrenia and bipolar ...

Reviews & ratings

6.1 / 10

[1,187 Reviews](#)

[View more](#)

Seroquel

Seroquel is used to treat the symptoms of schizophrenia, bipolar disorder and major depressive ...

Reviews & ratings

7.0 / 10

[694 Reviews](#)

[View more](#)

What other drugs will affect Risperdal?

Taking Risperdal with other drugs that make you sleepy or slow your breathing can cause dangerous or life-threatening side effects. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your other medicines, especially:

- blood pressure medication;
- [carbamazepine](#);
- [clozapine](#);

- [fluoxetine \(Prozac\)](#) or [paroxetine \(Paxil\)](#); or
- [levodopa](#).

This list is not complete. Other drugs may interact with risperidone, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible drug interactions are listed here.

i [Risperdal drug interactions](#) (more detail)

Does Risperdal interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Risperdal

+

Enter a drug name

Add

Popular FAQ

What are the strongest sleeping pills?



More FAQ

- [What drugs cause tardive dyskinesia?](#)
- [What is the difference between Perseris and Risperdal Consta?](#)

[View more FAQ...](#)

More about Risperdal (risperidone)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(241\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)
- [Generic availability](#)
- [Support group](#)

- [FDA approval history](#)
- [Drug class: atypical antipsychotics](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

Other brands

[Uzedy](#), [Perseris](#), [Risvan](#), [Rykindo](#)

Professional resources

- [Risperdal prescribing information](#)
- [Risperidone \(AHFS Monograph\)](#)

Other brands

[Uzedy](#), [Perseris](#), [Risvan](#), [Rykindo](#)

Other formulations

- [Risperdal Consta](#)

Related treatment guides

- [Autism](#)
- [Schizoaffective Disorder](#)
- [Schizophrenia](#)
- [Bipolar Disorder](#)
- [Asperger Syndrome](#)
- [Mania](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Risperdal only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 21.03.

Availability

Rx Prescription only

Pregnancy & Lactation

 Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+ years FDA approved 1993



User Reviews & Ratings

6.0 / 10

[241 Reviews](#)

Images

[Risperdal 1 mg \(JANSSEN R 1\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)





[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.